



Page	Content
2	Surveillance programme description
3	Comments
4-6	Systematic Influenza-like illness (ILI) surveillance Influenza Respiratory syncytial virus <i>Bordetella pertussis</i>
7	Influenza-like illness (ILI) Viral Watch Influenza
8-10	National syndromic surveillance for pneumonia Influenza Respiratory syncytial virus <i>Bordetella pertussis</i>
11	Private Hospital Consultation Outpatient consultations In patient consultations

Respiratory Pathogen Surveillance

Programme Descriptions

Programme	Influenza-like illness (ILI)	Viral Watch	National syndromic surveillance for pneumonia	Private hospital consultations
Start year	2012	1984	2009	2002
Provinces*	KZ NW WC**	EC FS GP LP MP NC NW WC	GP KZ MP NW WC	EC FS GP LP MP NW WC
Type of site	Primary health care clinics	General practitioners	Public hospitals	Private hospitals
Case definition	An acute respiratory illness with a temperature ($\geq 38^{\circ}\text{C}$) and cough, & onset ≤ 10 days	An acute respiratory illness with a temperature ($\geq 38^{\circ}\text{C}$) and cough, & onset ≤ 10 days	Acute or chronic lower respiratory tract infection	ICD codes J10-J18
Specimens collected	Oropharyngeal & nasopharyngeal swabs	Throat and/or nasal swabs or Nasopharyngeal swabs	Oropharyngeal & nasopharyngeal swabs	Not applicable
Main pathogens tested***	INF RSV BP	INF RSV BP	INF RSV BP	Not applicable

Epidemic Threshold

Thresholds are calculated using the Moving Epidemic Method (MEM), a sequential analysis using the R Language, available from: <http://CRAN.R-project.org/web/package=mem>) designed to calculate the duration, start and end of the annual influenza epidemic. MEM uses the 40th, 90th and 97.5th percentiles established from available years of historical data to calculate thresholds of activity. Thresholds of activity for influenza and RSV are defined as follows: Below seasonal threshold, Low activity, Moderate activity, High activity, Very high activity. For influenza, thresholds from outpatient influenza like illness (Viral Watch Programme) are used as an indicator of disease transmission in the community and thresholds from pneumonia surveillance are used as an indicator of impact of disease.

* EC: Eastern Cape; FS: Free State; GP: Gauteng; KZ: KwaZulu-Natal; LP: Limpopo; MP: Mpumalanga; NC: Northern Cape; NW: North West; WC: Western Cape

**Started in 2019

***INF: influenza virus; RSV: respiratory syncytial virus; BP: *Bordetella pertussis*

Respiratory Pathogen Surveillance

Reporting period 01/01/2019 to 06/10/2019

Results until end of epidemiologic week 40 (2019)

Comments:

Influenza

The 2019 season started in week 16 (week ending 21 April) when influenza detection in the Viral Watch programme rose above the seasonal threshold, as determined by the Moving Epidemic Method. The season ended in week 33 (week ending 18 August) after which transmission dropped below seasonal threshold levels.

ILI programme: In 2019 to date, specimens from 1414 patients were received from 4 ILI sites. Influenza was detected in 152 specimens, 23 (15%) were identified as influenza A(H1N1)pdm09, 126 (83%) as influenza A(H3N2) and three (2%) A subtype inconclusive.

Viral Watch programme: During the same period, specimens were received from 1326 patients from Viral Watch sites in 7 provinces. Influenza was detected in 779 patients, of which 48 (6%) were influenza A(H1N1)pdm09, 718 (92%) influenza A(H3N2), 15 (2%) A subtype inconclusive, one (<1%) influenza B(Victoria) and one (<1%) influenza B(Yamagata). The above breakdown by type and subtype includes four patients testing dual positive for influenza: three were dual positive for influenza A(H1N1)pdm09 and A(H3N2) in week 25 and one dual positive for influenza A(H3N2) and influenza B(Yamagata) in week 27.

Pneumonia surveillance: Since the beginning of 2019, specimens from 3525 patients with severe respiratory illness (SRI) were received from the 6 sentinel sites. Influenza was detected in 220 patients, of which 15 (7%) were influenza A(H1N1)pdm09, 196 (89%) influenza A(H3N2), seven (3%) A subtype inconclusive, one (<1%) influenza B(Victoria) and two (1%) influenza B(Yamagata). This includes one dual positive patient for influenza A(H1N1)pdm09 and A(H3N2) in week 21.

Respiratory syncytial virus

The 2019 RSV season which started in week 8 (week starting 18 February) when RSV detections in pneumonia surveillance rose above the seasonal threshold as determined by the Moving Epidemic Method ended in week 25 (week starting 17 June). However, sporadic detections of RSV are still being made.

In 2019 to date, 1414 specimens were tested and RSV was detected in specimens of 144 (10%) patients in ILI programme.

Pneumonia surveillance: 3525 specimens were tested and RSV was detected in specimens of 775 (22%) patients.

Viral Watch programme: 1326 specimens were tested and RSV was detected in specimens of 30 (2%) patients.

Bordetella pertussis

ILI programme: From 1 January 2019 to date, combined nasopharyngeal and oropharyngeal specimens were tested from 1408 patients for *B. pertussis*, nine (0.6%) tested positive.

Pneumonia surveillance: During the same period, combined nasopharyngeal and oropharyngeal specimens were tested from 3504 patients for *B. pertussis*, which was detected in 31 (1.0%) specimens.

In addition, *B. pertussis* was detected in two of 137 (1.5 %) specimens from patients who did not meet the pneumonia/ILI surveillance case definition, but who did meet the suspected pertussis case definition.


Respiratory Pathogen Surveillance

Reporting period 01/01/2019 to 06/10/2019

Results until end of epidemiologic week 40 (2019)

Influenza-like illness (ILI) surveillance primary health care clinics

Figure 1. Number of positive samples* by influenza subtype and lineage and detection rate by week**



*Specimens from patients with influenza-like illnesses at 3 sentinel sites in 3 provinces

**Only reported for weeks with >10 specimens submitted

Inconclusive: insufficient viral load in sample and unable to characterise further

Table 1. Cumulative number of influenza subtype and lineage and total number of samples tested by clinic and province

Clinic (Province)	A(H1N1)pdm09	A(H3N2)	A subtype inconclusive	B/Victoria	B/Yamagata	B lineage inconclusive	Total samples
Eastridge (WC)	16	55	0	0	0	0	755
Edendale Gateway (KZ)	3	22	3	0	0	0	109
Jouberton (NW)	2	44	0	0	0	0	460
Mitchell's Plain (WC)	2	5	0	0	0	0	90
Total:	23	126	3	0	0	0	1414

KZ: KwaZulu-Natal; NW: North West; WCP: Western Cape

Inconclusive: insufficient viral load in sample and unable to characterise further


Respiratory Pathogen Surveillance

Reporting period 01/01/2019 to 06/10/2019

Results until end of epidemiologic week 40 (2019)

Influenza-like illness (ILI) surveillance primary health care clinics

Figure 2. Number of samples testing positive for respiratory syncytial virus by subgroup and detection rate by week



Inconclusive: insufficient viral load in sample and unable to characterise further
 RSV AB: Both RSV A and B subgroup identified

Table 2. Cumulative number of respiratory syncytial virus subgroups identified and total number of samples tested by clinic and province

Clinic (Province)	RSVA	RSVB	RSVAB	RSV subgroup inconclusive	Total samples
Eastridge (WC)	28	65	2	0	755
Edendale Gateway (KZ)	5	0	0	0	109
Jouberton (NW)	38	0	1	0	460
Mitchell's Plain (WC)	0	5	0	0	90
Total	71	70	3	0	1414

KZ: KwaZulu-Natal; NW: North West; WC: Western Cape
 Inconclusive: insufficient viral load in sample and unable to characterise further
 RSV AB: Both RSV A and B subgroup identified

Reporting period 01/01/2019 to 06/10/2019

Results until end of epidemiologic week 40 (2019)

Influenza-like illness (ILI) surveillance primary health care clinics

Figure 3. Number of samples testing positive for *B. pertussis* and detection rate by month




Table 3. Cumulative number of *B. pertussis* identified and total number of samples tested by province**

Clinic (Province)	<i>B. pertussis</i> Positive**	Total samples
Eastridge (WC)	4	754
Edendale Gateway (KZ)	2	107
Jouberton (NW)	3	458
Mitchell's Plain (WC)	0	89
Total:	9	1408

KZ: KwaZulu-Natal; NW: North West; WC: Western Cape

**26 cases met the suspected pertussis case definition but did not meet Influenza-like illness (ILI) case definition. These are not included in the table or the epidemiological curve


Respiratory Pathogen Surveillance

Reporting period 01/01/2019 to 06/10/2019

Results until end of epidemiologic week 40 (2019)

Influenza-like illness (ILI) surveillance Viral Watch

Figure 4. Number of positive samples* by influenza subtype and lineage and detection rate by week**



*Specimens from patients with Influenza-like illnesses at 90 sentinel sites in 7 provinces

** Only reported for weeks with >10 specimens submitted.

Inconclusive: insufficient viral load in sample and unable to characterise further
Dual positives are included in the graph cumulatively

Table 4. Cumulative number of influenza subtype and lineage and total number of samples tested by province

Province	A(H1N1)pdm09	A(H3N2)	A subtype inconclusive	B/Victoria	B/Yamagata	B lineage inconclusive	Total samples
Eastern Cape	2	48	3	0	0	0	67
Free State	1	54	0	0	0	0	79
Gauteng	14	374	8	0	1	0	675
Limpopo	2	31	0	0	0	0	48
Mpumalanga	5	28	1	0	0	0	77
North West	0	4	0	0	0	0	8
Western Cape	24	179	3	1	0	0	372
Total:	48	718	15	1	1	0	1326

*Inconclusive: insufficient viral load in sample and unable to characterise further

Included in the table are 3 dual specimens positive for influenza A(H1N1)pdm01 & influenza A(H3N2) in week25 and one dual positive for influenza A(H3N2) & influenza B(Yamagata) in week27

From 01 January 2019 to date, 36 patients were tested for influenza at the time of entry into South Africa following travel abroad and influenza was detected in five patients, of which one influenza A(H1N1)pdm09, two influenza A(H3N2), one influenza A (subtype inconclusive) and one influenza B (lineage inconclusive).

Patients known to have acquired influenza abroad are not included in the table or epidemiological curve.


Respiratory Pathogen Surveillance

Reporting period 01/01/2019 to 06/10/2019

Results until end of epidemiologic week 40 (2019)

National syndromic surveillance for pneumonia

Figure 5. Number of positive samples* by influenza subtype and lineage and detection rate by week**



*Specimens from patients hospitalised with pneumonia at 6 sentinel sites in 5 provinces

**Only reported for weeks with >10 specimens submitted

Inconclusive: insufficient viral load in sample and unable to characterise further

Dual positives are included in the graph cumulatively

Table 5. Cumulative number of identified influenza subtype and lineage and total number of samples tested by hospital

Hospital (Province)	A(H1N1)pdm09	A(H3N2)	A subtype inconclusive	B/Victoria	B/Yamagata	B lineage inconclusive	Total samples
Edendale (KZ)	7	32	2	0	0	0	581
Helen Joseph-Rahima Moosa (GP)	0	38	2	1	0	0	821
Klerksdorp-Tshepong (NW)	1	50	0	0	0	0	512
Mapulaneng-Matikwana (MP)	2	29	0	0	1	0	362
Red Cross (WC)	3	32	1	0	1	0	935
Mitchell's Plain (WC)	2	15	2	0	0	0	314
Total:	15	196	7	1	2	0	3525

GP: Gauteng; KZ: KwaZulu-Natal; NW: North West; MP: Mpumalanga; WC: Western Cape

Inconclusive: insufficient viral load in sample and unable to characterise further

From the influenza positive specimens reflected on the above table we detected one dual specimen positive for influenza A(H1N1)pdm01 & influenza A(H3N2) in week21


Respiratory Pathogen Surveillance

Reporting period 01/01/2019 to 06/10/2019

Results until end of epidemiologic week 40 (2019)

National syndromic surveillance for pneumonia

Figure 6. Number of samples testing positive for respiratory syncytial virus by subgroup and detection rate by week



Inconclusive: insufficient viral load in sample and unable to characterise further
 RSV AB: Both RSV A and B subgroup identified

Table 6. Cumulative number of respiratory syncytial virus subgroups identified and total number of samples tested by hospital

Hospital (Province)	RSVA	RSVB	RSVAB	RSV subgroup inconclusive	Total samples
Edendale (KZ)	123	5	0	1	581
Helen Joseph-Rahima Moosa (GP)	109	61	2	0	821
Klerksdorp-Tshepong (NW)	51	8	0	1	512
Mapulaneng-Matikwana (MP)	54	1	0	0	362
Red Cross (WC)	69	215	4	3	935
Mitchell's Plain (WC)	21	47	0	0	314
Total:	427	337	6	5	3525

GP: Gauteng; KZ: KwaZulu-Natal; NW: North West; MP: Mpumalanga; WC: Western Cape
 Inconclusive: insufficient viral load in sample and unable to characterise further
 RSV AB: Both RSV A and B subgroup identified

Respiratory Pathogen Surveillance

Reporting period 01/01/2019 to 06/10/2019

Results until end of epidemiologic week 40 (2019)

National syndromic surveillance for pneumonia

Figure 7. Number of samples testing positive for *B. pertussis* and detection rate by month




Table 7. Cumulative number of *B. pertussis* identified and total number of samples tested by hospital and province


Hospital (Province)	<i>B. pertussis</i> Positive**	Total samples
Edendale (KZ)	5	573
Helen Joseph-Rahima Moosa (GP)	8	814
Klerksdorp-Tshepong (NW)	3	511
Mapulaneng-Matikwana (MP)	4	361
Red Cross (WC)	11	932
Mitchell's Plain (WC)	0	313
Total:	31	3504

GP: Gauteng; KZ: KwaZulu-Natal; NW: North West; MP: Mpumalanga; WC: Western Cape

**111 cases met the suspected pertussis case definition but did not meet Pneumonia Surveillance case definition. These are not included in the table and epidemiologic curve.

Private hospital consultations


Figure 8. Number of private hospital outpatient consultations* with a diagnosis of pneumonia and influenza (P&I) and viral isolates**



* Hospital outpatient data from weekly reports of consultations to the Netcare hospital group. Discharge diagnosis is according to International Statistical Classification of Diseases and Related Health Problems coding by clinicians and does not represent laboratory confirmation of aetiology

** Influenza positive specimens from the Viral Watch surveillance programme

Figure 9. Number of private hospital admissions* with a discharge diagnosis of pneumonia and influenza (P&I) and viral isolates**



*Hospitalisation admission data from weekly reports of consultations to the Netcare hospital group. Discharge diagnosis is according to International Statistical Classification of diseases and Related Health Problems/ ICD by clinicians and does not represent laboratory confirmation of aetiology ** Influenza positive specimens from the national syndromic surveillance for pneumonia.