

Clinical management of suspected or confirmed COVID-19 disease

– version 3 (27th March 2020)

1. BACKGROUND	3
2. EPIDEMIOLOGY AND CLINICAL CHARACTERISTICS	3
3. MANAGEMENT OF SUSPECTED COVID-19 CASES	5
4. MANAGEMENT OF CONFIRMED COVID-19 CASES	9
5. INFECTION PREVENTION AND CONTROL (IPC)	15
6. RECORDING AND REPORTING	16
APPENDIX 1 – EXAMPLE OF A PATIENT INFORMATION SHEET	19

Writing committee (in alphabetical order): David Anderson, Lesley Bamford, Tom Boyles, Lucille Blumberg, Cheryl Cohen, Andrew Gray, Ahmad Haeri Mazanderani, Tendesayi Kufa-Chakeza, Halima Dawood, Fikile Mabena, Shaheen Mehtar, Natalie Mayet, Marc Mendelson, Jeremy Nel, Wolfgang Preiser, Jantjie Taljaard

Version 3: What's New?

- New de-isolation criteria for confirmed cases (section 4.5)
- Statement on therapeutics for hospitalized cases (section 4.3)
- Statement on the use of ACE-inhibitors, angiotensin receptor blockers, and nonsteroidal anti-inflammatory drugs (section 4.3).

Guideline Summary

Testing

- Apply the latest case definition from the NICD to determine testing eligibility.
<http://www.nicd.ac.za/diseases-a-z-index/covid-19/>
- Combined nasopharyngeal and oropharyngeal swabs should be sent in all suspected cases. Lower respiratory tract samples (e.g. sputum) can also be sent if present (do not perform sputum induction however).
- Ensure that the specimen is labelled and packaged correctly, and stays between 2-8°C during specimen storage and transport.

Suspected COVID-19 cases

- Any suspected case should be identified as soon as possible (ideally prior to entering the facility). Such cases should immediately be given a surgical mask, and be isolated. Good hand hygiene and cough etiquette should be taught, and appropriate samples obtained.
- A broad differential diagnosis should be entertained for suspected cases, and appropriate testing for alternative diagnoses should be undertaken.
- Suspected COVID-19 cases who are medically well, or who are assessed as having only mild disease, may be managed at home while awaiting test results.

Confirmed COVID-19 cases

- Patients with mild disease may be considered for management at home, provided they are able to safely self-isolate and are not at risk of developing severe disease (see criteria in table 2).
- Supportive oxygen therapy is the cornerstone of management for patients with severe disease – target oxygen saturations of $\geq 90\%$ for most patients, using nasal prong oxygen, a simple face mask, or a face mask with a reservoir bag.
- For intubated patients with ARDS, use lung-protective ventilation strategies.
- There is currently no good evidence for any specific therapy for COVID-19. Any investigational drugs or therapeutics should be reserved for hospitalized patients. It should ideally be administered as part of a trial, but at a minimum it should be administered under the Monitored Emergency Use of Unregistered Interventions (MEURI) framework.
- Patients may be de-isolated without the need for repeat PCR tests. Those with mild disease may be de-isolated 14 days after symptom onset, while those with severe disease may be de-isolated 14 days after achieving clinical stability (e.g. once supplemental oxygen is discontinued).
- There is currently no good evidence to suggest that patients on ACE-inhibitors, angiotensin-receptor blockers need to discontinue these agents.

Healthcare worker personal protective equipment (PPE)

- For the majority of direct COVID-19 patient interactions, appropriate healthcare worker person protective equipment consists of gloves, a gown or apron, and a surgical mask.
- When performing aerosol-generating procedures (e.g. taking nasopharyngeal swabs, performing CPR, or intubating a patient), an N95 respirator should be used in place of a surgical mask, and eye protection (shield or goggles) should be added to the above.

Clinical management of suspected or confirmed COVID-19 disease

1. Background

On 31st December 2019, the World Health Organization (WHO) was alerted to a cluster of pneumonia of unknown aetiology in patients in Wuhan City, Hubei Province of China. One week later the novel coronavirus (severe acute respiratory syndrome coronavirus 2: SARS-CoV-2) was identified as the cause. The resulting illness was named COVID-19 on the 11th February 2020. The clinical spectrum of COVID-19 ranges from an asymptomatic or mild flu-like illness to a severe pneumonia requiring critical care. These guidelines describe the clinical management of cases of COVID-19 disease and covers clinical care in and outside health care facilities. It is intended for health care practitioners taking care of symptomatic patients with suspected or confirmed COVID-19.

2. Epidemiology and clinical characteristics

SARS-CoV-2 is a betacoronavirus closely related to SARS-CoV and MERS-CoV. It is an enveloped, non-segmented, positive sense RNA virus. It is thought to have originated in bats but the animal that mediated transmission to humans remains unknown.

2.1 Epidemiology

The median incubation period for COVID-19 is estimated to be 4-5 days, with an interquartile range of 2-7 days.^{3,4} Transmission from asymptomatic patients has been postulated, but the extent of this is unknown.⁵ The reproductive number for the virus is approximately 2.2 (meaning that on average each person spread the infection to two others).³ In the early reported cases, the median age of reported cases was 50 years with a male preponderance of cases (~60%). Very few severe cases which required hospitalisation have been reported among children under the age of 15 years (~1%), although school closures may have influenced this figure. Risk factors for severe disease include older age and cardiopulmonary comorbidities.

2.2 Clinical characteristics – what to look for

80% of symptomatic patients develop mild disease, an estimated 15% develop severe disease (with hypoxaemia, dyspnoea and tachypnoea) while 5% become critically ill (with respiratory failure, septic shock and/or multiorgan dysfunction).⁶ The proportion of asymptomatic carriers is currently unknown.

The most common presenting symptom has been fever (~90%, but only present in 44% on admission). Other common symptoms include cough (68%), fatigue (38%), sputum production (34%), shortness of

breath (19%), myalgia or arthralgia (15%), sore throat (14%), headache (13.6%) and chills (12%).⁴ Gastrointestinal symptoms such as nausea or vomiting (5.0%) and diarrhoea (3.8%) appear to be uncommon.

Abnormalities are visible on chest X-ray in approximately 60% of COVID-19 patients, and on 85% of patients' chest CT scans.⁴ These are typically patchy ground glass opacities, though other patterns have been described.⁴

2.3 Outcomes and prognosis

The vast majority of cases will make a full recovery, though this may take several weeks, particularly in severe cases. In a minority of cases, COVID-19 has been associated with rapid progression to acute respiratory distress syndrome (ARDS), multiple organ failure and sometimes death. The case fatality ratio is currently unknown, but is estimated to be within the range of 0.5-4%.

3. Management of Suspected COVID-19 Cases

3.1 Early identification/triage

Patients fulfilling the latest case definition for suspected COVID-19 case (a “person under investigation”, PUI) should ideally phone ahead of time to their doctor or emergency room, so that adequate precautions can be taken ahead of time. PUIs who do not self-identify should be screened and identified as soon as possible upon entering a health facility, to avoid prolonged contact with other patients and healthcare workers.

- The criteria for a “person under investigation” (PUI) are dynamic and will change with time. For the latest criteria, please see the NICD’s website: <http://www.nicd.ac.za/diseases-a-z-index/covid-19/>

As of 26th March 2020, the NICD’s case definition is:

A hospitalized patient with severe acute respiratory illness (fever and at least one sign/symptom of respiratory disease, e.g. cough, shortness of breath) AND the absence of an alternative diagnosis that fully explains the clinical presentation

OR

Any person with acute respiratory illness with sudden onset of at least one of the following: cough, sore throat, shortness of breath or fever [$\geq 38^{\circ}\text{C}$ (measured) or history of fever (subjective)] irrespective of admission status **AND** in the 14 days prior to onset of symptoms, met at least one of the following epidemiological criteria:

- Were in close contact with a confirmed or probable case of SARS-CoV-2 infection;

OR

- Had a history of travel to areas with local transmission of SARS-CoV-2 (the list of these countries will change with time – consult the NICD website)

OR

- Worked in or attended a health care facility where patients with SARS-CoV-2 infections were being treated

- Measures that may facilitate early identification of suspected COVID-19 cases include:
 - Posters, pamphlets, billboards or staff members outside and within the healthcare facility asking patients who fulfil criteria for a PUI to identify themselves to healthcare workers as soon as possible (rather than remaining in line in a waiting area).
 - Including a screening questionnaire for COVID-19 as part of the standard triage form at healthcare facilities.
- Any patient who fulfils criteria for a suspected COVID-19 case should immediately have the following measures taken:
 - Give the patient a medical (surgical) mask (N95 respirators are NOT required for patients).

- Direct the patient to a separate area, preferably an isolation room if available. Where an individual isolation room is not available, a 1-2 metre distance should be kept between suspected COVID-19 cases and other patients.
- Instruct the patient to cover his/her nose and mouth during coughing or sneezing with a tissue or a flexed elbow. The patient should perform hand hygiene after contact with respiratory secretions (wash hands or use alcohol-based hand rub, which should be readily available at the point of triage).
- Limit the movement of the patient (e.g. use portable X-rays rather than sending the patient to the X-ray department). If the patient has to be moved, ensure that (s)he wears a mask.
- The patient should have a dedicated bathroom (where this is possible).
- Patients should be quickly triaged in terms of clinical severity. Routine emergency department triage systems may be used. In the context of COVID-19, triaging is essential because:
 - It allows for rapid initiation of supportive therapy (e.g. oxygen supplementation)
 - It has implications for whether or not the patient can be allowed home to await results of the COVID-19 testing (see below).
 - It protects both patients and staff.

3.2 Testing

All persons under investigation require testing for SARS-CoV-2 by means of reverse transcriptase PCR (RT-PCR). Samples to be sent are:

- *Upper respiratory tract samples* – nasopharyngeal and oropharyngeal swabs (combined in the same universal transport medium tube) in all patients.
- *Lower respiratory tract samples* – may not be possible depending on the patient's symptoms. Where available, send sputum, tracheal aspirates, or bronchoalveolar lavage fluid. Sputum induction should not be performed.

Appropriate personal protective equipment (PPE) should be worn by all healthcare workers when obtaining specimens (see IPC section below).

The **differential diagnosis** of suspected cases includes influenza (remembering the seasonality in patients from the northern hemisphere differs from those of the southern hemisphere), both conventional and atypical bacterial pneumonias, and in patients with HIV and a CD4 count <200 cells/mm³ (or equivalent immunosuppression), *Pneumocystis jirovecii* pneumonia. Depending on the patient, appropriate samples may include:

- Full blood count + differential count
- Blood culture
- Nasopharyngeal swabs or aspirates and oropharyngeal swabs for detection of viral and atypical pathogens
- Chest radiography
- Sputum for MCS and *Mycobacterium tuberculosis* detection (GeneXpert MTB/RIF Ultra).
- Urine for lipoarabinomannan (LAM) test if HIV positive

Obtaining samples for SARS-CoV-2 testing

- Healthcare workers obtaining respiratory samples require appropriate personal protective equipment, including those for contact, droplet and aerosol precautions (see infection prevention and control section below).
- Collecting a good quality specimen is vital. For details on how to properly obtain good quality specimens, please see Appendix 5 of the NICD's COVID-19 guidelines for case-finding, diagnosis, management and public health response in South Africa: <http://www.nicd.ac.za/wp-content/uploads/2020/02/Guidelines-for-case-finding-diagnosis-management-and-public-health-response-in-South-Africa.pdf>

Transport of specimens

- Ensure that samples are kept between 2-8°C until they are processed.

3.3 Empiric treatment of other pathogens

Where the patient fits the appropriate clinical syndrome, consider treatment of other pathogens such as:

- Conventional community-acquired pneumonia pathogens (or hospital-acquired pneumonia pathogens if appropriate) – e.g. ceftriaxone [see [EDL guidelines](#)]
- Atypical pneumonia pathogens – e.g. azithromycin [see [EDL guidelines](#)]
- Influenza (if influenza epidemiology fits and has severe illness or if patient is at risk of severe influenza) – oseltamivir [see [NICD influenza guidelines](#)]
- PJP (if appropriate risk factors present, e.g. HIV with low CD4 count)

3.4 Managing patients at home while awaiting COVID-19 test results

Suspected COVID-19 cases who are medically well, or who are assessed as having only mild disease, may be managed at home while awaiting test results.

Table 1 – Criteria for “mild” disease (for age >12 years)¹

Criteria for "mild" disease
<ul style="list-style-type: none"> • SpO₂ ≥95% • Respiratory rate <25 • Heart rate <120 • Temp 36-39°C • Mental status normal

¹For age 5-12, use respiratory rate <30, and heart rate <130. For younger ages, use age-appropriate normal values.

Such patients should be instructed to self-isolate at home and be given appropriate advice about reducing possible transmission to others:

- Patients should stay in a specific room and use their own bathroom (if possible). Patients should avoid unnecessary travel and unnecessary contact with other people.
- Where contact is unavoidable, the patient should wear a facemask, and maintain a distance of at least 1 metre (preferably 2 metres) from other people
- Patients should clean their hands with soap and water frequently. Alcohol-based sanitizers may also be used, provided they contain at least 70% alcohol.

- Patients should practice good cough and sneeze hygiene, by using a tissue, and then immediately discarding the tissue in a lined trash can, followed by washing hands immediately.
- Patients should not have visitors in their home. Only those who live in their home should be allowed to stay.
- At home, the patient should stay in a specific room and use his/her own bathroom (if possible). If they live in shared accommodation (university halls of residence or similar) with a communal kitchen, bathroom(s) and living area, they should stay in their room with the door closed, only coming out when necessary, wearing a facemask if they do so.
- Patients should avoid sharing household items like dishes, cups, eating utensils and towels. After using any of these, the items should be thoroughly washed with soap and hot water.
- All high-touch surfaces like table tops, counters, toilets, phones, computers, etc. should be appropriately and frequently cleaned.
- If patients need to wash laundry at home before the results are available, then they should wash all laundry at the highest temperature compatible for the fabric using laundry detergent. This should be above 60° C. If possible, they should tumble dry and iron using the highest setting compatible with the fabric. Disposable gloves and a plastic apron should be used when handling soiled materials if possible and all surfaces and the area around the washing machine should be cleaned. Laundry should not be taken to a laundrette. The patient should wash his/her hands thoroughly with soap and water after handling dirty laundry (remove gloves first if used).
- Patients should know who to call if they develop any worsening symptoms, so that they can be safely reassessed.
- In addition to being given the above advice, a patient information sheet can be given if possible (see Appendix 1 for an example).

See also the NICD's page on self-isolation advice, available at: <https://www.nicd.ac.za/what-to-do-if-i-test-positive-for-coronavirus-disease-and-i-am-asked-to-home-isolate%E2%80%8B/>

4. Management of Confirmed COVID-19 Cases

The goal in clinical management of cases is to reduce morbidity and mortality and minimise transmission to uninfected contacts. Triaging patients and early identification of patients who are severely or critically ill and require hospital or ICU admission will be essential in reducing morbidity and mortality while isolation and implementation of infection prevention and control (IPC) measures within facilities as well as contact tracing, education on good cough hygiene and IPC at home will help minimise onward transmission of the virus. Key management principles include:

4.1 Rapid triage of cases – in order that appropriate IPC measures and an appropriate level of supportive care can be commenced.

- Cases triaged as having moderate or severe disease will require admission for medical reasons.
- Patients with mild disease may be considered for management at home, provided they are able to safely self-isolate and are not at risk of developing severe disease (see criteria in table 2).
- If patients are to be managed at home, it is imperative that all appropriate measures are taken to prevent onward transmission of the disease to others - give advice as in section 3.1 above.
- Note also that in 10-15% of cases, those patients assessed as having “mild” disease may continue to worsen over the course of a week or more and become severely ill. **Patients managed from home need to be given the contact details of their doctor or healthcare facility that they can reach out to in case of any clinical deterioration.**

Table 2 - Criteria for management at home (for age >12 years¹):




Mild disease
<ul style="list-style-type: none">• SpO₂ ≥95%• Respiratory rate <25• HR <120• Temp 36-39°C• Mental status normal
Able to safely self-isolate
<ul style="list-style-type: none">• Separate bedroom available for patient to self-isolate in• Patient able to contact, and return to, healthcare facility in case of deterioration
Not at high risk of deterioration
<ul style="list-style-type: none">• Age <65 years• No severe cardiac or pulmonary comorbidities• No other debilitating comorbidities (e.g. cancer)

¹For age 5-12, use respiratory rate <30, and heart rate <130. For younger ages, use age-appropriate normal values.

4.2 Early supportive therapy in hospitalised COVID-19 patients

✓ Give supplemental oxygen therapy immediately to patients with low oxygen saturation.⁷

- Oxygen therapy is likely to be the single most effective supportive measure in COVID-19 patients overall. Target SpO₂ ≥90% in non-pregnant adults and SpO₂ ≥92% in pregnant patients.⁷ Children with emergency signs (obstructed or absent breathing, severe respiratory distress, central cyanosis, shock, coma or convulsions) should receive oxygen therapy during resuscitation to target SpO₂ ≥94%; otherwise, the target SpO₂ is ≥92%.
- Titrate oxygen therapy up and down to reach targets by means of nasal cannula, a simple face mask or a face mask with reservoir bag, as appropriate:

		
O ₂ dose 1–5 L/min	O ₂ dose 6–10 L/min	O ₂ dose 10–15 L/min
FiO ₂ estimate 0.25–0.40	FiO ₂ estimate 0.40– 0.60	FiO ₂ estimate 0.60–0.95
Nasal cannula	Simple face mask	Face mask with reservoir bag

✓ Use conservative fluid management in patients with COVID-19 when there is no evidence of shock.

Aggressive fluid resuscitation may worsen oxygenation, especially in settings where there is limited availability of mechanical ventilation.^{8,9}

✓ If a clinical suspicion for co-infection exists, consider empiric antimicrobials to treat co-pathogens causing the syndrome, particularly in severe cases. This may include conventional and atypical bacterial pathogens, influenza and PJP (see section 3.3 above).

✓ Closely monitor patients with SARI for signs of clinical deterioration, such as rapidly progressive respiratory failure and sepsis, and apply supportive care interventions immediately.

4.3 Specific therapies

✗ Do not routinely give systemic corticosteroids for treatment of COVID-19 unless they are indicated for another reason.

A systematic review of observational studies of corticosteroids administered to patients with SARS reported no survival benefit and possible harms (avascular necrosis, psychosis, diabetes, and delayed viral clearance).¹⁰ A systematic review of observational studies in influenza found a higher risk of mortality and secondary infections with corticosteroids; the evidence was judged as very low to low quality due to confounding by indication.¹¹ A subsequent study that addressed this limitation by adjusting for time-varying confounders found no effect on mortality.¹² Finally, a recent study of patients receiving corticosteroids for MERS used a similar statistical approach and found no effect of corticosteroids on mortality but delayed lower respiratory tract (LRT) clearance of MERS-CoV.¹³

Given lack of effectiveness and possible harm, routine corticosteroids should be avoided unless they are indicated for another reason.

! There is no current evidence from RCTs to recommend any specific treatment for patients with suspected or confirmed COVID-19 infection. This is an area of active study. Candidate drugs undergoing investigation include remdesivir, lopinavir/ritonavir, chloroquine, interferon, and tocilizumab. To date, published clinical data on most of these agents consists largely of *in vitro* studies, with little or no human data. Hence we do not know whether these medicines benefit or cause harm to patients with COVID-19. One exception is lopinavir/ritonavir, which was studied in a recent randomized control trial of patients with severe COVID-19.¹⁴ No statistically significant benefit was seen with respect to viral load, time to clinical improvement, or mortality. Chloroquine has received considerable interest, given its relatively low cost, (limited) local availability, known side-effect profile (at registered doses), and some promising *in vitro* data. Published data from human trials is currently lacking.¹⁵ Given the state of evidence, we suggest consideration of the following:

- Where possible, consideration should be given to enroll hospitalized patients in clinical trials. This provides both adequate monitoring and ethics oversight, and affords the opportunity to contribute to the therapeutics evidence base for future patients.
- Where investigational therapeutics are given outside of a clinical trial, this should be done under the Monitored Emergency Use of Unregistered Interventions (MEURI) framework, whereby it can be ethically appropriate to offer individuals investigational interventions on an emergency basis in the context of an outbreak characterized by high mortality.¹⁶ The principles of this include:
 - Data providing preliminary support for the intervention's efficacy and safety are available, at least from laboratory or animal studies.
 - The relevant human research ethics committee has approved the therapeutics' use.
 - The patient's informed consent is obtained.
 - Adequate resources are devoted to minimizing the therapeutics' risk.
 - The results of the intervention are documented and shared with the wider medical and scientific community.
- Where therapeutics are given to patients outside of a clinical trial, these should be reserved only for *hospitalized patients* (rather than given to mild cases, the vast majority of whom will recover fully without any intervention).

! There is no evidence for the use of any drug or vaccine to prevent COVID-19 infection. Prevention consists of non-pharmaceutical interventions, such as good hand hygiene and social distancing.

Statement on the use of angiotensin-converting enzyme inhibitors (ACEi), angiotensin receptor blockers (ARBs), and nonsteroidal anti-inflammatory drugs in COVID-19 patients

Recent work suggested that patients on ACEi or ARBs upregulate ACE2 receptors, the binding site for SARS-CoV-2, within tissues including the lung and heart, prompting concern that this might place patients at risk of worse outcomes with COVID-19.¹ For the moment, this remains purely theoretical, with no evidence of a linkage to poor clinical outcomes. In addition, discontinuing or switching ACEi or ARBs to alternative agents may be deleterious to patient care. **Pending further evidence we therefore do not recommend switching patients off ACEi or ARBs unless there are other medical reasons to do so.**

The evidence regarding nonsteroidal anti-inflammatory drugs (NSAIDs) in COVID-19 is similarly lacking. Nonetheless, where short-term fever or pain relief is required, it may be prudent to use drugs from other classes, such as paracetamol. For patients requiring NSAIDs for other indications however, the evidence is not definitive enough to recommend discontinuation. The South African Health Products Regulatory Authority (SAHPRA) has released a communication on the topic of NSAIDs and COVID-19.²

4.4 Management of hypoxemic respiratory failure and ARDS

✔ **Recognize severe hypoxemic respiratory failure when a patient with respiratory distress is failing standard oxygen therapy.** Patients may continue to have increased work of breathing or hypoxemia ($\text{SpO}_2 < 90\%$, $\text{PaO}_2 < 60 \text{ mmHg}$ [$< 8.0 \text{ kPa}$]) even when oxygen is delivered via a face mask with reservoir bag. Hypoxemic respiratory failure in ARDS commonly results from intrapulmonary ventilation-perfusion mismatch or shunt and usually requires mechanical ventilation.

⚠ **High-flow nasal oxygen (HFNO) or non-invasive ventilation (NIV) should only be used in selected patients with hypoxemic respiratory failure. The risk of treatment failure was high in patients with MERS treated with NIV, and patients treated with either HFNO or NIV should be closely monitored for clinical deterioration. In addition, HFNO and NIV carry the risk of aerosolization of viral particles against which adequate precautions need to be taken.** Patients with hypercapnia (exacerbation of obstructive lung disease, cardiogenic pulmonary oedema), hemodynamic instability, multi-organ failure, or abnormal mental status should generally not receive HFNO, although emerging data suggest that HFNO may be safe in patients with mild-moderate and non-worsening hypercapnia.¹⁷ Patients receiving HFNO should be in a monitored setting and cared for by experienced personnel capable of endotracheal intubation in case the patient acutely deteriorates or does not improve after a short trial (about 1-2 hrs).

- Risks of NIV include delayed intubation, large tidal volumes, and injurious transpulmonary pressures. Limited data suggest a high failure rate when MERS patients receive NIV.¹³
- *A single patient room and airborne precautions should be taken whenever HFNO and NIV is used.*

✔ **For intubated patients with ARDS use lung-protective ventilation strategies.** Always consult an expert intensivist if possible. Detailed recommendations on mechanical ventilation strategies are beyond the scope of the guideline. Nonetheless, the general principles in patients with ARDS include:

- Aim for an initial tidal volume of 6ml/kg.¹⁸ Higher tidal volume up to 8 ml/kg predicted body weight may be needed if undesirable side effects occur (e.g. dyssynchrony, pH <7.15).
- Strive to achieve the lowest plateau pressure possible. Plateau pressures above 30cm H₂O are associated with an increased risk of pulmonary injury.¹⁸
- Hypercapnia is permitted if meeting the pH goal of 7.30-7.45.
- Application of prone ventilation >12 hours a day is strongly recommended for patients with severe ARDS.¹⁸
- In patients with moderate or severe ARDS, moderately higher PEEP instead of lower PEEP is suggested.¹⁸
- The use of deep sedation may be required to control respiratory drive and achieve tidal volume targets.
- In patients with moderate-severe ARDS (PaO₂/FiO₂ <150), neuromuscular blockade by continuous infusion should not be routinely used.¹⁹ Continuous neuromuscular blockade may still be considered in patients with ARDS in certain situations: ventilator dyssynchrony despite sedation, such that tidal volume limitation cannot be reliably achieved; or refractory hypoxemia or hypercapnia.
- In settings with access to expertise in extracorporeal life support (ECLS), consider referral of patients with refractory hypoxemia despite lung protective ventilation.²⁰⁻²²
- Avoid disconnecting the patient from the ventilator, which results in loss of PEEP and atelectasis. Use in-line catheters for airway suctioning and clamp endotracheal tube when disconnection is required (for example, transfer to a transport ventilator). A high efficiency particulate filter on the expiratory limb of the ventilator circuit should be used.

4.5 De-isolation criteria

Patients can be de-isolated 14 days after the onset of their symptoms (mild cases), or 14 days after achieving clinical stability (moderate-severe cases).

Most patients with mild COVID-19 infection continue to shed SARS-CoV-2 from their upper airways for approximately 7-12 days.²³⁻²⁵ The duration of shedding is longer in severe cases, though in both mild and severe cases, significant variation is seen.^{23,24,26}


Viral shedding does not necessarily equate to infectiousness however. Viral shedding may decline to a level below the infectious threshold before it ceases completely, and/or non-viable virus may be shed. In a small cohort of mild COVID-19 cases from Germany (n=9), viral loads and viral cultures were performed on a variety of specimens simultaneously.²⁷ The virus was readily culturable from specimens taken during the first week of symptoms, but no positive cultures were obtained from samples taken after day 8. Importantly, this was despite ongoing high viral loads being detected at the time. The authors estimated that there would be a <5% chance of successful culture by day 10.

Given the very small sample size of the German cohort, we suggest a cautious approach of de-isolating patients with mild disease 14 days after symptom onset.

Patients with severe disease (i.e. requiring admission due to clinical instability) may continue to shed virus at higher levels for longer periods. We therefore suggest de-isolating such patients 14 days after clinical stability has been achieved (e.g. after supplemental oxygen was discontinued).

Patients who remain asymptomatic after a positive COVID-19 result can be de-isolated 14 days after their positive test. Although asymptomatic patients might be expected to be less infectious than symptomatic patients, in one study the two groups' viral loads were shown to be similar, and we believe a similarly cautious approach to de-isolation is warranted.²⁶

Patients admitted to hospital can continue their isolation period at home once clinical stability has been achieved, provided that the criteria in table 2 are met.



5. Infection prevention and control (IPC)

IPC is a critical and integral part of clinical management of patients and should be initiated at the point of entry of the patient to hospital (typically the Emergency Department). **A combination of standard, contact and droplet precautions should be practiced for all COVID-19 cases, and further precautions when performing aerosol-generating procedures (AGP).**

Standard precautions are used to prevent or minimize transmission of pathogens at all times, and should be applied to all patients in healthcare facilities irrespective of their diagnosis or status. These include hand hygiene, appropriate use of PPE, safe handling of sharps, linen and waste, disinfection of patient care articles, respiratory hygiene, occupational health and injection safety.

Transmission-based precautions - droplet, and contact:

- Hand hygiene is the first and most essential aspect
- Healthcare worker PPE consists of gloves, gown (or apron), and a medical mask.
- Safe waste management
- Use either disposable or dedicated equipment (e.g. stethoscopes, blood pressure cuffs and thermometers). If equipment needs to be shared among patients, clean and disinfect between each patient use.
- Limit patient movement within the institution (e.g. where possible, use portable X-rays rather than sending the patient to the X-ray department), and ensure that patients wear medical masks when outside their rooms.

Aerosol-generating Procedures:

Aerosol precautions are required when performing aerosol-generating procedures. These include taking respiratory tract samples for SARS-CoV-2 testing (such as nasopharyngeal and oropharyngeal swabs), intubation, bronchoscopy, open suctioning of the respiratory tract, and cardiopulmonary resuscitation.

Aerosol precautions for healthcare workers:

- Healthcare worker PPE consists of gloves, gown (or apron), a fit-tested particulate respirator (N95 respirator), and eye protection (goggles or face shield).
- Use an adequately ventilated single room when performing aerosol-generating procedures, with spacing between beds of at least 1-1.5 metres.

For more details, refer to the World Health Organization IPC guidelines:

<https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance/infection-prevention-and-control>

6. Recording and reporting

The goal of clinical management is to reduce morbidity and mortality from COVID-19. It is important to record and report the cases of COVID-19 disease in order to track the size and severity of the epidemic, the care received by patients in and out of hospital and identify areas for improvement in current and future outbreaks. There are different tools which will be needed to record and report clinical cases of COVID-19.

Tool	When to complete	Comments
Person of interest form	To be completed for all individuals <u>suspected</u> of COVID 19 disease and have a specimen taken	
NMC case notification	To be completed for all cases who meet the case definition for COVID-19	Can be completed online using NICD NMC mobile or web based app
Admission form (For <u>inpatients</u>)	To be completed for all <u>confirmed</u> patients admitted to a health care facility at admission or as soon as possible after admission	This form will document presence of co-morbidities, severity of illness at admission
Daily monitoring form (separate forms for inpatients and outpatients)	To be completed for all <u>confirmed</u> patients for each day until they are considered cured (by PCR criteria).	This form will document the daily symptoms, signs and severity of disease during admission
Discharge form (different forms for inpatients and outpatients)	To be completed for all <u>confirmed</u> patients	This form will document patient outcomes such as death, transfer or discharge.
Homecare form (for <u>outpatients</u>)	To be completed for all <u>confirmed</u> patients admitted with mild disease managed at home.	This form will document presence of co-morbidities, severity of illness at admission

The latest version of these forms are available from www.nicd.ac.za

References

1. Fang L, Karakiulakis G, Roth M. Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? *Lancet Respir Med*. 2020.
2. SAHPRA. The Use of Non-Steroidal Anti-Inflammatory Drugs in patients with Covid-19. Media release. 2020.
3. Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al. Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus-Infected Pneumonia. *N Engl J Med*. 2020.
4. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med*. 2020.
5. Yu P, Zhu J, Zhang Z, Han Y, Huang L. A familial cluster of infection associated with the 2019 novel coronavirus indicating potential person-to-person transmission during the incubation period. *J Infect Dis*. 2020.
6. Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA*. 2020.
7. World Health Organization. Clinical management of severe acute respiratory infection when novel coronavirus (nCoV) infection is suspected 2020 [Available from: [https://www.who.int/publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-\(ncov\)-infection-is-suspected](https://www.who.int/publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-(ncov)-infection-is-suspected)].
8. Schultz MJ, Dunser MW, Dondorp AM, Adhikari NK, Iyer S, Kwizera A, et al. Current challenges in the management of sepsis in ICUs in resource-poor settings and suggestions for the future. *Intensive Care Med*. 2017;43(5):612-24.
9. Rhodes A, Evans LE, Alhazzani W, Levy MM, Antonelli M, Ferrer R, et al. Surviving Sepsis Campaign: International Guidelines for Management of Sepsis and Septic Shock: 2016. *Intensive Care Med*. 2017;43(3):304-77.
10. Stockman LJ, Bellamy R, Garner P. SARS: systematic review of treatment effects. *PLoS Med*. 2006;3(9):e343.
11. Lansbury L, Rodrigo C, Leonardi-Bee J, Nguyen-Van-Tam J, Lim WS. Corticosteroids as adjunctive therapy in the treatment of influenza. *Cochrane Database Syst Rev*. 2019;2:CD010406.
12. Delaney JW, Pinto R, Long J, Lamontagne F, Adhikari NK, Kumar A, et al. The influence of corticosteroid treatment on the outcome of influenza A(H1N1pdm09)-related critical illness. *Crit Care*. 2016;20:75.
13. Arabi YM, Mandourah Y, Al-Hameed F, Sindi AA, Almekhlafi GA, Hussein MA, et al. Corticosteroid Therapy for Critically Ill Patients with Middle East Respiratory Syndrome. *Am J Respir Crit Care Med*. 2018;197(6):757-67.
14. Cao B, Wang Y, Wen D, Liu W, Wang J, Fan G, et al. A Trial of Lopinavir–Ritonavir in Adults Hospitalized with Severe Covid-19. *New England Journal of Medicine*. 2020.
15. Cortegiani A, Ingoglia G, Ippolito M, Giarratano A, Einav S. A systematic review on the efficacy and safety of chloroquine for the treatment of COVID-19. *J Crit Care*. 2020.
16. World Health Organization. Guidance For Managing Ethical Issues In Infectious Disease Outbreaks 2016. Available from: <https://apps.who.int/iris/bitstream/handle/10665/250580/9789241549837-eng.pdf;jsessionid=2C3A0BBB41D97192E283FF36FF1D7644?sequence=1>.
17. Lee MK, Choi J, Park B, Kim B, Lee SJ, Kim SH, et al. High flow nasal cannulae oxygen therapy in acute-moderate hypercapnic respiratory failure. *Clin Respir J*. 2018;12(6):2046-56.
18. Fan E, Del Sorbo L, Goligher EC, Hodgson CL, Munshi L, Walkey AJ, et al. An Official American Thoracic Society/European Society of Intensive Care Medicine/Society of Critical Care Medicine

- Clinical Practice Guideline: Mechanical Ventilation in Adult Patients with Acute Respiratory Distress Syndrome. *Am J Respir Crit Care Med*. 2017;195(9):1253-63.
19. National Heart L, Blood Institute PCTN, Moss M, Huang DT, Brower RG, Ferguson ND, et al. Early Neuromuscular Blockade in the Acute Respiratory Distress Syndrome. *N Engl J Med*. 2019;380(21):1997-2008.
 20. Combes A, Hajage D, Capellier G, Demoule A, Lavoue S, Guervilly C, et al. Extracorporeal Membrane Oxygenation for Severe Acute Respiratory Distress Syndrome. *N Engl J Med*. 2018;378(21):1965-75.
 21. Goligher EC, Tomlinson G, Hajage D, Wijeyesundera DN, Fan E, Juni P, et al. Extracorporeal Membrane Oxygenation for Severe Acute Respiratory Distress Syndrome and Posterior Probability of Mortality Benefit in a Post Hoc Bayesian Analysis of a Randomized Clinical Trial. *JAMA*. 2018;320(21):2251-9.
 22. Alshahrani MS, Sindi A, Alshamsi F, Al-Omari A, El Tahan M, Alahmadi B, et al. Extracorporeal membrane oxygenation for severe Middle East respiratory syndrome coronavirus. *Ann Intensive Care*. 2018;8(1):3.
 23. World Health Organization. Report of the WHO-China Joint Mission on Coronavirus Disease 2019 (COVID-19)2020. Available from: <https://www.who.int/docs/default-source/coronaviruse/who-china-joint-mission-on-covid-19-final-report.pdf>.
 24. Liu Y, Yan L-M, Wan L, Xiang T-X, Le A, Liu J-M, et al. Viral dynamics in mild and severe cases of COVID-19. *Lancet Infect Dis* [Internet]. 2020. Available from: [https://www.thelancet.com/journals/laninf/article/PIIS1473-3099\(20\)30232-2/fulltext](https://www.thelancet.com/journals/laninf/article/PIIS1473-3099(20)30232-2/fulltext).
 25. Young BE, Ong SWX, Kalimuddin S, Low JG, Tan SY, Loh J, et al. Epidemiologic Features and Clinical Course of Patients Infected With SARS-CoV-2 in Singapore. *JAMA*. 2020.
 26. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020.
 27. Woelfel R, Corman VM, Guggemos W, Seilmaier M, Zange S, Mueller MA, et al. Clinical presentation and virological assessment of hospitalized cases of coronavirus disease 2019 in a travel-associated transmission cluster. *medRxiv*. 2020:2020.03.05.20030502.

Appendix 1 – Example of a patient information sheet

Example of a patient information sheet for use with suspected cases who are being sent home to await test results for SARS-CoV-2 (COVID-19).

While awaiting test results for COVID-19 (the novel coronavirus), you have been assessed as being medically well enough to be managed at home.

However, please consider yourself as potentially infectious until the final results are available. You will need to abide by the following:





- You should quarantine yourself at home. Don't go to work, avoid unnecessary travel, and as far as possible avoid close interactions with other people.
- You should clean your hands with soap and water frequently. Alcohol-based sanitizers may also be used, provided they contain at least 60% alcohol.
- Do not have visitors in your home. Only those who live in your home should be allowed to stay. If it is urgent to speak to someone who is not a member of your household, do this over the phone.
- You should wear a facemask when in the same room (or vehicle) as other people.
- At home, you should stay in a specific room and use your own bathroom (if possible). If you live in shared accommodation (university halls of residence or similar) with a communal kitchen, bathroom(s) and living area, you should stay in your room with the door closed, only coming out when necessary, wearing a facemask if one has been issued to you.
- You should practice good cough and sneeze hygiene by coughing or sneezing into a tissue, discarding the tissue immediately afterwards in a lined trash can, and then wash your hands immediately.
- If you need to wash the laundry at home before the results are available, then wash all laundry at the highest temperature compatible for the fabric using laundry detergent. This should be above 60° C. If possible, tumble dry and iron using the highest setting compatible with the fabric. Wear disposable gloves and a plastic apron when handling soiled materials if possible and clean all surfaces and the area around the washing machine. Do not take laundry to a laundrette. Wash your hands thoroughly with soap and water after handling dirty laundry (remove gloves first if used).
- You should avoid sharing household items like dishes, cups, eating utensils and towels. After using any of these, the items should be thoroughly washed with soap and water.
- All high-touch surfaces like table tops, counters, toilets, phones, computers, etc. that you may have touched should be appropriately and frequently cleaned.
- Monitor your symptoms - Seek prompt medical attention if your illness is worsening, for example, if you have difficulty breathing, or if the person you are caring for symptoms are worsening. If it's not an emergency, call your doctor or healthcare facility at the number below. If it is an emergency and you need to call an ambulance, inform the call handler or operator that you are being tested for SARS-CoV-2.

While awaiting the results, if your symptoms worsen:

- Call:
- Or come to:

For more information on COVID-19, see the NICD's FAQ page:
<http://www.nicd.ac.za/diseases-a-z-index/covid-19/frequently-asked-questions/>

Standard precautions to prevent transmission of COVID-19

	<p>Keep your hands clean</p> <p>When?</p> <ul style="list-style-type: none"> • After visiting the bathroom • Before and after eating • After blowing your nose • Whenever you think your hands are dirty <p>How? Use alcohol hand rub or wash hands with soap and water</p> <p>Caution Never touch your eyes, nose or mouth with unwashed hands</p>
	<p>Cough etiquette</p> <ul style="list-style-type: none"> • Keep a distance of 2 meters between you and a person with a cough • Cover your own cough or sneeze with a tissue • Once used, throw the tissue away in a closed container • Clean your hands afterwards
	<ul style="list-style-type: none"> • Do not share items with other people (clothing, blankets, pillows, towels, mobile phones, uncovered food, magazines, books) • Do not keep the toilet lid up when you flush the toilet (you can transmit the virus from all body excretions)
	<p>Keep your immediate environment clean</p> <ul style="list-style-type: none"> • Wipe frequently-touched areas regularly with a disinfectant cloth • Discard all waste immediately