



Page	Content
2	Surveillance programme description
3	Comments
4-5	Systematic Influenza-like illness (ILI) surveillance Influenza Respiratory syncytial virus
6-7	Influenza-like illness (ILI) Viral Watch Influenza
8-10	National syndromic surveillance for pneumonia Influenza Respiratory syncytial virus
11	Private Hospital Consultation Outpatient consultations In patient consultations

Influenza and Respiratory Syncytial Virus Surveillance Report

Programme Descriptions

Programme	Influenza-like illness (ILI)	Viral Watch	National syndromic surveillance for pneumonia	Private hospital consultations
Start year	2012	1984	2009	2002
Provinces*	KZ NW WC**	EC FS GP LP MP NC NW WC	GP KZ MP NW WC	EC FS GP LP MP NW WC
Type of site	Primary health care clinics	General practitioners	Public hospitals	Private hospitals
Case definition	An acute respiratory illness with a temperature ($\geq 38^{\circ}\text{C}$) and cough, & onset ≤ 10 days	An acute respiratory illness with a temperature ($\geq 38^{\circ}\text{C}$) and cough, & onset ≤ 10 days	Acute (symptom onset ≤ 10 days) or chronic (symptom onset > 10) lower respiratory tract infection	ICD codes J10-J18
Specimens collected	Oropharyngeal & nasopharyngeal swabs	Throat and/or nasal swabs or Nasopharyngeal swabs	Oropharyngeal & nasopharyngeal swabs	Not applicable
Main pathogens tested***	INF RSV BP	INF RSV BP	INF RSV BP	Not applicable

Epidemic Threshold

Thresholds are calculated using the Moving Epidemic Method (MEM), a sequential analysis using the R Language, available from: <http://CRAN.R-project.org/web/package=mem>) designed to calculate the duration, start and end of the annual influenza epidemic. MEM uses the 40th, 90th and 97.5th percentiles established from available years of historical data to calculate thresholds of activity. Thresholds of activity for influenza and RSV are defined as follows: Below seasonal threshold, Low activity, Moderate activity, High activity, Very high activity. For influenza, thresholds from outpatient influenza like illness (Viral Watch Programme) are used as an indicator of disease transmission in the community and thresholds from pneumonia surveillance are used as an indicator of impact of disease.

* EC: Eastern Cape; FS: Free State; GP: Gauteng; KZ: KwaZulu-Natal; LP: Limpopo; MP: Mpumalanga; NC: Northern Cape; NW: North West; WC: Western Cape

**Started in 2019

***INF: influenza virus; RSV: respiratory syncytial virus; BP: *Bordetella pertussis*

Influenza and Respiratory Syncytial Virus Surveillance Report

Reporting period 01/01/2020 to 05/04/2020

Results until end of epidemiologic week 14(2020)

Comments:

Influenza

The 2020 influenza season has not yet started although sustained detections of influenza A(H1N1)pdm09 and to a lesser extent influenza B(Victoria) have been made, from Western Cape Province, in all surveillance programmes. Influenza has only occasionally been detected from other provinces. The increase in cases in the Western Cape Province began as a localised increase in case numbers with clusters identified in universities and schools but now appears to be more widespread as indicated by influenza detection in all surveillance programmes.

ILI programme: In 2020 to date, specimens from 409 patients were received from 4 ILI sites. Influenza was detected in 52 specimens (all from Western Cape province), 36 (69%) were identified as influenza A(H1N1)pdm09, one (2%) influenza A subtype inconclusive, 12 (23%) as influenza B (Victoria) and three (6%) B lineage inconclusive.

Viral Watch programme: During the same period, specimens were received from 183 patients from Viral Watch sites in 8 provinces. Influenza was detected in 77 patients (76 from Western Cape and one imported case from Gauteng provinces), of which 74 (97%) were influenza A(H1N1)pdm09, one (1%) influenza A(H3N2) and one (1%) influenza B(Victoria). Seven (9%; 7/75) of the positive cases had history of international travel.

Pneumonia surveillance: Since the beginning of 2020, specimens from 875 patients with severe respiratory illness (SRI) were received from the 6 sentinel sites. Influenza was detected in 24 patients (all from Western Cape province), of which 20 (83%) were influenza A(H1N1)pdm09, one (4%) influenza A subtype inconclusive and three (13%) influenza B(Victoria).

Respiratory syncytial virus

The 2020 RSV season has not yet started. However, sporadic detections of RSV are being made. In the previous 3 years the average start of RSV season ranged between week 7 to week 9, therefore the start of the season is delayed compared to previous years in 2020.

ILI programme: In 2020 to date, 409 specimens were tested and RSV was detected in specimens of seven (2%) patients.

Viral Watch programme: During the same period, 183 specimens were tested and RSV was not detected.

Pneumonia surveillance: Since the beginning of 2020, 875 specimens were tested and RSV was detected in specimens of 60 (7%) patients.


Influenza and Respiratory Syncytial Virus Surveillance Report

Reporting period 01/01/2020 to 05/04/2020

Results until end of epidemiologic week 14(2020)

Influenza-like illness (ILI) surveillance primary health care clinics

Figure 1. Number of positive samples* by influenza subtype and lineage and detection rate** by week



*Specimens from patients with influenza-like illnesses at 4 sentinel sites in 3 provinces

**Only reported for weeks with >10 specimens submitted

Inconclusive: insufficient viral load in sample and unable to characterise further

Table 1. Cumulative number of influenza subtype and lineage and total number of samples tested by clinic and province

Clinic (Province)	A(H1N1)pdm09	A(H3N2)	A subtype inconclusive	B/Victoria	B/Yamagata	B lineage inconclusive	Total samples
Eastridge (WC)	33	0	1	12	0	3	194
Edendale Gateway (KZ)	0	0	0	0	0	0	80
Jouberton (NW)	0	0	0	0	0	0	80
Mitchell's Plain (WC)	3	0	0	0	0	0	55
Total:	36	0	1	12	0	3	409

KZ: KwaZulu-Natal; NW: North West; WC: Western Cape

Inconclusive: insufficient viral load in sample and unable to characterise further


Influenza and Respiratory Syncytial Virus Surveillance Report

Reporting period 01/01/2020 to 05/04/2020

Results until end of epidemiologic week 14 (2020)

Influenza-like illness (ILI) surveillance primary health care clinics

Figure 2. Number of samples testing positive for respiratory syncytial virus by subgroup and detection rate by week



Inconclusive: insufficient viral load in sample and unable to characterise further
 RSV AB: Both RSV A and B subgroup identified

Table 2. Cumulative number of respiratory syncytial virus subgroups identified and total number of samples tested by clinic and province

Clinic (Province)	RSVA	RSVB	RSVAB	RSV subgroup inconclusive	Total samples
Eastridge (WC)	0	2	0	0	194
Edendale Gateway (KZ)	3	2	0	0	80
Jouberton (NW)	0	0	0	0	80
Mitchell's Plain (WC)	0	0	0	0	55
Total	3	4	0	0	409

KZ: KwaZulu-Natal; NW: North West; WC: Western Cape
 Inconclusive: insufficient viral load in sample and unable to characterise further
 RSV AB: Both RSV A and B subgroup identified


Influenza and Respiratory Syncytial Virus Surveillance Report

Reporting period 01/01/2020 to 05/04/2020

Results until end of epidemiologic week 14(2020)

Influenza-like illness (ILI) surveillance Viral Watch

Figure 4. Number of positive samples* by influenza subtype and lineage and detection rate** by week




*Specimens from patients with Influenza-like illnesses at 92 sentinel sites in 8 provinces

** Only reported for weeks with >10 specimens submitted.

Inconclusive: insufficient viral load in sample and unable to characterise further

Figure 5. ILI surveillance Viral Watch percentage influenza detections and epidemic thresholds*



*Thresholds based on 2010-2019 data

Influenza and Respiratory Syncytial Virus Surveillance Report

Reporting period 01/01/2020 to 05/04/2020

Results until end of epidemiologic week 14(2020)

Table 4. Cumulative number of influenza subtype and lineage and total number of samples tested by province

Province	A(H1N1)pdm09	A(H3N2)	A subtype inconclusive	B/Victoria	B/Yamagata	B lineage inconclusive	Total samples
Eastern Cape	0	0	0	0	0	0	4
Free State	0	0	0	0	0	0	0
Gauteng	1	0	0	0	0	0	36
Limpopo	0	0	0	0	0	0	1
Mpumalanga	0	0	0	0	0	0	2
North West	0	0	0	0	0	0	0
Northern Cape	0	0	0	0	0	0	4
Western Cape	74	1	0	1	0	0	136
Total:	75	1	0	1	0	0	183

Inconclusive: insufficient viral load in sample and unable to characterise further

From 01 January 2020 to date, 10 patients were tested for influenza at the time of entry into South Africa following travel abroad and influenza was detected in three patients, of which one influenza A(H1N1)pdm09 and two influenza A(H3N2).

Patients known to have acquired influenza abroad are not included in the table or epidemiological curve.


Influenza and Respiratory Syncytial Virus Surveillance Report

Reporting period 01/01/2020 to 05/04/2019

Results until end of epidemiologic week 14(2020)

National syndromic surveillance for pneumonia

Figure 6. Number of positive samples* by influenza subtype and lineage and detection rate** by week




*Specimens from patients hospitalised with pneumonia at 6 sentinel sites in 5 provinces

**Only reported for weeks with >10 specimens submitted

Inconclusive: insufficient viral load in sample and unable to characterise further

Figure 7. National syndromic surveillance for pneumonia percentage influenza detections and epidemic thresholds*



*Thresholds based on 2010-2019 data

Influenza and Respiratory Syncytial Virus Surveillance Report

Reporting period 01/01/2020 to 05/04/2020

Results until end of epidemiologic week 14(2020)

Table 5. Cumulative number of identified influenza subtype and lineage and total number of samples tested by hospital

Hospital (Province)	A(H1N1)pdm09	A(H3N2)	A subtype inconclusive	B/Victoria	B/Yamagata	B lineage inconclusive	Total samples
Edendale (KZ)	0	0	0	0	0	0	115
Helen Joseph-Rahima Moosa (GP)	0	0	0	0	0	0	212
Klerksdorp-Tshepong (NW)	0	0	0	0	0	0	117
Mapulaneng-Matikwana (MP)	0	0	0	0	0	0	91
Red Cross (WC)	18	0	0	1	0	0	252
Mitchell's Plain (WC)	2	0	1	2	0	0	88
Total:	20	0	1	3	0	0	875

GP: Gauteng; KZ: KwaZulu-Natal; NW: North West; MP: Mpumalanga; WC: Western Cape

Inconclusive: insufficient viral load in sample and unable to characterise further


Influenza and Respiratory Syncytial Virus Surveillance Report

Reporting period 01/01/2020 to 05/04/2020

Results until end of epidemiologic week 14(2020)

National syndromic surveillance for pneumonia

Figure 8. Number of samples testing positive for respiratory syncytial virus by subgroup and detection rate by week



Inconclusive: insufficient viral load in sample and unable to characterise further
 RSV AB: Both RSV A and B subgroup identified

Table 6: Cumulative number of respiratory syncytial virus subgroups identified and total number of samples tested by hospital

Hospital (Province)	RSVA	RSVB	RSVAB	RSV subgroup inconclusive	Total samples
Edendale (KZ)	2	5	0	1	115
Helen Joseph-Rahima Moosa (GP)	19	5	0	0	212
Klerksdorp-Tshepong (NW)	0	0	0	0	117
Mapulaneng-Matikwana (MP)	0	0	0	0	91
Red Cross (WC)	18	10	0	0	252
Mitchell's Plain (WC)	0	0	0	0	88
Total:	39	20	0	1	875

GP: Gauteng; KZ: KwaZulu-Natal; NW: North West; MP: Mpumalanga; WC: Western Cape
 Inconclusive: insufficient viral load in sample and unable to characterise further
 RSV AB: Both RSV A and B subgroup identified


Influenza and Respiratory Syncytial Virus Surveillance Report

Reporting period 08/04/2019 to 05/04/2020

Results until end of epidemiologic week 14(2020)

Private hospital consultations


Figure 9. Number of private hospital outpatient consultations* with a diagnosis of pneumonia and influenza (P&I) and viral isolates, week 15 2019 to week 14 2020**



* Hospital outpatient data from weekly reports of consultations to the Netcare hospital group. Discharge diagnosis is according to International Statistical Classification of Diseases and Related Health Problems coding by clinicians and does not represent laboratory confirmation of aetiology

** Influenza positive specimens from the Viral Watch surveillance programme

Figure 10. Number of private hospital admissions* with a discharge diagnosis of pneumonia and influenza (P&I) and viral isolates week 15 2019 to week 14 2020**



*Hospitalisation admission data from weekly reports of consultations to the Netcare hospital group. Discharge diagnosis is according to International Statistical Classification of diseases and Related Health Problems/ ICD by clinicians and does not represent laboratory confirmation of aetiology ** Influenza positive specimens from the national syndromic surveillance for pneumonia.