



## CONTENTS

### PAGE

### ZOONOTIC AND VECTOR-BORNE DISEASES

An update on rabies in South Africa 2

### INTERNATIONAL OUTBREAKS OF IMPORTANCE

A description of COVID-19 clusters in Nelson Mandela Bay District, Eastern Cape Province 3

An update on Ebola virus disease outbreak, Democratic Republic of Congo 4

### SEASONAL DISEASES

Respiratory syncytial virus and influenza seasons, 2020 5

### BEYOND OUR BORDERS

6

### WHO-AFRO: OUTBREAKS AND EMERGENCIES

9

## Editor's Note



Dr Ann Mathews

In this month's Communicable Diseases Communiqué, we provide a description of the coronavirus disease 2019 (COVID-19) clusters in the Eastern Cape Province, including overlapping events and geographical locations.

We also explore the respiratory syncytial virus (RSV) and influenza seasons, noting that the detection rate of RSV has remained below the seasonal threshold as compared to trends over the last ten years, and that the influenza season, which occurs mainly during the winter months of May to August, has not yet started.

The Ebola outbreak in the Democratic Republic of Congo (DRC) continues since the resurgence of cases on 10 April 2020, though there have been no new cases since early May 2020.

The routine rabies report describes the only recognised case of rabies in the country for this year, noting that the COVID-19 pandemic will have had an impact on reporting.

Internationally, the few other outbreaks of interest include cases of human plague in the DRC, cases of Japanese encephalitis in India, cholera in Kenya, Ethiopia and Somalia, and West Nile fever in Brazil.

## ZOONOTIC AND VECTOR-BORNE DISEASES

### An update on rabies in South Africa


In May 2020, we report a probable case of human rabies from the Eastern Cape Province. The case involved a 10-year-old child who died shortly after admission in hospital on 13 May. On 17 April 2020, the child was attacked by a dog and received rabies vaccine following the exposure. It was unclear if she also received rabies immunoglobulin as part of the rabies post-exposure prophylaxis (PEP) regimen. Although completely well prior to presentation to hospital, on admission, she was unconscious and had seizures with associated neurological fall-out. The NICD obtained saliva and cerebrospinal fluid samples for rabies testing on the day of death. These samples tested negative for rabies by PCR. No post-mortem brain samples were obtained to confirm rabies in this child. Given the history of exposure to a dog prior to illness and a clinical history that is compatible with the diagnosis of rabies, this case was classified as a probable rabies case in the absence of laboratory confirmation.

For South Africa this year, only one case of human rabies, reported here, has been identified. The COVID-19 pandemic and its response have led to a setback of several public health programmes, including rabies. South Africa may have unrecognised cases, considering that during 2019 16 cases were reported, of which seven were from the Eastern Cape Province and the other nine were from KwaZulu-Natal and Limpopo provinces. As an example, in the Eastern Cape Province, sampling of animals that were euthanised or found dead reported 33 animal rabies cases in 2020 (Figure 1). Canine rabies remains a veterinary and public health crisis in South Africa's east coast. Nearly all human deaths are caused

by dog bites. Children are at high risk, with more than half of 2019's reported rabies cases aged less than 18 years (56%, 9/16). This is a recognised phenomenon as children are more vulnerable to not disclosing dog or other animal contact or bite, and are therefore unlikely to receive appropriate and timely anti-rabies PEP.

With appropriate post-exposure prophylaxis, rabies is 100% preventable. There is a need to promote continued awareness in the community and at healthcare facilities. Details on rabies, its treatment and prevention, and collection of specimens for diagnosis can be found on the NICD website: [www.nicd.ac.za](http://www.nicd.ac.za).

#### EASTERN CAPE RABIES 21 May 2020



**Figure 1.** Canine and human rabies detection in Eastern Cape Province.

**Source:** Kevin le Roux, Department of Agriculture, Environment and Rural Development, KwaZulu-Natal Province.

**Article source:** Centre for Emerging Zoonotic and Parasitic Diseases, NICD-NHLS; [januszp@nicd.ac.za](mailto:januszp@nicd.ac.za)