

# PRIVATE CONSULTATIONS SURVEILLANCE REPORT

SOUTH AFRICA WEEK 33 2020

## CONTENTS

Programme descriptions	2
Interpretation of data presented	3-4
Data from virologic surveillance programmes to aid in interpretation of consultation trends	5
Number of consultations - all respiratory including confirmed or suspected COVID-19 and other by week	6
All ages	7
0-4 years of age	8
5-19 years of age	9
20-49 years of age	10
≥50 years of age	11
Eastern Cape Province	12
Gauteng Province	13
KwaZulu-Natal Province	14
Western Cape Province	15

## HIGHLIGHTS: WEEK 33

- The increase in total numbers of respiratory hospitalisations observed in previous weeks has gradually declined in the past four weeks, although the numbers are still slightly higher than pre-lock down period.
- The proportion of respiratory hospitalisations in all ages remains at the very high level, though there has been a decrease in the past two weeks in all provinces evaluated. The decline was especially noted among individuals in 20-49 and ≥50 year age groups.
- Percentage of casualty visits coded as respiratory remains in very high levels in individuals aged 20-49 years and ≥50 years.
- Differences by province and age group should be interpreted with caution due to low numbers in some groups.

# PRIVATE CONSULTATIONS SURVEILLANCE REPORT

WEEK 33 2020

## PROGRAMME DESCRIPTIONS

Inpatient data from a large national private hospital group and outpatient data from a general practitioner network linked to the same hospital group were received for the last week. Data were obtained from eight provinces (**Eastern Cape**, Free State, **Gauteng**, Limpopo, **KwaZulu-Natal**, Mpumalanga, North West, **Western Cape**). Sufficient numbers for province-level reporting were available for four of these (bold). Consultations and admissions were coded based on discharge diagnosis using the International Classification of Diseases and Related Health Problems, 10th revision (ICD-10). Data were analysed using the indicator: All respiratory and confirmed or suspected COVID-19 (J00-J99 & U07.1 & U07.2)/Total consultations. Data on the indicator Pneumonia and Influenza (J10-J18)/Total consultations are available on request but were not included in this report.

Data were categorised in the following age groups: All ages, <5 years, 5-19 years, 20-49 years, ≥50 years

### Epidemic Threshold

Thresholds were calculated using the Moving Epidemic Method (MEM), a sequential analysis using the R Language, available from: <http://CRAN.R-project.org/web/package=mem>, designed to calculate the duration, start and end of the annual influenza epidemic. MEM uses the 40th, 90th and 97.5th percentiles established from historical data (2015-2019 for inpatients, 2016-2019 for outpatients) to calculate thresholds of activity, defined as follows:

- Epidemic threshold: Median of weekly values for all baseline years
- Low activity: Between epidemic threshold including 40th percentile
- Moderate activity: Between 40th and 90th percentile
- High activity: Between 90th and 97.5th percentile
- Very high activity: 97.5th percentile and above

Hospitalization data for recent weeks are adjusted for delayed reporting (diagnosis codes assigned on discharge delayed for prolonged hospitalisations). Adjustment accounts for the probability of being admitted, but not yet discharged at the time of data drawdown using the age- and syndrome-specific probability distribution of duration of admission obtained from all hospitalizations that occurred during 2015-2019 and applied to the most recent weeks in 2020.

# PRIVATE CONSULTATIONS SURVEILLANCE REPORT

WEEK 33 2020

## INTERPRETATION OF DATA PRESENTED

**Total admissions** reduced from week 13 when lockdown was implemented and have remained below pre-lockdown levels.

**Total respiratory admissions** reduced from week 13 when lockdown was implemented and slowly increased exceeding pre-lockdown levels in week 26 and continued to increase up to week 29 reaching approximately double the level before the lockdown. From week 30, the numbers started to decrease. The proportion of admissions coded as confirmed COVID-19 (out of suspected) increased from week 15, exceeding 60% from week 26, peaked at ~ 75% in week 30 and started to come down.

**Total and respiratory outpatient (general practitioner) consultations** reduced from week 13. Respiratory consultations recovered to levels slightly lower than those preceding the lockdown from week 26 and 28, with a gradual decrease since week 29. The proportion of general practitioner consultations coded as confirmed COVID-19 (out of suspected) increased from week 15, with a sharp increase from week 29 onwards.

**Total and respiratory outpatient (casualty) consultations** reduced from week 13. Respiratory consultations recovered to levels similar to those preceding the lockdown from week 26. The proportion of casualty consultations coded as confirmed COVID-19 (out of suspected) increased from week 15, exceeding 50% from week 25.

**Proportion of admissions respiratory or suspected COVID-19** overall remained below threshold until week 21, following which it increased rapidly reaching the very high threshold in week 25 onwards with a decrease from week 29 but remains above the very high threshold to date.

By age group, percent admissions respiratory or suspected COVID for 0-4 years, has increased in recent weeks, remaining below the seasonal threshold. Among individuals aged 5-19 years, continuously increased since week 18, reaching low levels of activity in week 26.

Among individuals 20-49 years and ≥50 years, percent respiratory admissions has continuously increased since week 13, reaching very high level from week 21.

**Proportion of outpatient (general practitioner) consultations respiratory or suspected COVID-19** overall increased from week 11, peaking in week 13 then dropping well below the threshold, increased again crossing the seasonal threshold in week 25, peaked in week 30 and now below threshold.

By age group, percent outpatient visits (general practitioner) has breached the seasonal threshold among individuals aged 5-19 in week 27 and dropped again to below seasonal threshold from week 29 onwards. Among individuals aged 20-49 years and ≥50 years percent outpatient visits (general practitioner) breached seasonal threshold in week 25, peaked in week 28 at moderate levels for individuals aged 20-49 years and at low levels for individuals aged ≥50 years.

**Proportion of outpatient (casualty) consultations respiratory or suspected COVID-19** overall dropped from week 13 during the lockdown but then increased from week 23 reaching very high levels in week 28, now decreasing reaching low levels in week 32.

By age group, percent outpatient visits (casualty) showed similar trends, breached the seasonal threshold in age group 5-19 years and reaching very high levels in individuals aged 20-49 years and ≥50 years.

Trends in proportion of admissions and outpatient consultations respiratory or COVID varied by **province** with proportion inpatients respiratory reaching very high levels in all provinces evaluated. However, all four provinces evaluated experienced a downward trend, from week 28 in Eastern Cape and from week 30 in Gauteng, KwaZulu-Natal and Western Cape Provinces.

# PRIVATE CONSULTATIONS SURVEILLANCE REPORT

WEEK 33 2020

## Limitations

Thresholds are established based on the proportion of consultations which are respiratory. If numbers of non-respiratory consultations drop substantially because of changes in health-seeking behaviour, this could lead to elevated respiratory proportions. Delays in coding of consultations may lead to changes in data from previous weeks.

## Assessment

The increase in total numbers of respiratory hospitalisations observed in previous weeks has been decreasing in the past three weeks although the numbers are still slightly higher than pre-lock down period. The proportion of respiratory hospitalisations among 20-49 and  $\geq 50$  years, remains at very high levels, with a downward trend the past two weeks, this may suggest a decrease in respiratory cases including COVID-19 cases. Changes in health-seeking behaviours and/or effects of lockdown-related reductions may also have contributed.

The percentage of casualty visits coded as respiratory has decreased, with individuals aged 20-49 years and  $\geq 50$  years still in the very high threshold. Proportion outpatient visits to general practitioners are also showing a decline.

Differences by province and age group should be interpreted with caution due to low numbers in some groups.




# PRIVATE CONSULTATIONS SURVEILLANCE REPORT

WEEK 33 2020


## DATA FROM VIROLOGIC SURVEILLANCE PROGRAMMES TO AID IN INTERPRETATION OF CONSULTATION TRENDS

### NUMBER OF INFLUENZA POSITIVE SAMPLES BY SUBTYPE/LINEAGE AND DETECTION RATE BY WEEK


**Inpatients**




**Outpatients**



**Number of respiratory syncytial virus positive by groups and subgroups per week (Inpatients)**



**Number of samples tested and results for SARS-COV-2 per week (Inpatients)**





# PRIVATE CONSULTATIONS SURVEILLANCE REPORT

WEEK 33 2020

## NUMBER OF CONSULTATIONS - ALL RESPIRATORY INCLUDING CONFIRMED OR SUSPECTED COVID-19 AND OTHER CONSULTATIONS BY WEEK



(SOLID ARROW INDICATES FIRST WEEK OF LOCKDOWN, DASHED ARROW FIRST WEEK OF LEVEL 4)

### Inpatients





### Outpatients

#### A) General Practitioners



#### B) Casualty




# PRIVATE CONSULTATIONS SURVEILLANCE REPORT

WEEK 33 2020

## ALL AGES


All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Inpatients



All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Outpatients  
A) General Practitioners



B) Casualty




# PRIVATE CONSULTATIONS SURVEILLANCE REPORT


WEEK 33 2020

## 0-4 YEARS OF AGE


All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Inpatients



All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Outpatients  
A) General Practitioners



B) Casualty






# PRIVATE CONSULTATIONS SURVEILLANCE REPORT


WEEK 33 2020

## 5-19 YEARS OF AGE


All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Inpatients



All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Outpatients  
A) General Practitioners



B) Casualty




# PRIVATE CONSULTATIONS SURVEILLANCE REPORT


WEEK 33 2020

## 20-49 YEARS OF AGE


All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Inpatients



All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Outpatients  
A) General Practitioners



B) Casualty




# PRIVATE CONSULTATIONS SURVEILLANCE REPORT


WEEK 33 2020

## ≥50 YEARS OF AGE


All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Inpatients



All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Outpatients  
A) General Practitioners



B) Casualty




# PRIVATE CONSULTATIONS SURVEILLANCE REPORT

WEEK 33 2020

## EASTERN CAPE PROVINCE

All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Inpatients



All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Outpatients  
A) General Practitioners



B) Casualty






# PRIVATE CONSULTATIONS SURVEILLANCE REPORT


WEEK 33 2020

## GAUTENG


All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Inpatients



All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Outpatients  
A) General Practitioners



B) Casualty




# PRIVATE CONSULTATIONS SURVEILLANCE REPORT


WEEK 33 2020

## KWA ZULU NATAL


All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Inpatients



All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Outpatients  
A) General Practitioners



B) Casualty




# PRIVATE CONSULTATIONS SURVEILLANCE REPORT


WEEK 33 2020

## WESTERN CAPE PROVINCE

All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Inpatients



All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Outpatients  
A) General Practitioners



B) Casualty

