

Cumulative invasive pneumococcal disease case numbers reported by the GERMS-SA surveillance programme, 1 January 2012 to 31 August 2020

GERMS-SA surveillance programme

- GERMS-SA is a national, active, laboratory-based surveillance system initiated in 2003.
- Invasive pneumococcal disease (IPD) cases defined as hospitalised individuals with *Streptococcus pneumoniae* detected from normally sterile-site specimens (e.g. cerebrospinal fluid, blood or joint fluid).
- Repeat isolates from the same individual within 21 days were excluded.
- ~190 laboratories each year send reports and isolates.
- Age, sex, date of specimen collection, and source of specimen were captured.
- Pneumococcal isolates were serotyped by Quellung reaction using specific antisera (Statens Serum Institute, Copenhagen, Denmark). Culture-negative/bacterial antigen detection test positive, or isolates that lost viability were confirmed positive using a real-time *lytA* PCR¹ and serotyped using an adaption from the method described by Azzari *et al.*² This molecular assay includes targets for 38 serotypes (42 serotypes prior to 2014) and covers all serotypes included in PCV13. Only samples with an initial *lytA* PCR ct value of ≤ 35 were included. Where ct value was ≤ 35 but no serotype could be identified by including the 38 targets (42 targets prior to 2014), serotype was classified as non-vaccine type. Where *lytA* PCR ct value was ≥ 36 , serotype was classified as unknown and was not included in graphs. Where the PCR target could not distinguish between vaccine and non-vaccine serotype, serotype was classified as unknown and not included in the figures (targets: 18ABC, 18ABCF, 7AF, 9ALVN and 9AV).
- **Cumulative graph case numbers include viable isolates and those non-viable but characterised using molecular diagnostic techniques.**
- Figures 1 – 3 are for cases < 5 years, and Figures 4 – 6 for cases 5 years and older. Cases with unknown age were excluded from the figures.
- There are three graphs for each age group:
 - Disease caused by any of the seven serotypes in PCV7 (4, 6B, 9V, 14, 18C, 19F and 23F)
 - Disease caused by any of the six additional serotypes in PCV13 but not in PCV7 (1, 3, 5, 6A, 7F, 19A)
 - Disease caused by any serotypes not in PCV13
- Figures showing number of viable isolates submitted to GERMS-SA from 2008 to 2012 can be found in the appendix at the end of this report.
- More information on the GERMS-SA system available at:
<https://www.nicd.ac.za/centres/division-of-public-health-surveillance-and-response/>

PCV vaccine introduction in South Africa

- The 7-valent pneumococcal conjugate vaccine (PCV7) was introduced into the South African Expanded Programme on Immunisation in April 2009, with no catch-up vaccination campaign.
- There was a graded replacement of PCV7 by 13-valent pneumococcal conjugate vaccine (PCV13) in 2011. By June 2011 all provinces were using PCV13.

- There was a limited PCV13 catch-up campaign in 2011 and 2012.
- WHO/UNICEF vaccine coverage estimates for receiving a third dose of the PCV vaccine in South Africa are 10% in 2008, 58% in 2009, 62% in 2011, 75% in 2012, 77% in 2013, 85% in 2014, 85% 2015, 77% in 2016, 68% in 2017, and 73% in 2018.³
- The effect of the vaccine on IPD in South Africa has been described.⁴




Figure 1. Cumulative weekly number of disease episodes of invasive pneumococcal disease due to any of the seven serotypes (4, 6B, 9V, 14, 18C, 19F and 23F) in PCV7: children <5 years of age in South Africa, from 2012 to date. Viable isolates and those serotyped using molecular techniques included.




Figure 2. Cumulative weekly number of disease episodes of invasive pneumococcal disease due to any of the six additional (1, 3, 5, 6A, 7F, 19A) serotypes in PCV13 but not in PCV7: children <5 years of age in South Africa, from 2012 to date. Viable isolates and those serotyped using molecular techniques included. (Note: There is reported cross protection between 6A and 6B which is included in PCV7⁵)

Data are provisional as reported to date.




Figure 3. Cumulative weekly number of disease episodes of invasive pneumococcal disease due to any of the serotypes not in PCV13: children <5 years of age in South Africa, from 2012 to date. Viable isolates and those serotyped using molecular techniques included.




Figure 4. Cumulative weekly number of disease episodes of invasive pneumococcal disease due to any of the seven serotypes (4, 6B, 9V, 14, 18C, 19F and 23F) in PCV7: individuals ≥ 5 years of age in South Africa, from 2012 to date. Viable isolates and those serotyped using molecular techniques included.

Data are provisional as reported to date.




Figure 5: Cumulative weekly number of disease episodes of invasive pneumococcal disease due to any of the six additional (1, 3, 5, 6A, 7F, 19A) serotypes in PCV13 but not in PCV7: individuals ≥ 5 years of age in South Africa, from 2012 to date. Viable isolates and those serotyped using molecular techniques included (Note: There is reported cross protection between 6A and 6B which is included in PCV7⁵)




Figure 6. Cumulative weekly number of disease episodes of invasive pneumococcal disease due to any of the serotypes not in PCV13: individuals ≥5 years of age in South Africa, from 2012 to date. Viable isolates and those serotyped using molecular techniques included.

Missing information

Age was unknown for 954 of the cases (Table 1). By the time that this report was produced there were 7 viable isolates with pending serotype results (Table 2). For 315 isolates in the 9-year period, serotype could not be identified due to high ct value during *lytA* PCR, or PCR serotype target not distinguishing between vaccine and non-vaccine serotype.

Table 1. Isolates with missing age; number of viable, non-viable isolates and audit cases identified, January 2012 to date.

| | Age missing, n(%) | Viable, n(%) | Non-viable, n(%) | Audit/missing isolates, n(%) | Capture delays*, n(%) | Total |
|------|----------------------|--------------|------------------|---------------------------------|--------------------------|--------|
| 2012 | 248 (8) | 2,160 (67) | 273 (8) | 789 (24) | 0 (0) | 3222 |
| 2013 | 138 (5) | 1,932 (67) | 268 (9) | 665 (23) | 0 (0) | 2865 |
| 2014 | 165 (6) | 1,752 (64) | 291 (11) | 688 (25) | 0 (0) | 2731 |
| 2015 | 157 (6) | 1,700 (65) | 208 (8) | 727 (28) | 0 (0) | 2635 |
| 2016 | 41 (2) | 1,578 (65) | 197 (8) | 658 (27) | 0 (0) | 2433 |
| 2017 | 53 (2) | 1,535 (63) | 280 (11) | 625 (26) | 0 (0) | 2440 |
| 2018 | 55 (2) | 1,336 (58) | 327 (14) | 652 (28) | 0 (0) | 2315 |
| 2019 | 59 (2) | 1,387 (59) | 343 (15) | 632 (27) | 0 (0) | 2362 |
| 2020 | 38 (5) | 493 (62) | 125 (16) | 156 (20) | 24 (3) | 798 |
| All | 954 (5) | 13,873 (62) | 2312 (16) | 5592 (20) | 24 (3) | 21,801 |

*For 24 cases reported to CRDM, viability is unknown due to capturing delays.

Table 2. Cases where serotype was not available at the time this report was produced

| | Not typed | Unknown serotype | Viable, serotype pending | Non- viable, serotype pending | Viability unknown, serotype pending | Total serotypes pending |
|--------------|--------------|---------------------|--------------------------------|--|--|-------------------------------|
| 2012 | 38 | 9 | 0 | 0 | 0 | 0 |
| 2013 | 38 | 12 | 0 | 0 | 0 | 0 |
| 2014 | 1 | 39 | 0 | 0 | 0 | 0 |
| 2015 | 0 | 38 | 0 | 0 | 0 | 0 |
| 2016 | 2 | 30 | 0 | 0 | 0 | 0 |
| 2017 | 3 | 44 | 0 | 0 | 0 | 0 |
| 2018 | 0 | 38 | 0 | 0 | 0 | 0 |
| 2019 | 0 | 69 | 0 | 0 | 0 | 0 |
| 2020 | 0 | 36 | 7 | 24 | 24 | 55 |
| Total | 82 | 315 | 7 | 24 | 24 | 55 |

* Viability unknown due to capturing delays

Discussion

Compared to previous years, there was a marked reduction in IPD episodes in 2020. Due to the coronavirus disease 2019 (COVID-19) pandemic, a nation-wide lockdown was implemented on 26 March 2020. The reduction in IPD episodes could be related to reduced healthcare seeking behaviour, or closures of work-places, schools, universities, physical distancing and mandatory wearing of masks.⁶

Data Source

National Institute for Communicable Diseases | GERMS-SA

References

1. Carvalho MdGS, Tondella ML, McCaustland K, Weidlich L, McGee L, Mayer LW, Steigerwalt A, Whaley M, Facklam RR, Fields B, Carlone G, Ades EW, Dagan R, Sampson JS. Evaluation and improvement of real-time PCR assays targeting *lytA*, *ply*, and *psaA* genes for detection of pneumococcal DNA. *J. Clin. Microbiol.* 2007;45(8):2460-6.
2. Azzari C, Moriondo M, Indolfi G, Cortimiglia M, Canessa C, Becciolini L, Lippi F, de Martino M, Resti M. Realtime PCR is more sensitive than multiplex PCR for diagnosis and serotyping in children with culture negative pneumococcal invasive disease. *PLoS One.* 2010;5(2):e9282.
3. World Health Organization [Internet] WHO-UNICEF estimates of PCV3 coverage. 2019 [updated 15 July 2019; cited 23 September 2019]. Available from: http://apps.who.int/immunization_monitoring/globalsummary/timeseries/tswucoveragepcv3.html.
4. von Gottberg A, de Gouveia L, Tempia S, Quan V, Meiring S, von Mollendorf C, Madhi SA, Zell ER, Verani JR, O'Brien KL, Whitney CG, Klugman KP, Cohen C, Investigators G-S. Effects of vaccination on invasive pneumococcal disease in South Africa. *N. Engl. J. Med.* 2014;371(20):1889-99.
5. Whitney CG, Pilishvili T, Farley MM, Schaffner W, Craig AS, Lynfield R, Nyquist A-C, Gershman KA, Vazquez M, Bennett NM, Reingold A, Thomas A, Glode MP, Zell ER, Jorgensen JH, Beall B, Schuchat A. Effectiveness of seven-valent pneumococcal conjugate vaccine against invasive pneumococcal disease: A matched case-control study. *The Lancet.* 2006;368(9546):1495-502.
6. National Institute for Communicable D. Reduction in invasive pneumococcal disease in South Africa, January through July 2020. Available from <https://www.Nicd.Ac.Za/wp-content/uploads/2020/08/pneumococcal-disease.Pdf> accessed 14 September 2020. *Communicable Diseases Communiqué.* 2020;19(8):5.

Last updated: 11 September 2020

Next update: 30 November 2020

Appendix: Cumulative invasive pneumococcal disease case numbers reported by the GERMS-SA surveillance programme, January 2005 to December 2012

GERMS-SA surveillance programme

- GERMS-SA is a national, active, laboratory-based surveillance system initiated in 2003.
- Invasive pneumococcal disease (IPD) cases defined as hospitalised individuals with *Streptococcus pneumoniae* cultured from normally sterile site specimens (e.g. cerebrospinal fluid, blood or joint fluid).
- Repeat isolates from the same individual within 21 days were excluded.
- ~190 laboratories each year send reports and isolates.
- Age, sex, date of specimen collection, and source of specimen were captured.
- Pneumococci were serotyped by Quellung reaction using specific antisera (Statens Serum Institute, Copenhagen, Denmark).
- **Only viable isolates are included in cumulative graph case numbers as molecular diagnostic techniques were only introduced in 2007.**
- Figures 1 – 3 are for cases < 5 years, and Figures 4 – 6 for cases 5 years and older. Cases with unknown age were excluded from the figures.
- There are three graphs for each age group:
 - Disease caused by any of the seven serotypes in PCV7 (4, 6B, 9V, 14, 18C, 19F and 23F)
 - Disease caused by any of the six additional serotypes in PCV13 but not in PCV7 (1, 3, 5, 6A, 7F, 19A)
 - Disease caused by any serotypes not in PCV13
- More information on the GERMS-SA system available at:
<http://www.nicd.ac.za/centres/division-of-public-health-surveillance-and-response/>

PCV vaccine introduction in South Africa

- The 7-valent pneumococcal conjugate vaccine (PCV-7) was introduced to the South African Expanded Programme on Immunization in April 2009, with no catch-up vaccination campaign.
- There was a graded replacement of PCV-7 by 13-valent pneumococcal conjugate (PCV-13) in 2011. By June 2011 all provinces were using PCV-13.
- There was a limited PCV-13 catch-up campaign in 2011 and 2012.
- WHO/UNICEF vaccine coverage estimates for receiving a third dose of the PCV vaccine in South Africa are 10% in 2008, 58% in 2009, 62% in 2011, 75% in 2012, 77% in 2013, 85% in 2014, 85% 2015 and 77% in 2016.¹

The effect of the vaccine on IPD in South Africa has been described.²

Appendix: Cumulative invasive pneumococcal disease case numbers reported by the GERMS-SA surveillance programme, 1 January 2005 to December 2012




Figure 1. Cumulative weekly number of disease episodes of invasive pneumococcal disease due to any of the seven serotypes (4, 6B, 9V, 14, 18C, 19F and 23F) in PCV-7: children <5 years of age in South Africa, from 2005 to 2012. Only viable isolates serotyped using Quellung method included.

Appendix: Cumulative invasive pneumococcal disease case numbers reported by the GERMS-SA surveillance programme, 1 January 2005 to December 2012




Figure 2. Cumulative weekly number of disease episodes of invasive pneumococcal disease due to any of the six additional (1, 3, 5, 6A, 7F, 19A) serotypes in PCV-13 but not in PCV7: children <5 years of age in South Africa, from 2005 to 2012. Only viable isolates serotyped using Quellung method included. (Note: There is reported cross protection between 6A and 6B which is included in PCV7⁵)




Figure 3. Cumulative weekly number of disease episodes of invasive pneumococcal disease due to any of the serotypes not in PCV13: Children <5 years of age in South Africa, from 2005 to 2012. Only viable isolates serotyped using Quellung method included.




Figure 4. Cumulative weekly number of disease episodes of invasive pneumococcal disease due to any of the seven serotypes (4, 6B, 9V, 14, 18C, 19F and 23F) in PCV-7: Individuals ≥ 5 years of age in South Africa, from 2005 to 2012. Only viable isolates serotyped using Quellung method included.




Figure 5: Cumulative weekly number of disease episodes of invasive pneumococcal disease due to any of the six additional (1, 3, 5, 6A, 7F, 19A) serotypes in PCV-13 but not in PCV-7: individuals ≥ 5 years of age in South Africa, from 2005 to 2012. Only viable isolates serotyped using Quelling method included. (Note: There is reported cross protection between 6A and 6B which is included in PCV-7⁵)




Figure 6. Cumulative weekly number of disease episodes of invasive pneumococcal disease due to any of the serotypes not in PCV-13: individuals ≥ 5 years of age in South Africa, from 2005 to 2012. Only viable isolates serotyped using Quellung method included.

Appendix: Cumulative invasive pneumococcal disease case numbers reported by the GERMS-SA surveillance programme, 1 January 2005 to December 2012

Missing information

Table 1. Isolates with missing age; number of viable and non-viable isolates and audit cases identified, 2005-2012

| | Age missing, n (%) | Viable, n (%) | Non-viable, n (%) | Audit, n (%) | Total |
|------|--------------------|---------------|-------------------|--------------|--------|
| 2005 | 236 (5) | 3,650 (75) | 380 (8) | 856 (18) | 4,886 |
| 2006 | 223 (5) | 3,419 (72) | 444 (9) | 868 (18) | 4,731 |
| 2007 | 217 (5) | 3,329 (70) | 597 (13) | 816 (17) | 4,742 |
| 2008 | 208 (4) | 3,327 (69) | 576 (12) | 932 (19) | 4,835 |
| 2009 | 161 (3) | 3,387 (71) | 532 (11) | 841 (18) | 4,760 |
| 2010 | 141 (3) | 2,873 (68) | 515 (12) | 809 (19) | 4,197 |
| 2011 | 218 (6) | 2,409 (63) | 451 (12) | 944 (25) | 3,804 |
| 2012 | 248 (8) | 2,160 (67) | 344 (11) | 718 (22) | 3,222 |
| All | 1,652 (5) | 24,554 (67) | 3,839 (12) | 6,784 (20) | 35,177 |

References

1. World Health Organization [Internet] WHO-UNICEF estimates of PCV3 coverage. 2017 [cited 21 July 2017]. Available from: http://apps.who.int/immunization_monitoring/globalsummary/timeseries/tswucoveragepcv3.html.
2. von Gottberg A, de Gouveia L, Tempia S, Quan V, Meiring S, von Mollendorf C, et al. Effects of vaccination on invasive pneumococcal disease in South Africa. *N. Engl. J. Med.* 2014;371(20):1889-99.