WEEKLY RESPIRATORY PATHOGENS SURVEILLANCE **REPORT**



SOUTH AFRICA WEEK 38 2020

CONTENTS

Programme descriptions	2
Comments	3
Influenza- like illness (ILI) surveillance primary health care clinics	4-7
Influenza-like illness (ILI) surveillance: Viral Watch	8-9
National syndromic surveillance for pneumonia	10-13
Summary of laboratory	14-15

CUMULATIVE DATA FROM



2020

HIGHLIGHTS: WEEK 38

The 2020 influenza season has not yet started. Only one detection of influenza A(H1N1)pdm09 has been made in week 24 (week starting 8 June 2020), since the localised outbreak of influenza A(H1N1) pdm09 and to a lesser extent influenza B(Victoria) in the Western Cape Province, in the first three months of the year.

Although the 2020 RSV season has not started, with only sporadic detections since mid April, there has been a moderate increase in detection since week 29 (week ending 19 July) in the pneumonia surveillance programme and to a lesser extent in the ILI programme.

This week additional 11 patients tested positive for SARS-CoV-2 in SARI (9), ILI (2) and Viral Watch (0). To date, 576 cases have been detected from all surveillance programmes.

WEEK 38 2020 REPORTING PERIOD 01 JANUARY 2020 TO 20 SEPTEMBER 2020

PROGRAMME DESCRIPTIONS

Programme	Influenza-like illness (ILI)	Viral Watch	National syndromic surveillance for pneumonia
Start year	2012	1984	2009
Provinces*	KZ NW WC**	EC FS GP LP MP NC NW	GP KZ MP NW WC
Type of site	Primary health care clinics	General practitioners	Public hospitals
Case definition	An acute respiratory illness with a temperature (≥38°C) and cough, & onset ≤10 days	An acute respiratory illness with a temperature (≥38°C) and cough, & onset ≤10 days	Acute (symptom onset≤10 days) or chronic (symptom onset >10) lower respiratory tract infection
Specimens collected	Oropharyngeal & nasopharyngeal swabs	Throat and/or nasal swabs or Nasopharyngeal swabs	Oropharyngeal & nasopharyngeal swabs
Main pathogens tested***	INF RSV BP SARS-CoV-2****	INF RSV BP SARS-CoV-2****	INF RSV BP SARS-CoV-2****

Epidemic Threshold

Thresholds are calculated using the Moving Epidemic Method (MEM), a sequential analysis using the R Language, available from http://CRAN.R-project.org/web/package=mem, designed to calculate the duration, start and end of the annual influenza epidemic. MEM uses the 40th, 90th and 97.5th percentiles established from available years of historical data to calculate thresholds of activity. Thresholds of activity for influenza and RSV are defined as follows: Below seasonal threshold, Low activity, Moderate activity, High activity, Very high activity. For influenza, thresholds from outpatient influenza like illness (Viral Watch Programme) are used as an indicator of disease transmission in the community and thresholds from pneumonia surveillance are used as an indicator of impact of disease.

^{*} EC: Eastern Cape; FS: Free State; GP: Gauteng; KZ: KwaZulu-Natal; LP: Limpopo; MP: Mpumalanga: NC: Northern Cape; NW: North West; WC: Western Cape

^{**}Started in 2019

^{***}INF: influenza virus; RSV: respiratory syncytial virus; BP: Bordetella pertussis

^{****}SARS-CoV-2: Severe acute respiratory syndrome coronavirus 2

WEEK 38 2020 REPORTING PERIOD 01 JANUARY 2020 TO 20 SEPTEMBER 2020

COMMENTS

Influenza

The 2020 influenza season has not yet started although sustained detections of influenza A(H1N1)pdm09 and to a lesser extent influenza B(Victoria) were made from Western Cape Province, in all surveillance programmes from week 2 to week 15. In week 24 (week ending 14th June), one influenza case was detected in Gauteng province.

ILI programme: In 2020 to date, specimens from 1 094 patients were received from 4 ILI sites. Influenza was detected in 52 specimens (all from Western Cape province), 36 (69%) were identified as influenza A(H1N1) pdm09, one (2%) influenza A subtype inconclusive, 12 (23%) as influenza B (Victoria) and three (6%) B lineage inconclusive (Figure 1 and Table 1).

Viral Watch programme: During the same period, specimens were received from 347 patients from Viral Watch sites in 8 provinces. Influenza was detected in 78 patients, one is an imported case and one is a new case of week 24 since week 14 influenza detections were made. Of the 77 local cases; 75 (98%) were influenza A(H1N1)pdm09, one (1%) influenza A(H3N2) and one (1%) influenza B(Victoria) (Figure 4 and Table 4).

Pneumonia surveillance: Since the beginning of 2020, specimens from 2 952 patients with severe respiratory illness (SRI) were received from the 6 sentinel sites. Influenza was detected in 26 patients (all from Western Cape Province), of which 21 (81%) were influenza A(H1N1) pdm09, one (4%) influenza A subtype inconclusive and four (15%) influenza B(Victoria) (Figure 7 and Table 6).

Respiratory syncytial virus

Although the 2020 RSV season has not started, with only sporadic detections since mid April, there has been a moderate increase in detection since week 29 (week ending 19 July) in the pneumonia surveillance programme and to a lesser extent in the ILI programme.

ILI programme: In 2020 to date, 1 094 specimens were tested and RSV was detected in specimens of 29 (3%) patients.

Viral Watch programme: During the same period, 347 specimens were tested and RSV has not been detected.

Pneumonia surveillance: Since the beginning of 2020, 2 952 specimens were tested and RSV was detected in specimens of 277 (9%) patients.

SARS-CoV-2 (Severe acute respiratory syndrome coronavirus 2)

Testing for SARS-CoV-2 was initiated in all three surveillance programmes in week 10 (week starting 2 March 2020).

ILI programme: In 2020 to date, specimens from 860 patients were tested and SARS-CoV-2 was detected in 125 (15%) patients.


Viral Watch programme: In 2020 to date, specimens were tested from 243 patients and SARS-CoV-2 was detected in 32 (13%) patients.

Pneumonia surveillance: In 2020 to date, specimens from 2 428 patients with severe respiratory illness (SRI) were tested and SARS-CoV-2 was detected in 419 (17%) patients.

WEEK 38 2020 REPORTING PERIOD 01 JANUARY 2020 TO 20 SEPTEMBER 2020

INFLUENZA-LIKE ILLNESS (ILI) SURVEILLANCE PRIMARY HEALTH CARE CLINICS

Figure 1. Number of positive samples* by influenza subtype and lineage and detection rate** by week



^{*}Specimens from patients with influenza-like illnesses at 4 sentinel sites in 3 provinces
**Only reported for weeks with >10 specimens submitted
Inconclusive: insufficient viral load in sample and unable to characterise further

Table 1. Cumulative number of influenza subtype and lineage and total number of samples tested by clinic and province


Clinic (Province)	A(H1N1) pdm09	A(H3N2)	A subtype inconclusive	B/Victoria	B/ Yamagata	B lineage inconclusive	Total samples
Eastridge (WC)	33	0		12	0	3	451
Edendale Gateway (KZ)	0	0	О	0	0	0	136
Jouberton (NW)	0	0	О	0	0	0	236
Mitchell's Plain (WC)	3	0	О	0	0	0	271
Total:	36	0	1	12	0	3	1 094

KZ: KwaZulu-Natal; NW: North West; WC: Western Cape Inconclusive: insufficient viral load in sample and unable to characterise further

WEEK 38 2020 REPORTING PERIOD 01 JANUARY 2020 TO 20 SEPTEMBER 2020

INFLUENZA-LIKE ILLNESS (ILI) SURVEILLANCE PRIMARY HEALTH CARE CLINICS

Figure 2. Number of samples testing positive for respiratory syncytial virus by subgroup and detection rate by week



Inconclusive: insufficient viral load in sample and unable to characterise further

Table 2. Cumulative number of respiratory syncytial virus subgroups identified and total number of samples tested by clinic and province


Clinic (Province)	RSVA	RSVB	RSVAB	RSV subgroup inconclusive	Total samples
Eastridge (WC)	13	3	0	0	451
Edendale Gateway (KZ)		5	0	0	136
Jouberton (NW)		0	0	0	236
Mitchell's Plain (WC)	6	О	0	0	271
Total:	21	8	0	0	1 094

KZ: KwaZulu-Natal; NW: North West; WC: Western Cape Inconclusive: insufficient viral load in sample and unable to characterise further RSV AB: Both RSV A and B subgroup identified

WEEK 38 2020 REPORTING PERIOD 02 MARCH 2020 TO 20 SEPTEMBER 2020

INFLUENZA-LIKE ILLNESS (ILI) SURVEILLANCE PRIMARY HEALTH CARE CLINICS

Figure 3. Number of samples testing positive for SARS-CoV-2*, and detection rate by week



^{*}Specimens from nationts with influenza-like illnesses at 4 sentinel sites in 3 provinces

Table 3. Cumulative number of SARS-CoV-2 identified and total number of samples tested by clinic and province


Clinic (Province)	SARS-CoV-2 positive	Total samples tested
Eastridge (WC)	26	348
Edendale Gateway (KZ)	21	86
Jouberton (NW)	40	187
Mitchell's Plain (WC)	38	239
Total:	125	860

KZ: KwaZulu-Natal; NW: North West; WC: Western Cape

WEEK 38 2020 REPORTING PERIOD 01 JANURY 2020 TO 20 SEPTEMBER 2020

INFLUENZA-LIKE ILLNESS (ILI) SURVEILLANCE VIRAL WATCH

Figure 4. Number of positive samples* by influenza subtype and lineage and detection rate** by week



^{*}Specimens from patients with influenza-like illnesses at 92 sentinel sites in 8 provinces
** Only reported for weeks with >10 specimens submitted.

Table 4. Cumulative number of influenza subtype and lineage and total number of samples tested by province

Province	A(H1N1) pdm09	A(H3N2)	A subtype inconclusive	B/Victoria	B/ Yamagata	B lineage inconclusive	Total samples
Eastern Cape	0	0	0	0	0	0	4
Free State	0	0	0	0	0	0	11
Gauteng	2	0	О	0	0	0	152
Limpopo	0	0	О	О	0	0	3
Mpumalanga	0	0	О	0	0	0	7
North West	0	0	О	0	0	0	0
Northern Cape	0	0	О	0	0	0	4
Western Cape	74	1	0	1	0	0	166
Total:	76	1	0	1	0	0	347

Inconclusive: insufficient viral load in sample and unable to characterise further

From 01 January 2020 to date, 10 patients were tested for influenza at the time of entry into South Africa following travel abroad and influenza was detected in three patients, of which one influenza A(H1N1)pdm09 and two influenza A(H3N2).

Patients known to have acquired influenza abroad are not included in the table or epidemiological curve.

www.nicd.ac.za

TOLL-FREE NUMBER 0800 029 999

PAGE 7

WEEK 38 2020 REPORTING PERIOD 01 JANUARY 2020 TO 20 SEPTEMBER 2020

INFLUENZA-LIKE ILLNESS (ILI) SURVEILLANCE VIRAL WATCH

Figure 5. ILI surveillance (Viral Watch) percentage influenza detections and epidemic thresholds




*Thresholds based on 2010-2019 data

WEEK 38 2020 REPORTING PERIOD 02 MARCH 2020 TO 20 SEPTEMBER 2020

INFLUENZA-LIKE ILLNESS (ILI) SURVEILLANCE: VIRAL WATCH

Figure 6. Number of samples testing positive for SARS-CoV-2*, and detection rate by week



Specimens from patients with Influenza-like illnesses at 92 sentinel sites in 8 provinces


Table 5. Cumulative number of SARS-CoV-2 identified and total number of samples tested by province

Province	SARS-CoV-2 positve	Total samples tested
Eastern Cape	0	3
Free State	0	11
Gauteng	24	141
Limpopo	0	2
Mpumalanga		5
North West	0	0
Northern Cape	0	2
Western Cape	7	79
Total:	32	243

WEEK 38 2020 REPORTING PERIOD 01 JANUARY 2020 TO 20 SEPTEMBER 2020

NATIONAL SYNDROMIC SURVEILLANCE FOR PNEUMONIA

Figure 7. Number of positive samples* by influenza subtype and lineage and detection rate** by week



^{*}Specimens from patients hospitalised with pneumonia at 6 sentinel sites in 5 provinces

Inconclusive: insufficient viral load in sample and unable to characterise further

Table 6. Cumulative number of identified influenza subtype and lineage and total number of samples tested by hospital

Hospital (Province)	A(H1N1) pdm09	A(H3N2)	A subtype inconclusive	B/ Victoria	B/ Yamagata	B lineage inconclusive	Total samples
Edendale (KZ)	0	0	0	0	0	0	484
Helen Joseph-Rahima Moosa (GP)	О	О	О	О	0	0	676
Klerksdorp-Tshepong (NW)	О	0	0	О	0	0	512
Mapulaneng-Matikwana (MP)	О	0	О	О	0	0	213
Red Cross (WC)	19	0	О	2	0	0	745
Mitchell's Plain (WC)	2	0	TO THE L	2	0	0	322
Total:	21	0	1	4	0	0	2 952

 ${\sf GP: Gauteng; KZ: KwaZulu-Natal; NW: North West; MP: Mpumalanga; WC: Western Cape}$


Inconclusive: insufficient viral load in sample and unable to characterise further

^{**}Only reported for weeks with >10 specimens submitted

WEEK 38 2020 REPORTING PERIOD 01 JANUARY 2020 TO 20 SEPTEMBER 2020

NATIONAL SYNDROMIC SURVEILLANCE FOR PNEUMONIA

Figure 8. National syndromic surveillance for pneumonia percentage influenza detections and epidemic thresholds*




*Thresholds based on 2010-2019 data

WEEK 38 2020 REPORTING PERIOD 01 JANUARY 2020 TO 20 SEPTEMBER 2020

NATIONAL SYNDROMIC SURVEILLANCE FOR PNEUMONIA

Figure 9. Number of samples testing positive for respiratory syncytial virus by subgroup and detection rate by week



Inconclusive: insufficient viral load in sample and unable to characterise further

Table 7: Cumulative number of respiratory syncytial virus subgroups identified and total number of samples tested by hospital


Hospital (Province)	RSVA	RSVB	RSVAB	RSV subgroup inconclusive	Total samples
Edendale (KZ)	2	9	О	1	484
Helen Joseph-Rahima Moosa (GP)	39	10	О	0	676
Klerksdorp-Tshepong (NW)	2	0	О	0	512
Mapulaneng-Matikwana (MP)	0	0	О	0	213
Red Cross (WC)	152	26	О	2	745
Mitchell's Plain (WC)	33	0	О	1	322
Total:	228	45	0	4	2 952

GP: Gauteng; KZ: KwaZulu-Natal; NW: North West; MP: Mpumalanga; WC: Western Cape Inconclusive: insufficient viral load in sample and unable to characterise further RSV AB: Both RSV A and B subgroup identified

WEEK 38 2020 REPORTING PERIOD 02 MARCH 2020 TO 20 SEPTEMBER 2020

NATIONAL SYNDROMIC SURVEILLANCE FOR PNEUMONIA

Figure 10. Number of samples testing positive for SARS-CoV-2*, and detection rate by week



^{*}Specimens from patients with pneumonia at 6 sentinel sites in 5 provinces

Table 8. Cumulative number of identified SARS-CoV-2 and total number of samples tested by hospital

Hospital (Province)	SARS-CoV-2 positve	Total samples tested
Edendale (KZ)	93	408
Helen Joseph-Rahima Moosa (GP)	117	541
Klerksdorp-Tshepong (NW)	122	437
Mapulaneng-Matikwana (MP)	9	164
Red Cross (WC)	27	603
Mitchell's Plain (WC)	51	275
Total:	419	2 428

GP: Gauteng: KZ: KwaZulu-Natal: NW: North West: MP: Mpumalanga: WC: Western Cape

WEEK 38 2020 REPORTING PERIOD 02 MARCH 2020 TO 20 SEPTEMBER 2020

SUMMARY OF LABORATORY-CONFIRMED SARS-CoV-2 CASES

Table 9. Characteristics of laboratory-confirmed cases of COVID-19, enrolled in influenza-like illness (ILI) and pneumonia surveillance programmes, South Africa, 2 March - 20 September 2020

Characteristic		Influenza -like illness (ILI), public sector, n =125 (%)	Pneumonia surveillance, n=419 (%)
Age group			
0-9		21/125 (17)	34/419 (8)
10-19		12/125 (10)	4/419 (1)
20-29		60/125 (48)	83/419 (20)
40-59)	23/125 (18)	160/419 (38)
60-79		9/125 (7)	130/419 (31)
≥80		0/125 (0)	8/419 (2)
Sex-female		65/125 (52)	258/419 (62)
Province*			
Gau	teng	N/A	117/419 (28)
KZN		21/125 (17)	93/419 (22)
Мри	ımalanga	N/A	9/419 (2)
Nort	:h West	40/125 (32)	122/419 (29)
Wes	stern Cape	64/125 (51)	78/419 (19)
Race			
Black		76/121 (63)	328/408 (80)
Colo	ured	45/121 (37)	58/408 (14)
Asiar	n/Indian	0/121 (0)	18/408 (4)
Othe		0/121 (0)	4/408 (1)
Presentation			
Feve		119/121 (98)	245/408 (60)
Coug	h	120/121 (99)	406/408 (99)
Shortness o	f breath	25/121 (21)	327/408 (80)
Tight ch	est	39/121 (32)	199/408 (49)
Diarrho	ea	13/121 (11)	23/408 (6)

WEEK 38 2020 REPORTING PERIOD 02 MARCH 2020 TO 20 SEPTEMBER 2020

SUMMARY OF LABORATORY-CONFIRMED SARS-CoV-2 CASES

Characteristic	Influenza -like illness (ILI), public sector, n =125 (%)	Pneumonia surveillance, n=419 (%)
Underlying conditions		
Hypertension	8/121 (7)	53/403 (13)
Cardiac	0/121 (0)	8/403 (2)
Respiratory	1/121 (1)	17/403 (4)
Diabetes	1/121 (1)	92/403 (23)
Cancer	0/121 (0)	1/403 (<1)
Tuberculosis	0/121 (0)	13/403 (3)
HIV-infection	14/121 (12)	83/403 (21)
Other **	4/121 (3)	70/403 (17)
Management		
Oxygen therapy	1/121 (1)	259/403 (64)
ICU admission	N/A	16/403 (4)
Ventilation	N/A	13/403 (3)
Outcome***		
Died	0/121 (0)	50/403 (12)

 $[^]st$ ILI surveillance not conducted in Gauteng & Mpumalanga provinces

Note: Children may be over-represented amongst hospitalised patients due to the inclusion of a large paediatric hospital in Cape Town.

Of the 50 patients who died, six were in the 20-39 year age group, 17 in the 40-59 year age group, and 27 were ≥60 years; 33/50 (66%) were female. All except four were known to have underlying medical conditions



www.nicd.ac.za

TOLL-FREE NUMBER 0800 029 999

PAGE **15**

^{**} Chronic lung, liver and kidney disease, organ transplant, pregnancy, malnutrition, obesity, tracheostomy, prematurity, seizure, stroke, anaemia, asplenia, burns, Systemic lupus erythematosus, seizures

 $[\]hbox{\it ****} \ {\tt Outcome} \ {\tt includes} \ {\tt patients} \ {\tt who} \ {\tt are} \ {\tt still} \ {\tt hospitalised}, have been \ {\tt discharged} \ {\tt or} \ {\tt referred}, and \ {\tt those} \ {\tt who} \ {\tt or} \ {\tt o$