WEEKLY RESPIRATORY PATHOGENS SURVEILLANCE **REPORT**



SOUTH AFRICA WEEK 41 2020

CONTENTS

Programme descriptions	2
Comments	3
Influenza- like illness (ILI) surveillance primary health care clinics	4-7
Influenza-like illness (ILI) surveillance: Viral Watch	8-9
National syndromic surveillance for pneumonia	10-13
Summary of laboratory confirmed SARS-CoV-2 cases	14-15

CUMULATIVE DATA FROM





HIGHLIGHTS: WEEK 41

The 2020 influenza season has not yet started. Only one detection of influenza A(H1N1)pdm09 has been made in week 24 (week starting 8 June 2020), since the localised outbreak of influenza A(H1N1) pdm09 and to a lesser extent influenza B(Victoria) in the Western Cape Province, in the first three months of the year.

Although the 2020 RSV season has not started, with only sporadic detections since mid April, there has been a moderate increase in detection since week 29 (week ending 19 July) in the pneumonia surveillance programme and to a lesser extent in the ILI programme.

This the last report 14 additional patients tested positive for SARS-CoV-2 of which 7 were detected in the current reporting week (week41) 4 in SARI and 3 in ILI surveillance. To date, 614 cases have been detected from all surveillance programmes. Of the 432 hospitalised COVID-19 cases with available data on outcome, 56 (13%) died.

WEEK 41 2020 REPORTING PERIOD 01 JANUARY 2020 TO 11 OCTOBER 2020

PROGRAMME DESCRIPTIONS

Programme	Influenza-like illness (ILI)	Viral Watch	National syndromic surveillance for pneumonia
Start year	2012	1984	2009
Provinces*	KZ NW WC**	EC FS GP LP MP NC NW	GP KZ MP NW WC
Type of site	Primary health care clinics	General practitioners	Public hospitals
Case definition	An acute respiratory illness with a temperature (≥38°C) and cough, & onset ≤10 days	An acute respiratory illness with a temperature (≥38°C) and cough, & onset ≤10 days	Acute (symptom onset≤10 days) or chronic (symptom onset >10) lower respiratory tract infection
Specimens collected	Oropharyngeal & nasopharyngeal swabs	Throat and/or nasal swabs or Nasopharyngeal swabs	Oropharyngeal & nasopharyngeal swabs
Main pathogens tested***	INF RSV BP SARS-CoV-2****	INF RSV BP SARS-CoV-2****	INF RSV BP SARS-CoV-2****

Epidemic Threshold

Thresholds are calculated using the Moving Epidemic Method (MEM), a sequential analysis using the R Language, available from http://CRAN.R-project.org/web/package=mem, designed to calculate the duration, start and end of the annual influenza epidemic. MEM uses the 40th, 90th and 97.5th percentiles established from available years of historical data to calculate thresholds of activity. Thresholds of activity for influenza and RSV are defined as follows: Below seasonal threshold, Low activity, Moderate activity, High activity, Very high activity. For influenza, thresholds from outpatient influenza like illness (Viral Watch Programme) are used as an indicator of disease transmission in the community and thresholds from pneumonia surveillance are used as an indicator of impact of disease.

^{*} EC: Eastern Cape; FS: Free State; GP: Gauteng; KZ: KwaZulu-Natal; LP: Limpopo; MP: Mpumalanga: NC: Northern Cape; NW: North West; WC: Western Cape

^{**}Started in 2019

^{***}INF: influenza virus; RSV: respiratory syncytial virus; BP: Bordetella pertussis

^{****}SARS-CoV-2: Severe acute respiratory syndrome coronavirus 2

WEEK 41 2020 REPORTING PERIOD 01 JANUARY 2020 TO 11 OCTOBER 2020

COMMENTS

Influenza

The 2020 influenza season has not yet started although sustained detections of influenza A(H1N1)pdm09 and to a lesser extent influenza B(Victoria) were made from Western Cape Province, in all surveillance programmes from week 2 to week 15. In week 24 (week ending 14th June), one influenza case was detected in Gauteng province.

ILI programme: In 2020 to date, specimens from 1 183 patients were received from 4 ILI sites. Influenza was detected in 52 specimens (all from Western Cape province), 36 (69%) were identified as influenza A(H1N1) pdm09, one (2%) influenza A subtype inconclusive, 12 (23%) as influenza B (Victoria) and three (6%) B lineage inconclusive (Figure 1 and Table 1).

Viral Watch programme: During the same period, specimens were received from 357 patients from Viral Watch sites in 8 provinces. Influenza was detected in 78 patients and five are imported cases detected in the first weeks of the year. Of the 73 local cases, all were influenza A(H1N1)pdm09. (Figure4 and Table4).

Pneumonia surveillance: Since the beginning of 2020, specimens from 3 238 patients with severe respiratory illness (SRI) were received from the 6 sentinel sites. Influenza was detected in 26 patients (all from Western Cape Province), of which 21 (81%) were influenza A(H1N1) pdm09, one (4%) influenza A subtype inconclusive and four (15%) influenza B(Victoria) (Figure 7 and Table 6).

Respiratory syncytial virus

Although the 2020 RSV season has not started, with only sporadic detections since mid April, there has been a moderate increase in detection since week 29 (week ending 19 July) in the pneumonia surveillance programme and to a lesser extent in the ILI programme.

ILI programme: In 2020 to date, 1 183 specimens were tested and RSV was detected in specimens of 48 (4%) patients

Viral Watch programme: During the same period, 357 specimens were tested and RSV has not been detected.

Pneumonia surveillance: Since the beginning of 2020, 3 238 specimens were tested and RSV was detected in specimens of 340 (11%) patients.

SARS-CoV-2 (Severe acute respiratory syndrome coronavirus 2)

Testing for SARS-CoV-2 was initiated in all three surveillance programmes in week 10 (week starting 2 March 2020

ILI programme: In 2020 to date, specimens from 949 patients were tested and SARS-CoV-2 was detected in 135 (14%) patients.


Viral Watch programme: In 2020 to date, specimens were tested from 253 patients and SARS-CoV-2 was detected in 35 (14%) patients

Pneumonia surveillance: In 2020 to date, specimens from 2 736 patients with severe respiratory illness (SRI) were tested and SARS-CoV-2 was detected in 444 (16%) patients.

WEEK 41 2020 REPORTING PERIOD 01 JANUARY 2020 TO 11 OCTOBER 2020

INFLUENZA-LIKE ILLNESS (ILI) SURVEILLANCE PRIMARY HEALTH CARE CLINICS

Figure 1. Number of positive samples* by influenza subtype and lineage and detection rate** by week



^{*}Specimens from patients with influenza-like illnesses at 4 sentinel sites in 3 provinces
**Only reported for weeks with >10 specimens submitted
Inconclusive: insufficient viral load in sample and unable to characterise further

Table 1. Cumulative number of influenza subtype and lineage and total number of samples tested by clinic and province


Clinic (Province)	A(H1N1) pdm09	A(H3N2)	A subtype inconclusive	B/Victoria	B/ Yamagata	B lineage inconclusive	Total samples
Eastridge (WC)	33	0		12	0	3	484
Edendale Gateway (KZ)	0	0	О	0	0	0	142
Jouberton (NW)	0	0	О	0	0	0	271
Mitchell's Plain (WC)	3	0	О	0	0	0	286
Total:	36	0	1	12	0	3	1 183

KZ: KwaZulu-Natal; NW: North West; WC: Western Cape Inconclusive: insufficient viral load in sample and unable to characterise further

WEEK 41 2020 REPORTING PERIOD 01 JANUARY 2020 TO 11 OCTOBER 2020

INFLUENZA-LIKE ILLNESS (ILI) SURVEILLANCE PRIMARY HEALTH CARE CLINICS

Figure 2. Number of samples testing positive for respiratory syncytial virus by subgroup and detection rate by week



Inconclusive: insufficient viral load in sample and unable to characterise further

Table 2. Cumulative number of respiratory syncytial virus subgroups identified and total number of samples tested by clinic and province


Clinic (Province)	RSVA	RSVB	RSVAB	RSV subgroup inconclusive	Total samples
Eastridge (WC)	26	2	0		484
Edendale Gateway (KZ)		5	0	0	142
Jouberton (NW)		0	0	2	271
Mitchell's Plain (WC)	10	0	0	0	286
Total	38	7	0	3	1 183

KZ: KwaZulu-Natal; NW: North West; WC: Western Cape Inconclusive: insufficient viral load in sample and unable to characterise further RSV AB: Both RSV A and B subgroup identified

WEEK 41 2020 REPORTING PERIOD 02 MARCH 2020 TO 11 OCTOBER 2020

INFLUENZA-LIKE ILLNESS (ILI) SURVEILLANCE PRIMARY HEALTH CARE CLINICS

Figure 3. Number of samples testing positive for SARS-CoV-2*, and detection rate by week



^{*}Specimens from natients with influenza-like illnesses at 4 sentinel sites in 3 provinces

Table 3. Cumulative number of SARS-CoV-2 identified and total number of samples tested by clinic and province


Clinic (Province)	SARS-CoV-2 positive	Total samples tested
Eastridge (WC)	26	381
Edendale Gateway (KZ)	21	92
Jouberton (NW)	50	222
Mitchell's Plain (WC)	38	254
Total:	135	949

KZ: KwaZulu-Natal; NW: North West; WC: Western Cape

WEEK 41 2020 REPORTING PERIOD 01 JANURY 2020 TO 11 OCTOBER 2020

INFLUENZA-LIKE ILLNESS (ILI) SURVEILLANCE VIRAL WATCH

Figure 4. Number of positive samples* by influenza subtype and lineage and detection rate** by week



^{*}Specimens from patients with influenza-like illnesses at 92 sentinel sites in 8 provinces
** Only reported for weeks with >10 specimens submitted.

Table 4. Cumulative number of influenza subtype and lineage and total number of samples tested by province

Province	A(H1N1) pdm09	A(H3N2)	A subtype inconclusive	B/Victoria	B/ Yamagata	B lineage inconclusive	Total samples
Eastern Cape	0	0	0	0	0	0	4
Free State	0	0	0	0	0	0	13
Gauteng	2	0	О	0	0	0	156
Limpopo	0	0	О	О	0	0	3
Mpumalanga	0	0	О	0	0	0	7
North West	0	0	О	0	0	0	0
Northern Cape	0	0	О	0	0	0	4
Western Cape	74	1	0	1	0	0	170
Total:	76	1	0	1	0	0	357

Inconclusive: insufficient viral load in sample and unable to characterise further

From 01 January 2020 to date, 10 patients were tested for influenza at the time of entry into South Africa following travel abroad and influenza was detected in three patients, of which one influenza A(H1N1)pdm09 and two influenza A(H3N2).

Patients known to have acquired influenza abroad are not included in the table or epidemiological curve.

www.nicd.ac.za

TOLL-FREE NUMBER 0800 029 999

PAGE 7

WEEK 41 2020 REPORTING PERIOD 01 JANUARY 2020 TO 11 OCTOBER 2020

INFLUENZA-LIKE ILLNESS (ILI) SURVEILLANCE VIRAL WATCH

Figure 5. ILI (Viral Watch) percentage influenza detections and epidemic thresholds*




*Thresholds based on 2010-2019 data

WEEK 41 2020 REPORTING PERIOD 02 MARCH 2020 TO 11 OCTOBER 2020

INFLUENZA-LIKE ILLNESS (ILI) SURVEILLANCE: VIRAL WATCH

Figure 6. Number of samples testing positive for SARS-CoV-2*, and detection rate by week



Specimens from patients with Influenza-like illnesses at 92 sentinel sites in 8 provinces


Table 5. Cumulative number of SARS-CoV-2 identified and total number of samples tested by province

Province	SARS-CoV-2 positve	Total samples tested
Eastern Cape	0	3
Free State		13
Gauteng	25	145
Limpopo	0	2
Mpumalanga		5
North West	0	0
Northern Cape	0	2
Western Cape	8	83
Total:	35	253

WEEK 41 2020 REPORTING PERIOD 01 JANUARY 2020 TO 11 OCTOBER 2020

NATIONAL SYNDROMIC SURVEILLANCE FOR PNEUMONIA

Figure 7. Number of positive samples* by influenza subtype and lineage and detection rate** by week



^{*}Specimens from patients hospitalised with pneumonia at 6 sentinel sites in 5 provinces

Inconclusive: insufficient viral load in sample and unable to characterise further

Table 6. Cumulative number of identified influenza subtype and lineage and total number of samples tested by hospital

Hospital (Province)	A(H1N1) pdm09	A(H3N2)	A subtype inconclusive	B/ Victoria	B/ Yamagata	B lineage inconclusive	Total samples
Edendale (KZ)	О	0	О	О	0	0	541
Helen Joseph-Rahima Moosa (GP)	0	Ο	О	О	0	0	730
Klerksdorp-Tshepong (NW)	0	0	0	О	0	0	556
Mapulaneng-Matikwana (MP)	0	0	О	Ο	0	0	253
Red Cross (WC)	19	0	0	2	0	0	789
Mitchell's Plain (WC)	2	0	To I do	2	0	0	369
Total:	21	0	1	4	0	0	3 238

 ${\sf GP: Gauteng; KZ: KwaZulu-Natal; NW: North\ West;\ MP:\ Mpumalanga;\ WC:\ Western\ Cape}$

Inconclusive: insufficient viral load in sample and unable to characterise further

^{**}Only reported for weeks with >10 specimens submitted

WEEK 41 2020 REPORTING PERIOD 01 JANUARY 2020 TO 11 OCTOBER 2020

NATIONAL SYNDROMIC SURVEILLANCE FOR PNEUMONIA

Figure 8. National syndromic surveillance for pneumonia percentage influenza detections and epidemic thresholds*




*Thresholds based on 2010-2019 data

WEEK 41 2020 REPORTING PERIOD 01 JANUARY 2020 TO 11 OCTOBER 2020

NATIONAL SYNDROMIC SURVEILLANCE FOR PNEUMONIA

Figure 9. Number of samples testing positive for respiratory syncytial virus by subgroup and detection rate by week



Inconclusive: insufficient viral load in sample and unable to characterise further

Table 7: Cumulative number of respiratory syncytial virus subgroups identified and total number of samples tested by hospital


RSVA	RSVB	RSVAB	RSV subgroup inconclusive	Total samples
2	9	0	1	541
42	10	О	0	730
2	О	О		556
0	О	О	0	253
179	25	0	6	789
58	0	0	5	369
283	45	0	13	3 238
	2 42 2 0 179 58	2 9 42 10 2 0 0 0 179 25 58 0	2 9 0 42 10 0 2 0 0 0 0 0 179 25 0 58 0 0	2 9 0 1 42 10 0 0 2 0 0 1 0 0 0 0 179 25 0 6 58 0 0 5

GP: Gauteng; KZ: KwaZulu-Natal; NW: North West; MP: Mpumalanga; WC: Western Cape Inconclusive: insufficient viral load in sample and unable to characterise further RSV AB: Both RSV A and B subgroup identified

WEEK 41 2020 REPORTING PERIOD 02 MARCH 2020 TO 11 OCTOBER 2020

NATIONAL SYNDROMIC SURVEILLANCE FOR PNEUMONIA

Figure 10. Number of samples testing positive for SARS-CoV-2*, and detection rate by week



^{*}Specimens from patients with pneumonia at 6 sentinel sites in 5 provinces

Table 8. Cumulative number of identified SARS-CoV-2 and total number of samples tested by hospital

Hospital (Province)	SARS-CoV-2 positve	Total samples tested
Edendale (KZ)	96	466
Helen Joseph-Rahima Moosa (GP)	122	595
Klerksdorp-Tshepong (NW)	128	481
Mapulaneng-Matikwana (MP)	11	204
Red Cross (WC)	27	647
Mitchell's Plain (WC)	60	343
Total:	444	2 736

GP: Gauteng: KZ: KwaZulu-Natal: NW: North West: MP: Mpumalanga: WC: Western Cape

WEEK 41 2020 REPORTING PERIOD 02 MARCH 2020 TO 11 OCTOBER 2020

SUMMARY OF LABORATORY-CONFIRMED SARS-CoV-2 CASES

Table 9. Characteristics of laboratory-confirmed cases of COVID-19, enrolled in influenza-like illness (ILI) and pneumonia surveillance programmes, South Africa, 2 March - 20 September 2020

Characteristic	Influenza -like illness (ILI), public sector, n =135 (%)	Pneumonia surveillance, n=444 (%)
Age group		
0-9	21/135 (16)	36/444 (8)
10-19	12/135 (9)	4/444 (1)
20-39	64/135 (47)	87/444 (20)
40-59	28/135 (21)	170/444 (38)
60-79	10/135 (7)	139/444 (31)
≥80	0/135 (0)	8/444 (2)
Sex-female	70/135 (52)	270/444 (61)
Province*		
Gauteng	N/A	122/444 (27)
KZN	21/135 (16)	96/444 (22)
Mpumalanga	N/A	11/444 (2)
North West	50/135 (37)	128/444 (29)
Western Cape	64/135 (47)	87/444 (20)
Race		
Black	86/131 (66)	354/436 (81)
Coloured	45/131 (34)	60/436 (14)
Asian/Indian	0/131 (0)	18/436 (4)
Other	0/131 (0)	4/436 (1)
Presentation		
Fever	128/131 (98)	264/436 (61)
Cough	130/131 (99)	432/436 (99)
Shortness of brea	ath 29/131 (22)	352/436 (81)
Tight chest	44/131 (34)	208/436 (48)
Diarrhoea	17/131 (13)	29/436 (7)

WEEK 41 2020 REPORTING PERIOD 02 MARCH 2020 TO 11 OCTOBER 2020

SUMMARY OF LABORATORY-CONFIRMED SARS-CoV-2 CASES

Characteristic		Influenza -like illness (ILI), public sector, n =135 (%)	Pneumonia surveillance, n=444 (%)
Underlying conditi	ions		
Нур	pertension	9/131 (7)	56/432 (13)
Car	diac	0/131 (0)	8/432 (2)
Res	spiratory	0/131 (0)	1/432 (<1)
Dia	betes	1/131 (<1)	97/432 (22)
Car	ncer	0/131 (0)	1/432 (<1)
Tub	perculosis	0/131 (0)	13/432 (3)
HI√	/-infection	15/131 (11)	90/432 (21)
Oth	ner **	4/131 (3)	72/432 (17)
Management			
Оху	gen therapy	1/131 (1)	281/432 (65)
ICU	admission	N/A	16/432 (4)
Ven	ntilation	N/A	13/432 (3)
Outcome***			
Die	d	0/131 (0)	56/432 (13)

^{*} ILI surveillance not conducted in Gauteng & Mpumalanga provinces

Note: Children may be over-represented amongst hospitalised patients due to the inclusion of a large paediatric hospital in Cape Town.

Of the 56 patients who died, six were in the 20-39 year age group, 19 in the 40-59 year age group, and 31 were ≥60 years; 37/56 (66%) were female. All except four were known to have underlying medical conditions.



^{**} Chronic lung, liver and kidney disease, organ transplant, pregnancy, malnutrition, obesity, tracheostomy, prematurity, seizure, stroke, anaemia, asplenia, burns, Systemic lupus erythematosus, seizures

^{***} Outcome includes patients who are still hospitalised, have been discharged or referred, and those who died