



NICD HANDBOOK

Title: NICD Laboratory Handbook

Document number: NIC0104

Version number: 18

Written by: NICD QA Steering Group

Checked by: Henry Julius and Beverley Singh

Approved by: Adrian Puren

Active date: 08-09-2020

<i>Date of next review</i>	<i>Date reviewed</i>	<i>Reviewed by</i>	<i>Action</i>
08-09-2022			

Date withdrawn:

LABORATORY TEST INFORMATION HANDBOOK

Table of Contents

Services Available 4

Purpose: 4

Scope: 4

Location of the NICD 4

Hotline: 5

Results:..... 5

Abbreviations 5

Help Us to Help You 6

Complaints and Queries..... 7

NOTIFIABLE MEDICAL CONDITION (NMC) 7

The NICD Testing Centres / Laboratories 8

Alphabetical order of tests offered at the NICD 10

Notifiable Medical Conditions (NMC) Disease List 18

CATEGORY 1 NOTIFIABLE MEDICAL CONDITIONS – 18

CATEGORY 2 NOTIFIABLE MEDICAL CONDITIONS – 19

CATEGORY 3 NOTIFIABLE MEDICAL CONDITIONS – 19

CATEGORY 4 NOTIFIABLE MEDICAL CONDITIONS – 20

Tests offered at the NICD per Centre 21

Centre for Vaccines and Immunology (CVI) 21

Centre for Tuberculosis.....	25
Centre for HIV and STI	29
Centre for Emerging Zoonotic and Parasitic Diseases	39
Centre for Enteric Diseases	51
Centre for Healthcare-Associated Infections, Antimicrobial Resistance and Mycoses.....	56
Centre for Respiratory Diseases and Meningitis.....	59
Public Health, Surveillance and Response.....	65
Outbreak Response Unit.....	65
GERMS-SA	65
GERMS-SA clinic-based surveillance (STI, HIV and TB)	66
Travel Health:	66
NICD Sequencing Core Facility	67
Key Contact Staff – National Institute for Communicable Diseases.....	69

LABORATORY TEST INFORMATION HANDBOOK

Services Available

Purpose:

The National Institute for Communicable Diseases (NICD) is a communicable disease (medical microbiology/virology) surveillance facility and laboratory and falls within the control of the National Health Service (NHLS).

The NICD has been established to function as a public health oriented, laboratory-based, national facility distinct from and independent of the existing microbiology/virology laboratories attached to academic centres throughout the country. The direction the NICD takes, is that of public health orientation, rather than a patient oriented clinical diagnostic entity and this is reflected in the service commitments and research directions carried out by the organization. The NICD is, to a large extent, modelled on internationally recognized public health laboratories such as the Centre for Disease Control & Prevention of the USA and Public Health England (former Health Protection Agency).

Scope:

To provide a guide to services offered at the NICD.

Location of the NICD

The NICD is a permanent laboratory and is located at: 1 Modderfontein Road, Sandringham, 2192.

Advice:

For medical advice contact the consultants from the relevant Centre.

For medical advice on investigation of haemorrhagic fevers contact the Medical Consultant to the Centre for Emerging Zoonotic and Parasitic diseases (contact details provided in Table)

Working Hours:

The laboratories at the NICD operate Monday to Friday from 07h30 to 16h30. For after-hour service, please contact the consultant. Please see Key Contact Staff Table

Special Pathogens Laboratory – for after-hour service, please contact the Medical consultant on call (or NICD Call/Emergency Phone number) or the Laboratory directly (contact details provided in Table).

Hotline:

For outbreak, clinical advice or public health queries:

NICD Clinical Advice Hotline +27 82 883 9920

Results:

Results queries please contact the Results Call Centre

011 386 6404

011 386 6314

011 386 6466

Fax 011 386 6342

Abbreviations

Abbreviation	Descriptions
BAL	Bronchoalveolar lavage
CDC	Centre for Disease Control and Prevention
CIF's	Case investigation forms
CMV	Cytomegalovirus
CSF	Cerebrospinal Fluid
ESKAPE	<i>Enterococcus faecium, Staphylococcus aureus, Klebsiella pneumonia, Acinetobacter baumannii, Pseudomonas auruginosa, E. coli and Enterobacter cloacae.</i>
GERMS-SA	Group for Enteric, Respiratory and Meningeal Surveillance in South Africa
HPV	Human papillomavirus
HSV	Herpes Simplex Virus
LGV	Lymphogranuloma venereum
LIMS	Laboratory Information Management System
ILI	Influenza like Illness
NICD	National Institute for Communicable Diseases
NS	Nasal swab
NPA	Nasopharyngeal aspirate
NPS	Nasopharyngeal swab
OPA	Oropharyngeal aspirate
OPS	Oropharyngeal swab
RSVP	Respiratory PCR test
SADC	Southern African Development Community
TA	Throat Aspirate

Abbreviation	Descriptions
TAT	Turnaround Time
TP	<i>Treponema pallidum</i>
TS	Throat swab
TV	<i>Trichomonas vaginalis</i>
UTM	Universal Transport medium
URTI	Upper respiratory Tract Infection
VHF	Viral Hemorrhagic Fever
VTM	Viral transport medium
WHO	World Health Organization

Help Us to Help You

1. All specimens must be accompanied by the following clearly legible data:
 - a) Patient name and hospital number (or referring laboratory reference number).
 - b) Name and address of requesting clinician.
 - c) Nature of specimen (specimen type)
 - d) Collection date and time (where relevant).
 - e) Tests requested.
 - f) Brief clinical history.
 - g) Case investigation forms (CIFs) to be completed for Polio and Measles surveillance, and VHF and rabies request (available on NICD webpage, A-Z disease list).
 - h) Contact number/email for providing the laboratory report
2. Hard copies of all reports are sent to the requesting clinician by post or electronically for referred samples.
3. Reports can be telefaxed to “safe haven” fax machines on request. Please ensure that where required, this service is clearly requested and that a suitable fax number and contact person is supplied.

Where possible, the sender of category “A” specimens must notify the NICD testing laboratory via the NICD Hotline or the provided telephone numbers of the shipment of such samples prior to its arrival at the NICD.

Complaints and Queries

1. Complaints and queries of a minor nature should be addressed to the Laboratory concerned. (See Key Contact Staff Table)
2. Complaints and queries of a serious nature should be addressed to the head of the relevant Laboratory (See Key Contact Staff Table)

Please note that the TAT (Turnaround Time) quoted in this Handbook is calculated from the sample reception at the NICD until a report is issued.

NOTIFIABLE MEDICAL CONDITION (NMC)

Notifiable Medical Conditions are diseases that are of public health importance because they pose significant public health risks that can result in disease outbreaks or epidemics with high case fatality rates both nationally and internationally.

FOR ANY NMC RELATED QUESTIONS/QUERIES/CONCERNS?

NMC helpline: 072 621 3805

Email: NMCsurveillanceReport@nicd.ac.za

Fax: 086 639 1638

Website: www.nicd.ac.za

The NICD Testing Centres / Laboratories

Centre /Laboratories/ Departments	Laboratory	Contact Number
Centre Enteric Diseases (CED)	CED Bacteriology	011 555-0333 /0334
	CED Virology	011 555-0370
Centre for Emerging Zoonotic and Parasitic Diseases (CEZPD)	Electron Microscopy Laboratory	011 386 6318
	Special Bacterial Pathogens Reference Laboratory (SBPRL)	011 555 0331
	Arbovirus Reference Lab	011 386 6424/6353
	Special Viral Pathogens Laboratory (SVPL)	011 386 6376
	Parasitology Reference Laboratory (PRL)	011 555 0311
	Vector control Reference Laboratory (VCRL)	011 386-6480
Centre for HIV & STI's (CHIVSTI)	Cell Biology Unit	011 386-6366
	HIV Virology	011 386 6341
	HIV Surveillance	011 386 6462
	STI Surveillance	011 555 0461
Centre for Vaccines and Immunology (CVI)	Hepatitis	011 386-6387
	Serology and Immunology Unit	011 386-6422
	Measles	011 386 6343
	Polio Isolation and Culture Laboratory	011 555-0504
Centre for Healthcare-Associated Infections, Antimicrobial Resistance and Mycoses (CHARM)	Antimicrobial Resistance Laboratory-Culture Collection (AMRL-CC)	011 555 0342
	Mycology Reference Laboratory (MRL)	011 555 0325
Centre for Respiratory Diseases & Meningitis (CRDM_	GERMS-SA/microbiology	011 555 0315/7
	Molecular	011 555 0356/0387/0352
	Virus isolation and serology	011 386 6390/6373
Centre for Tuberculosis (CTB)	TB Focus Programme & National TB Reference Laboratory	011 885 5316 /5317

Centre /Laboratories/ Departments	Laboratory	Contact Number
Sequencing Core Facility	Sequencing Core Facility	0113866459
Division of Public Health, Surveillance and Response	Notifiable Medical Conditions (NMC)	072 621 3805
	(Outbreak response Unit) Dr Kerrigan Mc Carthy	079 8717278 +27 11 555 0542
	24/7 Doctor on call/Outbreak Hotline	082 883 9920

Alphabetical order of tests offered at the NICD

Pathogen / Test Species Name (disease/ Syndrome)	Pathogen Species Name (disease/ syndrome, ICD-code)	Tests To Request	Department and Contact telephone no's
<i>Arbovirus</i>	Arbovirus: (Chikungunya, Dengue, West Nile, Rift Valley fever, Sindbis, Zika)	Arbovirus (Routine arbo screen, include serology for Chikungunya, West Nile, Sindbis, Rift Valley fever) Dengue and Zika for patient with travel history. PCR/Isolation for Acute cases only Arbovirus PCR (specify which arbovirus, for example "Dengue PCR") Arbovirus isolation Disease specific IgM or IgG ELISA	Arbovirus Reference Laboratory Dr O Hellferscee 011 386 6424/6353 082 908 8045
<i>Adenovirus</i>	Adenovirus Gastroenteritis: Diarrhoea, vomiting, fever	Enteric Adenovirus detection	CED Virology 011 555 0370
Adenovirus	Adenovirus	RSVP	CRDM 011 555 0488 011 386 6373
<i>Anthrax (ANTHR)</i>	<i>Bacillus anthracis</i> (anthrax)	Anthrax (ANTHR)	Special Bacterial Pathogens Reference Laboratory 011 555 0331 011 555 0306
Arthropods/ Insects	Arthropods/ Insects	Morphological identification, PCR identification and/or ELISA	Vector Control Reference Laboratory 011 386 6480
<i>Astrovirus</i>	<i>Astrovirus</i> Gastroenteritis: Diarrhoea, vomiting, fever	<i>Astrovirus</i> detection	CED Virology 011 555 0370
Atypical pneumonia-causing pathogens	<i>M. pneumoniae</i> , <i>C. pneumoniae</i> , <i>Legionella</i> spp.	Atypical pneumonia pathogens (<i>M. pneumoniae</i> , <i>C. pneumoniae</i> , <i>Legionella</i> spp.)	CRDM Bacteriology 011 555 0315/011 555 0317
Avian Influenza (H5, H7, H9)	Avian Influenza (H5, H7, H9)	Suspected Avian influenza	Hotline-Dr on call:082 883 9920 CRDM 011 386 6390 (HOD) 011 555 0488 /011 386 6373
Bacterial meningitis screen	<i>N. meningitidis</i> , <i>H. influenzae</i> and <i>S. pneumoniae</i> , Group B <i>Streptococcus</i> , <i>L. monocytogenes</i> , <i>S. aureus</i> , <i>E.coli</i>	Phenotypic identification and serotyping/grouping MIC Antigen detection PCR for identification and serotyping/ grouping (S.pn, Nm, Hi)	CRDM 011 555 0315/011 555 0317
<i>Botulinum</i>	<i>Clostridium botulinum</i> (botulism)	Botulinum	Special Bacterial Pathogens Reference Laboratory 011 555 0331/011 555 0306
<i>Campylobacter</i> spp	<i>Campylobacter</i> spp	Enteric surveillance and reference function	CED Bacteriology 011 386 6235/011 555 0348 011 555 0334/011 555 0426
<i>Candida (surveillance)</i>	<i>Candida</i> (surveillance)	<i>Candida</i> surveillance	CHARM Mycology Reference Laboratory

Pathogen / Test Species Name (disease/ Syndrome)	Pathogen Species Name (disease/ syndrome, ICD-code)	Tests To Request	Department and Contact telephone no's
			011 555 0325 /011 555 0381/011 555 0323
<i>Chlamydia trachomatis</i>	<i>Chlamydia trachomatis</i>	/ NAAT (PCR, TMA)	STI: 011 555 0461 \ 011 555 0468
<i>Cryptococcus (surveillance)</i>	Cryptococcus (surveillance)	Cryptococcal surveillance	CHARM Mycology Reference Laboratory 011 555 0325/011 555 0381/011 555 0323
Cryptococcus Meningitis	Identification of Cryptococcal antigen (CrAg)	Identification of Cryptococcal antigen (CrAg)	CHARM: Mycology Reference Laboratory 011 386-6430 /011 386-6431/011 555 0323
Diphtheria	<i>Corynebacterium diphtheriae</i>	Identification (phenotypic), and toxin production (Elek and PCR)	CRDM 011 555 0315/011 555 0317
<i>E. coli</i> characterization	Diarrhoeagenic <i>Escherichia coli</i> or suspected STEC	Enteric surveillance and reference function / <i>E. coli</i> characterization	CED Bacteriology 011 386 6235 /011 555 0348 011 555 0334/011 555 0426
<i>Enterococcus faecium</i> and <i>faecalis</i>	<i>Enterococcus faecium</i> and <i>faecalis</i>	AMR detection/confirmation	CHARM AMRL-CC 011 555 0344 / 011 555 0342
Enterovirus	Enterovirus	Enterovirus / PCR / Enterovirus genotyping	CVI Polio 011 386 6438 011 555 0504
Fungal pathogen from clinical sources which cannot be identified by academic/ referral laboratories	Fungal pathogen from clinical sources which cannot be identified by academic/ referral laboratories	Identification of fungal pathogen; Antifungal susceptibility testing	CHARM Mycology Reference Laboratory 011 555 0323 /011 555 0325 011 555 0353
Genital Discharge syndrome	<i>Neisseria gonorrhoeae</i> <i>Chlamydia trachomatis</i> <i>Trichomonas vaginalis</i> and <i>Mycoplasma genitalium</i>	Multiplex PCR for genital discharge pathogens Organism specific PCR available for confirmation of <i>Neisseria gonorrhoeae</i> <i>Chlamydia trachomatis</i> <i>Trichomonas vaginalis</i> and <i>Mycoplasma genitalium</i>	STI 011 555 0461 011 555 0468
Genital Ulcer syndrome	<i>Treponema pallidum</i> , Herpes simplex virus 1 and 2, <i>Haemophilus ducreyi</i> , Lymphogranuloma venereum	Multiplex PCR for genital ulcer pathogens Organism specific PCR available for confirmation	STI 011 555 0461 011 555 0468
Gram-negative organisms (*As per current GERMS-SA case definition)	Gram-negative organisms (*As per current GERMS-SA case definition)	Surveillance	CHARM AMRL-CC 0115550344 0115550342
Group A <i>Streptococcus</i>	Group A <i>Streptococcus</i>	Group A <i>Streptococcus</i> culture and PCR	CRDM 011 555 0315/011 555 0317
<i>Haemophilus ducreyi</i>	<i>Haemophilus ducreyi</i>	PCR	STI: 011 555 0461 \ 011 555 0468
Hepatitis	Hepatitis For research and surveillance only (Anti-HBs the staff immunity screening request form is completed and must accompany the specimen)	Hepatitis B: Architect System HBsAg, Anti-HBs, Anti-HBc, HBeAg, Anti-HBe, Anti-HBc IgM, HBV Viral load, genotyping Hepatitis C:	CVI hepatitis 011 386 6347 011 555 0534

Pathogen / Test Species Name (disease/ Syndrome)	Pathogen Species Name (disease/ syndrome, ICD-code)	Tests To Request	Department and Contact telephone no's
		Architect System Anti-HCV, HCV Viral load,genotyping	
Human Boca virus	Human Boca virus	Human bocaviruses	CRDM Results:011-386 6404 Queries:011 555 0488 011 386 6373 011 386 3690
Human Corona viruses (229E, OC43, NL63, HKU1)	Human Corona viruses (229E, OC43, NL63, HKU1)	Human coronaviruses	CRDM Results:011-386 6404 Queries:011 555 0488 011 386 6373 011 386 3690
Human Metapneumovirus (hMPV)	Human Metapneumovirus (hMPV)	RSVP	CRDM Results:011-386 6404 Queries:011 555 0488 011 386 6373 011 386 3690
Histoplasmosis	Histoplasma galactomannan antigen (HAT)	Histoplasma antigen	CHARM :Mycology Reference Laboratory 011 555 0325 /011 555 0323 011 555 0353
HIV	HIV	HIV-1 Pooled Plasma	HIV Sero-Molecular: 011 386 6435 /011 386 6437 / 011 386 6460
HIV	HIV	CD4	HIV Sero-Molecular: 011 386 6435 /011 386 6437 / 011 386 6460
HIV	HIV	HIV Viral Load	HIV Sero-Molecular: 011 386 6435 /011 386 6437 / 011 386 6460
HIV	HIV	HIV ELISA	HIV Sero-Molecular: 011 386 6435 /011 386 6437 / 011 386 6457
HIV	HIV	HIV-1/2 Antigen-Antibody EIA	HIV Sero-Molecular: 011 386 6435 /011 386 6437 / 011 386 6457
HIV	HIV	HIV-1 or HIV-2 Western Blot	HIV Sero-Molecular: 011 386 6435 /011 386 6437 / 011 386 6457
HIV	HIV	Geenius HIV ½ Confirmatory Test (On Request Only)	HIV Sero-Molecular: 011 386 6435 /011 386 6437 / 011 386 6457
HIV	HIV	Alere Determine Combo Rapid (On Request Only)	HIV Sero-Molecular: 011 386 6435 /011 386 6437 / 011 386 6457
HIV	HIV	HIV Incidence Testing	HIV Sero-Molecular: 011 386 6435 /011 386 6437 / 011 386 6457
HIV-1	HIV-1 (p24)	P24 – Culture Supernatant	Virology Research Laboratory Lab Manager: 011 386 6341
HIV-1	HIV-1	Neutralization assays a) Pseudovirion assay	Virology Research Laboratory Lab Manager: 011 386 6341
HIV-1	HIV-1	Drug resistance testing a) In-House assay	Virology Research Laboratory Lab Manager: 011 386 6341
HIV-1	HIV-1	Gag ELISA	Virology Research Laboratory Lab Manager: 011 386 6341

Pathogen / Test Species Name (disease/ Syndrome)	Pathogen Species Name (disease/ syndrome, ICD-code)	Tests To Request	Department and Contact telephone no's
HIV-1	HIV-1	HIV Virus isolation	Virology Research Laboratory Lab Manager: 011 386 6341
HPV	HPV	PCR / Hybridisation	STI :011 555 0461 \ 011 555 0468
HSV 1 and 2	HSV 1 and 2	PCR; Acyclovir resistance testing by PCR/ sequencing & viral culture; and serum ELISA for HSV2 IgG	STI :011 555 0461 \ 011 555 0468
Influenza A and Influenza B virus.	Influenza A and Influenza B virus.	RSVP	CRDM Results:011-386 6404 Queries:011 555 0488/ 011 386 6373 011 386 3690
<i>Leptospirosis (LEPTO)</i>	<i>Leptospira spp.</i> (Leptospirosis)	Leptospirosis (LEPTO)	Special Bacterial Pathogens Reference Laboratory 011 555 0331/ 011 555 0306
<i>Listeria spp</i>	<i>Listeria spp</i>	Enteric surveillance and reference function / <i>Listeria</i> characterization	CED Bacteriology 011 386 6235 /011 555 0348 011 555 0334 /011 555 0426
<i>Lympho granuloma Venereum (LGV)</i>	<i>Lympho granuloma Venereum (LGV)</i>	PCR	STI :011 555 0461 \ 011 555 0468
MALDI-TOF identification	MALDI-TOF identification	Identification	CHARM AMRL-CC 0115550344/0115550342
<i>Measles^{CVI1} Rash-based Surveillance</i>	Measles Rash-based Surveillance	Measles IgM	CVI measles working group 011 386 6343 011 386 6398 011 555 0534
MERS-Coronavirus (MERS-CoV)	MERS-Coronavirus (MERS-CoV)	Suspected MERS-Coronavirus (MERS-CoV)	Hotline-Dr on call:082 883 9920 CRDM Results:011-386 6404 Queries:011 386 6373 011 386 3690
Monkeypox		Monkeypox	Special Viral Pathogens Laboratory Dr J Weyer: 011 386 6376 / 082 903 9131 NICD Hotline
Multi-drug resistant <i>Enterobacteriaceae</i> and other <i>ESKAPE</i> pathogens for surveillance and detection (*As per current GERMS-SA case definition)	Multi-drug resistant <i>Enterobacteriaceae</i> (*As per current GERMS-SA case definition and case report forms)	Cultures on Dorset slopes viable for up to one month at room temperature for surveillance or reference diagnostic	CHARM AMRL-CC 0115550344 /0115550342
<i>Mycobacterium tuberculosis</i>	<i>Mycobacterium tuberculosis</i>	Microscopy for acid-fast bacilli	Centre for TB 011 885 5317/011 885 5316
<i>Mycobacterium tuberculosis</i>	<i>Mycobacterium tuberculosis</i> or Non-tuberculosis mycobacteria	Culture	Centre for TB 011 8855317/011 885 5316
<i>Mycobacterium tuberculosis</i>	<i>Mycobacterium tuberculosis</i>	Phenotypic drug susceptibility testing (Specify drugs to be tested)	Centre for TB 011 885 5317/011 885 5316
<i>Mycobacterium tuberculosis</i>	<i>Mycobacterium tuberculosis</i> or Non-tuberculosis mycobacteria	Line Probe Assay - First line drug susceptibility - Second line drug susceptibility - NTM speciation	Centre for TB 011 885 5317/011 885 5316

Pathogen / Test Species Name (disease/ Syndrome)	Pathogen Species Name (disease/ syndrome, ICD-code)	Tests To Request	Department and Contact telephone no's
		- MTBcomplex speciation - NTM- drug susceptibility	
<i>Mycobacterium tuberculosis</i>	<i>Mycobacterium tuberculosis</i>	Xpert MTB/RIF Assay	Centre for TB 011 885 5217/011 885 5316
<i>Mycobacterium tuberculosis</i>	<i>Mycobacterium tuberculosis</i> or Non-tuberculosis mycobacteria	-Broth Microdilution or MGIT MIC testing. -Extended Panel for TB MIC testing for NTM. -MIC testing for Bedaquiline and / or Delaminid.	Centre for TB 011 885 5317/011 885 5316
<i>Mycoplasma genitalium</i>	<i>Mycoplasma genitalium</i>	Polymerase chain reaction (PCR)	STI: 011 555 0461 /011 555 0468
<i>Mycoplasma pneumoniae</i> , <i>Chlamydia pneumoniae</i> , <i>Legionella spp.</i> , <i>Bordetella spp.</i>	<i>Mycoplasma pneumoniae</i> , <i>Chlamydia pneumoniae</i> , <i>Legionella spp.</i> , <i>Bordetella spp.</i>	Polymerase chain reaction (PCR) for identification	CRDM 011 555 0315 011 555 0317
<i>Neisseria gonorrhoeae</i>	<i>Neisseria gonorrhoeae</i>	Culture and Antimicrobial susceptibility testing / NAAT (PCR, TMA)/ Typing by PCR / Antimicrobial resistant determinants testing by PCR	STI: 011 555 0461 /011 555 0468
Bacterial meningitis screen	<i>N. meningitidis</i> , <i>H. influenzae</i> and <i>S. pneumoniae</i> , Group B <i>Streptococcus</i> , <i>L. monocytogenes</i> , <i>S. aureus</i> , <i>E.coli</i>	Phenotypic identification and serotyping/grouping MIC Antigen detection PCR for identification and serotyping/ grouping	CRDM 011 555 0315/011 555 0317
Norovirus	Norovirus Gastroenteritis: Diarrhoea, vomiting, fever	Norovirus detection	CED Virology 011 555 0370
Parainfluenza virus types 1, 2 and 3	Parainfluenza virus types 1, 2 and 3	RSVP	CRDM Results;011-386 6404 Queries:011 555 0488 011 386 6373 011 386 3690(HOD)
Parasite - <i>Acanthamoeba spp.</i>	Parasite - <i>Acanthamoeba spp.</i> (<i>Acanthamoeba keratitis</i>)	Acanthamoeba	Parasitology Reference Laboratory 011 555-0304 011 555-0311
Parasite - <i>Babesia species (Babesiosis)</i>	Parasite - <i>Babesia species (Babesiosis)</i>	Babesia	Parasitology Reference Laboratory 011 555-0304 011 555-0311
Parasite - <i>Borrelia duttoni</i> or <i>B. recurrentis (Relapsing fever)</i>	Parasite - <i>Borrelia duttoni</i> or <i>B. recurrentis (Relapsing fever)</i>	Relapsing fever borrelia	Parasitology Reference Laboratory 011 555-0304 011 555-0311
Parasite - <i>Entamoeba histolytica</i>	Parasite - <i>Entamoeba histolytica (Amoebiasis)</i>	Amoebiasis	Parasitology Reference Laboratory 011 555-0304 011 555-0311
Parasite - <i>Echinococcus species</i>	Parasite - <i>Echinococcus species (Echinococcosis, hydatid disease)</i>	Echinococcus	Parasitology Reference Laboratory 011 555-0304 011 555-0311
Parasite - <i>Enterobius vermicularis (pinworm)</i>	Parasite - <i>Enterobius vermicularis (pinworm)</i>	<i>Enterobius vermicularis/ (pinworm)</i>	Parasitology Reference Laboratory 011 555-0304 011 555-0311
Parasite - <i>Human cestodes (tapeworms) including: Taenia saginata, T. solium, H. nana, H. diminuta and D. latum</i>	Parasite - <i>Human cestodes (tapeworms) including: Taenia saginata, T. solium, H. nana, H. diminuta and D. latum</i>	Stool parasites (specify suspected parasite, if applicable)	Parasitology Reference Laboratory 011 555-0304 011 555-0311

Pathogen / Test Species Name (disease/ Syndrome)	Pathogen Species Name (disease/ syndrome, ICD-code)	Tests To Request	Department and Contact telephone no's
<i>Parasite -Human nematodes including: Ascaris lumbricoides (common roundworm), hookworms (Ancylostoma duodenale/ Necator americanus), Trichuris trichiura (whipworm), Strongyloides stercoralis</i>	<i>Parasite -Human nematodes including: Ascaris lumbricoides (common roundworm), hookworms (Ancylostoma duodenale/ Necator americanus), Trichuris trichiura (whipworm), Strongyloides stercoralis</i>	Stool parasites (specify suspected parasite, if applicable)	Parasitology Reference Laboratory 011 555-0304 011 555-0311
<i>Parasite -Human protozoa including: Cryptosporidium species, Cystoisosporal belli, Cyclospora cayetanensis, Giardia lamblia, Entamoeba coli, Blastocystis hominis</i>	<i>Parasite -Human protozoa including: Cryptosporidium species, Cystoisosporal belli, Cyclospora cayetanensis, Giardia lamblia, Entamoeba coli, Blastocystis hominis</i>	Stool parasites (specify suspected parasite, if applicable)	Parasitology Reference Laboratory 011 555-0304 011 555-0311
<i>Parasite -Human trematodes including: Schistosoma mansoni/ haematobium</i>	<i>Parasite -Human trematodes including: Schistosoma mansoni/ haematobium (Bilharzia, schistosomiasis)</i>	Stool parasites (specify suspected parasite, if applicable)	Parasitology Reference Laboratory 011 555-0304 011 555-0311
<i>Parasite -Leishmania species</i>	<i>Parasite -Leishmania species (Leishmaniasis)</i>	Leishmania investigation	Parasitology Reference Laboratory 011 555-0304 011 555-0311
<i>Parasite -Microfilaria species (W. bancrofti, L. loa, M. perstans etc)</i>	<i>Parasite -Microfilaria species (W. bancrofti, L. loa, M. perstans etc)</i>	Microfilaria investigation	Parasitology Reference Laboratory 011 555-0304 011 555-0311
<i>Parasite -Plasmodium falciparum</i>	<i>Parasite -Plasmodium falciparum (Malaria)</i>	Antigen test for <i>Plasmodium falciparum</i>	Parasitology Reference Laboratory 011 555-0304 011 555-0311
<i>Parasite -Plasmodium species</i>	<i>Parasite -Plasmodium species (Malaria)</i>	Malaria investigation	Parasitology Reference Laboratory 011 555-0304 011 555-0311
<i>Parasite -Plasmodium species</i>	<i>Parasite -Plasmodium species (Malaria)</i>	Malaria PCR	Parasitology Reference Laboratory 011 555-0304 011 555-0311
<i>Parasite -Pneumocystis jirovecii</i>	<i>Parasite -Pneumocystis jirovecii (Pneumocystis pneumonia, PCP)</i>	<i>Pneumocystis jirovecii</i> PCR	Parasitology Reference Laboratory 011 555-0304 011 555-0311
<i>Parasite -Schistosoma haematobium</i>	<i>Parasite -Schistosoma haematobium (Schistosomiasis, bilharzia)</i>	<i>Schistosoma haematobium</i>	Parasitology Reference Laboratory 011 555-0304 011 555-0311
<i>Parasite -Strongyloides stercoralis</i>	<i>Parasite -Strongyloides stercoralis</i>	<i>Strongyloides stercoralis</i>	Parasitology Reference Laboratory 011 555-0304 011 555-0311
<i>Parasite -Tapeworm (adult) identification</i>	<i>Parasite -Tapeworm (adult) identification</i>	Tapeworm identification	Parasitology Reference Laboratory 011 555-0304 011 555-0311
<i>Parasite -Trypanosoma species</i>	<i>Parasite -Trypanosoma species (Sleeping sickness or trypanosomiasis)</i>	Trypanosome investigation	Parasitology Reference Laboratory 011 555-0304 011 555-0311
<i>Parasite -Wuchereria bancrofti</i>	<i>Parasite -Wuchereria bancrofti (Lymphatic filariasis or elephantiasis)</i>	<i>W. bancrofti</i> antigen test	Parasitology Reference Laboratory 011 555-0304 011 555-0311
Pertussis	<i>Bordetella pertussis</i>	<i>B. pertussis</i> culture and/or PCR	CRDM 011 555 0315/011 555 0317
Poliovirus	Poliovirus	Real Time RT-PCR and/or Sequencing	Centre for Vaccines & Immunology: Polio Laboratory 011 386 6438 011 555 0504

Pathogen / Test Species Name (disease/ Syndrome)	Pathogen Species Name (disease/ syndrome, ICD-code)	Tests To Request	Department and Contact telephone no's
Poliovirus isolation. AFP surveillance.	Poliovirus (Acute Flaccid Paralysis (AFP) surveillance)	Poliovirus isolation. AFP surveillance.	Poliovirus Isolation 011 555 0504/011 386 6361 011 386 6358
Poliovirus Serology	Poliovirus Serology	Poliovirus Antibody Neutralization	Poliovirus Serology 011 555 0504/011 386 6361 011 386 6358
Rabies	Rabies (Rabies and rabies-related lyssaviruses) ICD-10 code is A82	Rabies	Center for Emerging Zoonotic and Parasitic Diseases, Special Viral Pathogens Laboratory Dr J Weyer:011 386 6376 082 903 9131 NICD Hotline
Rabies immunity (for individuals with pre- exposure rabies vaccination)	Rabies antibodies	Rabies immunity	Center for Emerging Zoonotic and Parasitic Diseases, Special Viral Pathogens Laboratory Dr J Weyer 011 386 6376/ 082 903 9131
Rhinovirus	Rhinovirus	RSVP	CRDM Results;011-386 6404 Queries:011 555 0488 011 386 6373 011 386 3690(HOD)
Rotavirus	Rotavirus Gastroenteritis: Diarrhoea, vomiting, fever	Rotavirus detection /Rotavirus genotyping	CED Virology 011 555 0370
RSV	RSV	RSVP	CRDM 011 555 0488/ 011386 6373
Rubella (done as a reflex test for Measles) <small>CV12, CV13</small>	See measles	See Measles	CVI measles working group 011 386 6398 011 555 0534 011 386 6343
<i>Salmonella</i>	<i>Salmonella</i> Typhi <i>Salmonella</i> Paratyphi A, B and C Nontyphoidal <i>Salmonella</i>	Enteric surveillance and reference function / <i>Salmonella</i> characterization	CED Bacteriology 011 386 6235 /011 555 0348 011 555 0334 /011 555 0426
Sapovirus	Sapovirus Gastroenteritis: Diarrhoea, vomiting, fever	Sapovirus detection	CED Virology 011 555 0370
SARS-CoV-2	COVID-19	COVID-19	Virology Research Laboratory Lab Manager L Nxumalo 011 386 6341
SARS-CoV-2 COVID-19 (Coronavirus disease 2019)	SARS-CoV-2 COVID-19	SARS-CoV-2	CRDM 011-386 6373 011 555 0315/17

Pathogen / Test Species Name (disease/ Syndrome)	Pathogen Species Name (disease/ syndrome, ICD-code)	Tests To Request	Department and Contact telephone no's
Severe fever with thrombocytopenia syndrome (SFTS)	SFTS	SFTS PCR	Special Viral Pathogens Laboratory Dr J Weyer 011 386 6376 082 903 9131 /Contact NICD Hotline
<i>Shigella spp</i>	<i>Shigella spp</i>	Enteric surveillance and reference function / <i>Shigella</i> characterization	011 386 6235 /011 555 0348 011 555 0334 /011 555 0426
<i>Staphylococcus aureus</i> , <i>Enterococcus faecium</i> , <i>Enterococcus faecalis</i> and other Gram-positive organisms surveillance and detection	<i>Staphylococcus aureus</i> <i>Enterococcus faecium</i> , <i>Enterococcus faecalis</i> and other Gram-positive organisms	AMR detection/confirmation	AMRL-CC 0115550344 /0115550342
<i>Treponema pallidum</i>	<i>Treponema pallidum</i>	RPR/TPPA/ PCR / Typing/ Antimicrobial resistant determinants testing by PCR / Macrolide resistance testing by PCR/ Sequencing	STI: 011 555 0461\ 011 555 0468
<i>Trichomonas vaginalis</i>	<i>Trichomonas vaginalis</i>	PCR\ Culture	STI: 011 555 0461\ 011 555 0468
Urinary antigen test for identification	<i>S. pneumoniae</i> , <i>Legionella pneumophila serogroup 1</i>	Urinary antigen test for identification	CRDM 011 555 0315/011 555 0317
Viral haemorrhagic fever(VHF)	Viral haemorrhagic fevers (Crimean-Congo haemorrhagic fever, Ebola, Marburg, Lassa, Hanta and Yellow fever) ICD-10 code for unspecified viral haemorrhagic fever, A99	VHF	Center for Emerging Zoonotic and Parasitic Diseases, Special Viral Pathogens Laboratory Dr J Weyer 011 386 6376 082 903 9131 /Contact NICD Hotline
Virus screening for infections of unknown aetiology	Virus identification to Family level	TEM negative staining for viral diagnostics	Centre for Emerging Zoonotic and Parasitic Diseases, Electron Microscopy laboratory Dr M Birkhead 011 386 6318
<i>Yersinia pestis</i> (Plague)	<i>Yersinia pestis</i> (Plague)	Plague (PLAGE) Plague serology (PLAGE) / Plague surveillance (RATS)	Special Bacterial Pathogens Reference Laboratory 011 555 0331/ 011 555 0306

CVI1: Please note that Measles IgG should not be requested by users of the service as it is only done for NICD staff immunity screening

CVI2: Please note that the Rubella IgM test should not be requested by users of the service as it is not offered as a single test, the Rubella IgM is done as part of the testing for national measles surveillance for the Expanded Programme on Immunization (EPI), every sample that is sent for a suspected measles case will be tested for both measles and rubella IgM

CVI3: Please note that Rubella IgG should not be requested by users of the service as it is only done for NICD staff immunity screening.

Notifiable Medical Conditions (NMC) Disease List

CATEGORY 1 NOTIFIABLE MEDICAL CONDITIONS –

Category 1 notifiable medical conditions that require immediate reporting by the most rapid means available upon diagnosis followed by a written or electronic notification to the Department of Health within 24 hours of diagnosis by health care providers, private health laboratories or public health laboratories

* Food -borne disease outbreak is the occurrence of two or more cases of a similar foodborne disease resulting from the ingestion of a common food.

* *Examples of novel respiratory pathogens include novel influenza A virus and MERS

coronavirus.

* ** Viral haemorrhagic fever diseases include Ebola or Marburg viruses, Lassa virus, Lujo virus, new world arena viruses, Crimean -Congo haemorrhagic fever or other newly identified viruses causing haemorrhagic fever.

Acute flaccid paralysis

Acute rheumatic fever

Anthrax

Botulism

Cholera

Diphtheria

Enteric fever (typhoid or paratyphoid fever)

Food borne disease outbreak*

Haemolytic uraemic syndrome (HUS)

Listeriosis

Malaria

Measles

Meningococcal disease

Pertussis

Plague

Poliomyelitis

Rabies (human)

Respiratory disease caused by a novel respiratory pathogen**

Rift Valley fever (human)

SARS-CoV-2 (COVID-19)

Smallpox

Viral haemorrhagic fever diseases***

Yellow fever

CATEGORY 2 NOTIFIABLE MEDICAL CONDITIONS –

Category 2 notifiable medical conditions to be notified through a written or electronic notification to the Department of Health within seven (7) days of clinical or laboratory diagnosis by health care providers, private health laboratories or public health laboratories

Agricultural or stock remedy poisoning

Bilharzia (schistosomiasis)

Brucellosis

Congenital rubella syndrome

Congenital syphilis

Haemophilus influenzae type B

Hepatitis A

Hepatitis B

Hepatitis C

Hepatitis E

Lead poisoning

Legionellosis

Leprosy

Maternal death (pregnancy, childbirth and puerperium)

Mercury poisoning

Soil transmitted helminths (*Ascaris Lumbricoides*, *Trichuris trichiuria*, *Ancylostoma*)

Tetanus

Tuberculosis: pulmonary

Tuberculosis: extra-pulmonary

Tuberculosis: multidrug-resistant (MDR-TB)

Tuberculosis: extensively drug-resistant (XDR-TB)

CATEGORY 3 NOTIFIABLE MEDICAL CONDITIONS –

Category 3 notifiable medical conditions to be notified through a written or electronic notification to the Department of Health within 7 days of diagnosis by private and public health laboratories

Ceftriaxone-resistant *Neisseria gonorrhoea*

West Nile virus, Sindbis virus, Chikungunya virus (i.e. endemic arboviruses)

Dengue fever virus, other imported arboviruses of medical importance (i.e. non-endemic)

Salmonella spp. other than *S. Typhi* and *S. Paratyphi*

Rubella virus

Shiga toxin-producing Escherichia coli
--

Shigella spp.

CATEGORY 4 NOTIFIABLE MEDICAL CONDITIONS –

Category 4 notifiable medical conditions to be notified through a written or electronic notification to the Department of Health within 1 month of diagnosis by private and public health laboratories
--

Healthcare-associated infections or multidrug-resistant organisms of public health importance	· Carbapenemase-producing Enterobacteriaceae
	· Vancomycin-resistant enterococci
	· Staphylococcus aureus: hGISA and GISA
	· Colistin-resistant Pseudomonas aeruginosa
	· Colistin-resistant Acinetobacter baumannii
	· Clostridium difficile

Tests offered at the NICD per Centre

Centre for Vaccines and Immunology (CVI)

Pathogen Species Name (disease/syndrome)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
<p>Poliovirus</p> <p>(Acute Flaccid Paralysis (AFP) surveillance)</p> <p>Note: For purposes of AFP surveillance, the Poliovirus Isolation Laboratory at NICD is the only laboratory in South Africa accredited by World Health Organization to perform this testing.</p>	<p>Two stool samples (adult thumb size) collected in universal sample containers, 24-48 hours apart and within 14 days of onset of paralysis. In the event a stool sample cannot be obtained, please forward a rectal swab as an alternative.</p>	<p>Maintenance of cold chain (2-8°C) from collection to receipt at NICD. Samples to reach laboratory within 3 days of collection.</p> <p>Note: These are guidelines met to comply with WHO requirements; however samples will be subjected to processing and testing even if these criteria are not met with relevant details captured accordingly.</p>	<p>Poliovirus isolation. AFP surveillance.</p>	<p>Virus Isolation</p>	<p>EPI Notification 14 days (80% of all results to be reported within 14 days of receipt in lab). Issue of LIMS report: 21 days of receipt in lab.</p>	<p>The following information is required for capture on the AFP database and should be completed on specific AFP case investigation forms:</p> <ul style="list-style-type: none"> • Epid number – unique number for epidemiology purposes • Name • Date of Birth • Date of onset of paralysis • Date of stool collection • Immunization history • Province • District • Case or contact • Specimen number • Date sent from field to National Level – where applicable • Date received at National Level – where applicable • Date sent to NICD • Testing requested and clinical diagnosis to be clearly stated to prevent incorrect / unnecessary tests being performed 	<p>Poliovirus Laboratory</p> <p>011 555 0504</p> <p>011 386 6361</p> <p>011 386 6358</p> <p>011 386 6438</p>

Pathogen Species Name (disease/ syndrome)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
Poliovirus	Virus isolate from stool or environmental sample, minimum 1ml. FTA Card inoculated with virus isolate	Transport frozen or at 4°C. Specimen must reach the laboratory within 3 days of dispatch. Transport at ambient temperature. Specimen must reach the laboratory within 3 days of dispatch.	Real Time RT-PCR Sequencing	Real Time ITD RT-PCR Real Time VDPV Screening RT-PCR Sequencing VP1 Region	7 days 7 days	AFP Investigation Form with at least the following information: <ul style="list-style-type: none"> • Epid Number • Patient Name • Date of onset of paralysis • District • Case or contact • Specimen Number • Referring Laboratory specimen number • Date sent to NICD • Cell Line (L-Arm or R-Arm) Real Time PCR Results (if requesting Sequencing)	Centre for Vaccines & Immunology: Polio Laboratory 011 386 6438 011 555 0504
Poliovirus Serology	Clotted Blood - 5ml, Sera - 0.5ml	Maintenance of cold chain (2-8°C)	Poliovirus Antibody Neutralization	Neutralization Test	21 days NB: Samples are batched for testing. This may result in TATs being exceeded if not stated as urgent.	Testing requested and clinical diagnosis to be clearly stated on request forms to prevent incorrect / unnecessary tests being performed	Poliovirus Serology 011 555 0504 011 386 6361 011 386 6358
Enterovirus	Hand foot and mouth disease: Dextran swab of vesicle/rash fluid with a stool sample Meningitis/encephalitis: CSF with a stool sample Respiratory disease: throat swab/lavage with a stool sample Gastroenteritis/diarrhoea: Stool sample/rectal swab	Transport frozen or at 4°C. Specimen must reach the laboratory within 3 days of dispatch.	Enterovirus PCR Enterovirus genotyping	Real-time PCR Sequencing of VP1 region	7 Days 14 days	Cold chain essential for the storage and transport of samples	Centre for Vaccines & Immunology: Polio Laboratory 011 386 6438 011 555 0504

Pathogen Species Name (disease/ syndrome)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
Measles (Rash-based Surveillance)	5 ml Clotted Blood or EDTA Blood (red top or yellow SST tube or EDTA/heparin Blood or Serum	The blood/serum samples can be transported at room temperature to NICD, Samples should reach the lab within three days of sample collection (If transportation is delayed the samples should be refrigerated	Measles IgM	Anti-Measles IgM antibody (EIA)	7 working days	A Measles Case Investigation Form (CIF) must be completed in full and must accompany specimens. Please ensure that the following information is captured on the CIF: <ul style="list-style-type: none"> • Name, age, Date of Birth , address of patient • Name and contact number of clinician • Symptoms marked in tick boxes • Complications marked in tick boxes • Date of onset of rash • Date seen at health facility • Date specimen collected • Medical history including measles vaccination history, number of doses and date of last measles vaccination Date of notification and response to case	CVI measles working group 011 386 6398 011 555 0534 011 386 6343
Rubella (Rash based surveillance)	5 ml Clotted Blood or EDTA Blood (red top or yellow SST tube or EDTA/heparin Blood or Serum	Same as for measles	See note on page 16 under Measles & Rubella	Anti-Rubella IgM antibody (EIA)	7 working days	Same as for Measles	CVI measles working group 011 386 6398 011 555 0534

Pathogen Species Name (disease/ syndrome)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
Measles	Throat swab Please note Once a measles outbreak is identified (a cluster of IgM-positive cases) please collect 5-10 throat swab specimens from other suspected cases in the area and place in VTM tube (Viral Transport Medium) The instruction to collect throat swabs will be issued by the CDCs. Throat swabs should not be collected from sporadic suspected cases.	Transport at 2°C-8°C Sample should reach laboratory within 3 days	For surveillance only	RT-PCR Sequencing	14 days 14 days	Surveillance only. Case Investigation Forms (CIF) must be completed in full and must accompany specimens.	CVI measles working group 011 386 6343 011 386 6398 011 555 0534
Hepatitis	Serology: 5 ml Clotted Blood or EDTA Blood (red top or yellow SST tube or EDTA Blood or Serum) Molecular: 5ml EDTA blood	Transport at 2°C-8°C Sample should reach laboratory within 3 days	For research and surveillance only (Anti-HBs for staff immunity screening)	Hepatitis B: Architect System HBsAg, Anti-HBs, Anti-HBc, HBeAg, Anti-HBe, Anti-HBc IgM, HBV Viral load, genotyping Hepatitis C: Architect System Anti-HCV, HCV Viral load, genotyping	Serology: 14 days Molecular: 21 days	For research and surveillance only (Anti-HBs the staff immunity screening request form is completed and must accompany the specimen)	CVI hepatitis working group 011 386 6347 011 555 0534

Centre for Tuberculosis

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
<i>Mycobacterium tuberculosis</i>	Sputum or other human sample	Cool ambient temperature Specimens should be collected in clean leak proof containers free from paraffin and other waxes or oils. Specimens should be kept cool during transportation but should not be frozen. If there is a delay in transport, the specimen should be refrigerated (2-8°C) An adequate sample should be about 5-10ml sputum.	Microscopy for acid-fast bacilli	Microscopy for acid-fast bacilli	90% within 48-72hours	Test limited for surveillance and projects	Centre for TB 011 885 5323 011 885 5316
<i>Mycobacterium tuberculosis</i> or Non-tuberculous mycobacteria	Sputum or other human samples type	Specimens should be collected in clean leak proof containers free of oils. Study specific containers may be used. Sputum should be from deep cough avoiding saliva and secretions from nasopharynx. Specimens should be kept cool during transportation but should not be frozen. If there is a delay in transport, the specimen should be refrigerated (2-8°C) An adequate sample should be about 5-10ml sputum.	Culture	Culture	Project dependant		Centre for TB 011 8855323 011 885 5316

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
<i>Mycobacterium tuberculosis</i> Non-tuberculous mycobacteria	Isolate in MGIT tubes (or other liquid media) and on solid media	Isolates should be packaged as a biohazard category A (triple packaging)	Drug susceptibility testing (Specify drugs to be tested)	Drug susceptibility testing (MGIT 960)	3-8 weeks from pure positive sub-culture or Project dependant	For Bedaquiline Surveillance a special request form should be completed Contact laboratory manager for form. Extended TB DST should be requested via email to the pathologist.	Centre for TB 011 885 5323 011 885 5316

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
<p><i>Mycobacterium Tuberculosis</i> Non-tuberculous mycobacteria</p>	<p>Sputum (or other samples type) and/or culture (liquid/solid)</p>	<p>Cool ambient temperature Specimens should be collected in clean leak proof containers free from paraffin and other waxes or oils. Specimens should be kept cool during transportation but should not be frozen. If there is a delay in transport, the specimen should be refrigerated (2-8°C) An adequate sample should be about 5-10ml sputum. For isolates these should be packaged as biohazard category A (triple packaging)</p>	<p>Line Probe Assay First line drug susceptibility Second line drug susceptibility</p> <ul style="list-style-type: none"> - NTM speciation - MTB complex speciation - NTM- drug susceptibility 	<p>Molecular testing methods GenoType MTBDRplus, GenoType MTBDRsl, , GenoType CM GenoType MTBC GenoType NTM DR</p>	<p>2-7 days specimen or AFB positive culture or Project dependant</p>	<p>Culture isolates these should be packaged as a biohazard category A (triple packaging)</p>	<p>Centre for TB 011 885 5323 011 885 5316</p>

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
<i>Mycobacterium tuberculosis</i>	Sputum or other specimen or decontaminated sediment	Cool ambient temperature Specimens should be collected in clean leak proof containers free from paraffin and other waxes or oils. Specimens should be kept cool during transportation but should not be frozen. If there is a delay in transport, the specimen should be refrigerated (2-8°C) An adequate sample should be about 5-10ml sputum.	Xpert MTB/RIF Assay	Molecular testing methods	48-72 hours or Project dependant		Centre for TB 011 885 5217 011 885 5316
<i>Mycobacterium tuberculosis</i> or Non-tuberculosis mycobacteria	Isolate in MGIT tubes (or other liquid media) and on solid media	Isolates should be packaged as a biohazard category A (triple packaging)	-Broth Microdilution or agar proportion or MGIT MIC testing. -Extended Panel for MTB -MIC testing for Bedaquiline and / or Delaminid. MIC testing for NTM.	Advanced DST Methods Bedaquiline MIC testing Delamamid MIC testing Extended TB DST Sensitre: <ul style="list-style-type: none"> • MYCOTB (MTB) • Rapid grower NTM • Slow grower NTM 	3-8 weeks from pure positive sub-culture or Project dependant	For Delaminid and Bedaquiline Surveillance, a special request form should be completed. Contact the Lab Manager for the forms.	Centre for TB 011 885 5323 011 885 5316

Centre for HIV and STI

Pathogen Species Name (disease/syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
Bacterial vaginosis	Vaginal smear	Transport at ambient temperature		Gram stain		Only available for surveillance and research purposes	STI 011 555 0461 011 555 0468
<i>Candida</i> species	Vaginal smear	Transport at ambient temperature		Gram stain		Only available for surveillance and research purposes	STI 011 555 0461 011 555 0468
Herpes Simplex Type 2	Clotted Blood EDTA Blood Serum Plasma	Transport on ice		HSV2 IgG EIA	Project Dependant	Only available for Surveillance purposes and research projects	STI 011 555 0461 011 555 0468
Herpes Simplex Virus Type 1 and 2	Ulcer swab (Dacron)	Transport on ice or frozen	HSV PCR	PCR	10 days	Only available for Surveillance purposes and research project Special cases: persistent/recurrent genital ulcer disease (require prior consultation with STI Section staff & completion of specific request form)	STI 011 555 0461 011 555 0468
Herpes Simplex Virus Type 1 and 2 acyclovir resistance testing	Ulcer swab (Dacron): 1 swab in viral transport medium 1 swab in dry sterile container	Transport on ice	HSV PCR and sequencing for thymidine kinase +/- DNA polymerase mutations HSV viral culture and acyclovir plaque reduction assay	PCR + sequencing and viral culture	Project Dependant	Only available for Surveillance purposes and research project Special cases: persistent/genital ulcer disease non-responsive to acyclovir therapy (require prior consultation with STI Section staff & completion of specific request form)	STI 011 555 0461 011 555 0468

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
Genital Ulcer Disease	Ulcer swab (Dacron)	Transport on ice or frozen	Multiplex PCR for genital ulcer pathogens Organism-specific PCR available for confirmation of HSV and LGV	PCR	10 days	Available for Surveillance purposes and research projects Special cases: persistent/ recurrent genital ulcer disease (require prior consultation with STI Section staff & completion of specific request form)	STI 011 555 0461 011 555 0468
Genital Discharge	Urethral swab Vaginal swab Other samples as indicated by clinical presentation Endocervical swab Rectal swab Pharyngeal swab (Dacron swabs)	Transport on ice or frozen	Multiplex PCR for genital discharge pathogens Organism specific PCR available for confirmation of <i>Neisseria gonorrhoeae</i> <i>Chlamydia trachomatis</i> <i>Trichomonas vaginalis</i> and <i>Mycoplasma genitalium</i>	PCR	10 days	Available for Surveillance purposes and research projects Special cases: persistent/ recurrent genital discharge or suspected treatment failure for gonorrhoea: *sexual abuse/ assault (require prior consultation with STI Section staff & completion of specific request form)	STI 011 555 0461 011 555 0468

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
<p><i>Neisseria gonorrhoeae</i> culture isolate</p>	<p>Viable culture isolate on sealed agar plate</p> <p>For AST and confirmation of ceftriaxone-resistance, additionally inoculate:</p> <ol style="list-style-type: none"> 1) chocolate agar slope (overnight incubation in CO₂) <p>AND Trypticase soy broth with 10% glycerol (frozen at -20°C or -70°C) and transported on ice</p>		<p>Identification</p> <p>Antimicrobial Susceptibility Testing (AST)</p> <p>Confirmation of extended-spectrum cephalosporin resistance</p>	<p>Culture, phenotypic ID, PCR</p> <p>Culture, ID, AST (E-test and agar dilution)</p> <p>Culture, ID, AST (E-test and agar dilution) + antimicrobial resistance determinants testing by PCR</p>	<p>7 days</p>	<p>Sexual abuse/ assault case</p> <ul style="list-style-type: none"> • Two different phenotypic confirmatory methods should be performed on culture isolate at the local NHLS laboratory • *Refer swab specimen for multiplex PCR for genital discharge pathogens, if required (see above) <p>Suspected treatment failure for gonorrhoea or extended-spectrum cephalosporin resistance (require prior consultation with STI Section staff & completion of specific request form)</p> <p>Ceftriaxone-resistant gonorrhoea is a notifiable medical condition (Category 3). All clinical isolates identified as resistant to ceftriaxone (+/- cefixime) on phenotypic AST should be referred to STI Reference laboratory, NICD, for confirmation of resistance (requires completion of specific request form)</p>	<p>STI 011 555 0461 011 555 0468</p>

Pathogen Species Name (disease/syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
HIV	Plasma in EDTA or ACD tube	Frozen 1ml Plasma (-20°C) or Whole blood to be separated within 24 hours of collection	HIV-1 Pooled Plasma	Roche HIV-1 Viral Load	4 weeks		HIV Sero-Molecular Laboratory 011 386 6330 011 386 6439
HIV	Plasma in EDTA or ACD tube	Frozen 1ml Plasma (-20°C) or Whole blood to be separated within 24 hours of collection	HIV-1 RNA PCR	Roche HIV-1 Viral load	5 days		HIV Sero-Molecular Laboratory 011 386 6330 011 386 6439
HIV	Whole blood (EDTA or ACD Tube)	4°C to room temperature Minimum 2ml whole blood DO NOT FREEZE	HIV DNA PCR	HIV-1 DNA PCR (Whole blood)	5 days		HIV Sero-Molecular Laboratory 011 386 6330 011 386 6439
HIV	Dried Blood Spots collected on 903 Specimen Whatman collection paper	Sealed in plastic bags with desiccant and indicator to prevent moisture Minimum of three (3) full spots per card must be sent	HIV DNA PCR	HIV-1 DNA PCR (DBS)	5 days		HIV Sero-Molecular Laboratory 011 386 6330 011 386 6439
HIV	Frozen EDTA whole blood	Minimum of 2 X 100ul of frozen whole blood	HIV TNA PCR	HIV TNA PCR	5 days		HIV Sero-Molecular Laboratory 011 386 6330 011 386 6439
HIV	Whole blood (EDTA or ACD Tube)	Whole blood at room temperature	CD4	CD4 (TCell lymphocyte subset)	24 – 48 hours	No cold chain transport, transport at ambient temperature	HIV Sero-Molecular Laboratory 011 386 6330 011 386 6439
HIV	Dried Blood Spots collected on 903 Specimen Whatman collection paper	Sealed in plastic bags with desiccant and indicator to prevent moisture Minimum of three (3) full spots per card must be sent	HIV DBS viral load	Roche HIV-1 quantitative test on DBS FVE protocol	5 days or Project dependent		HIV Sero-Molecular Laboratory 011 386 6330 011 386 6439

Pathogen Species Name (disease/syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
HIV	Dried Blood Spots collected on 903 Specimen Whatman collection paper	Sealed in plastic bags with desiccant and indicator to prevent moisture Minimum of three (3) full spots per card must be sent	HIV DBS viral load	ABBOTT RealTime HIV-1 quantitative PCR on DBS	5 days or Project dependent		HIV Sero-Molecular Laboratory 011 386 6330 011 386 6439
HIV	5 ml/10ml Clotted Blood or EDTA Blood and Dried Blood Samples (DBS) collected on Whatman 903 Cards - minimum of 3 full circles of whole blood, preferably fill all 5 circles	Transport blood samples at 2°C-8°C Maintenance of cold chain is required from collection to receipt at NICD. DBS samples can be transported at RT°C (stable at RT°C for 14 days)	HIV ELISA	HIV-1/2 Antibody EIA	5 -7 working days or project dependent	Project/Survey requirements must be provided in writing by Requestor and discussed with NICD prior to testing.	HIV Sero-Molecular 011 386 6435 011 386 6437 011 386 6457
HIV	5 ml/10ml Clotted Blood, EDTA Blood or serum or plasma	Transport blood samples at 2°C-8°C Maintenance of cold chain is required from collection to receipt at NICD	Alere HIV Combo Rapid (On Request Only)	HIV Rapid	5 -7 working days or project dependent	Alere HIV Combo Rapid (On Request Only)	HIV Sero-Molecular: 011 386 6435 /011 386 6437 / 011 386 6457
HIV	5 ml/10ml Clotted Blood – serum or EDTA Blood and Dried Blood Samples (DBS) collected on Whatman 903 Cards - minimum of 3 full circles of whole blood, preferably fill all 5 circles	Transport blood samples at 2°C-8°C Maintenance of cold chain is required from collection to receipt at NICD. DBS samples can be transported at RT°C (stable at RT°C for 14 days)	HIV ELISA	HIV – 1/2 Antigen-Antibody EIA	5 -7 working days or project dependent	Project/Survey requirements must be provided in writing by Requestor and discussed with NICD prior to testing.	HIV Sero-Molecular 011 386 6435 011 386 6437 011 386 6457

Pathogen Species Name (disease/syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
HIV	5 ml/10ml Clotted Blood – serum or EDTA Blood and Dried Blood Samples (DBS) collected on Whatman 903 Cards – minimum of 3 full circles of whole blood, preferably fill all 5 circles	Transport blood samples at 2°C-8°C Maintenance of cold chain is required from collection to receipt at NICD. DBS samples can be transported at RT°C (stable at RT°C for 14 days)	HIV-1 or HIV-2 Western Blot	HIV-1 Western Blot HIV-2 Western Blot Geenius HIV ½ Confirmatory Rapid Assay (Serum, Plasma or Whole blood only)	5 -7 working days or project dependent	Project/Survey requirements must be provided in writing by Requestor and discussed with NICD prior to testing.	HIV Sero-Molecular 011 386 6435 011 386 6437 011 386 6457
HIV	5 ml/10ml Clotted Blood or EDTA Blood and Dried Blood Samples (DBS) collected on Whatman 903 Cards – minimum of 3 full circles of whole blood, preferably fill all 5 circles	Transport blood samples at 2-8°C Maintenance of cold chain is required from collection to receipt at NICD DBS samples can be transported at Room Temperature for (stable for 14 days at RT°C after collection)	HIV Incidence Testing	Sedia HIV-1 BED Incidence EIA (Serum/Plasma only) Sedia HIV-1 LAg Avidity Incidence EIA (Serum/Plasma) Maximum HIV-1 LAg Avidity Incidence EIA (Serum/Plasma/DBS) (Research Purposes Only)	14 - 21 working days for up to 1000 samples or project dependent	Project/Survey requirements must be provided in writing by Requestor and discussed with NICD prior to testing.	HIV Serol-Molecular 011 3866475 011 386 6437
HIV-1	p24	Store at -70°C		Culture supernatants	2 months	All these tests are done for Research/Project purposes only.	Virology Research Laboratory Lab Manager L Nxumalo 011 386 6341
HIV-1	Neutralization assays a) Pseudovirion assay	Transported on dry ice		Serum /plasma, monoclonal antibodies	2 months or project dependent	All tests are done for Research / Project purposes	Virology Research Laboratory Lab Manager L Nxumalo 011 386 6341
HIV-1	Drug resistance testing a) In-House assay	Store at -70°C		Plasma from EDTA blood	2 months	Can be used for diagnostic purposes in special circumstances.	Virology Research Laboratory Lab Manager L Nxumalo 011 386 6341

In the event of a dispute concerning this document, the electronic version stored on Q-Pulse will be deemed to be the correct version

National Health Laboratory Services- All rights reserved

Pathogen Species Name (disease/syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
HIV-1	Gag ELISA	Store at -20°C		Serum	2 months	All these tests are done for Research/Project purposes only.	Virology Research Laboratory Lab Manager L Nxumalo 011 386 6341
HIV-1	HIV Virus isolation	LN storage		PBMC's	2 months	Whole blood received for testing to be stored at ambient temperature only	Virology Research Laboratory Lab Manager L Nxumalo 011 386 6341
HIV-1	5ml/10ml Clotted blood or EDTA blood and Dried blood samples (DBS) collected on Whatman 903 Cards – minimum of 3 full circles of whole blood, preferably fill all 5 circles or frozen plasma	Transport blood samples at 2-8°C DBS samples can be transported at room temperature (stable for 14 days at RT°C after collection) Transport can be on Dry Ice and stored at -70°C	Drug resistance testing In-house assay	Plasma from EDTA blood DBS	2 Months or project dependant	Project/Survey requirements must be provided in writing by Requestor and discussed with NICD prior to testing	Virology Research Laboratory Lab Manager L Nxumalo 011 386 6341
HIV-1	Mycoalert Mycoplasma	Store at -20°C		Cell supernatants	1 month		Virology Research Laboratory Lab Manager L Nxumalo 011 386 6341
<i>Treponema pallidum</i>	Clotted blood or EDTA Serum Plasma	Transport on ice		RPR	Project dependant	Only available for Surveillance purposes and research projects	STI 011 555 0461 011 555 0468
<i>Treponema pallidum</i>	Clotted blood or EDTA Serum Plasma	Transport on ice		TPPA	Project dependant	Only available for Surveillance purposes and research projects	STI 011 555 0461 011 555 0468
<i>Treponema pallidum</i>	Clotted blood or EDTA Serum Plasma	Transport on ice		TPAb	Project dependant	Only available for Surveillance purposes and research projects	STI 011 555 0461 011 555 0468

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
<i>Treponema pallidum</i>	Ulcer swab (Dacron)	Transport on ice or frozen	Syphilis/ <i>Treponema pallidum</i> PCR	PCR	10 days	Only available for Surveillance purposes and research projects and special cases (see above)	STI 011 555 0461 011 555 0468
<i>Treponema pallidum</i>	Ulcer swab (Dacron)	Transport on ice or frozen		Molecular Typing	Project dependant	Only available for Surveillance purposes and research projects	STI 011 555 0461 011 555 0468
<i>Treponema pallidum</i>	Ulcer swab (Dacron)	Transport on ice or frozen		Macrolide resistance testing by PCR/ Sequencing	Project dependant	Only available for Surveillance purposes and research projects	STI 011 555 0461 011 555 0468
<i>Lympho granuloma Venereum (LGV)</i>	Ulcer swab Genital swab (Dacron)	Transport on ice or frozen Transport on ice or frozen		PCR	10 days	Only available for Surveillance purposes and research projects And special cases (see above)	STI 011 555 0461 011 555 0468
HPV	Genital, rectal and oropharyngeal swabs	Transport on ice or frozen		PCR/ Hybridisation	Project dependent	Only available for Surveillance purposes and research project	STI 011 555 0461 011 555 0468
<i>Chlamydia trachomatis</i>	Urine Urethral Swab Vaginal swab Endocervical swab Oropharyngeal swab Rectal swab (Dacron swabs)	Transport on ice or frozen		NAAT (PCR, TMA)	10 days	Only available for Surveillance purposes and research project and special cases (see above)	STI 011 555 0461 011 555 0468
<i>Haemophilus. ducreyi</i>	Ulcer swab (Dacron swabs)	Transport on ice or frozen		PCR	10 days	Only available for Surveillance purposes and research project And special cases (see above)	STI 011 555 0461 011 555 0468
<i>Mycoplasma genitalium</i>	Urine Urethral swab Vaginal swab Endocervical swab (Dacron swabs)	Transport on ice or frozen		PCR	10 days	Only available for Surveillance purposes and research project And special cases (see above)	STI 011 555 0461 011 555 0468

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
<i>Neisseria gonorrhoeae</i>	Urethral swab Vaginal swab Other samples as indicated by clinical presentation Endocervical swab Rectal swab Pharyngeal swab Cultured strains (Dacron or nylon flocced swabs)	Use special transport media (Amies Stuarts). Transport at 4°C to reach lab within 48 hours.		Culture and Antimicrobial susceptibility testing	7 days	Only available for Surveillance purposes and research project And special cases (see above)	STI 011 555 0461 011 555 0468
<i>Neisseria gonorrhoeae</i>	Urine Urethral swab Vaginal swab Endocervical swab Penile swab Rectal swab Pharyngeal swab (Dacron or nylon flocced swabs)	Transport on ice or frozen		NAAT (PCR, TMA)	10 days	Only available for Surveillance purposes and research project And special cases (see above)	STI 011 555 0461 011 555 0468
<i>Neisseria gonorrhoeae</i>	Urethral swab Vaginal swab Other samples as indicated by clinical presentation Endocervical swab Rectal swab Pharyngeal swab (Dacron or nylon flocced swabs) Cultured strains	Transport on ice or frozen Except for cultured strains at ambient temperature		Typing by PCR / Sequencing	Project dependent	Only available for Surveillance purposes and research projects And special cases e.g. extensively-drug resistant ceftriaxone-resistant strains (see above)	STI 011 555 0461 011 555 0468
<i>Neisseria gonorrhoeae</i>	Urethral swab Vaginal swab Other samples as indicated by clinical presentation Endocervical swab Rectal swab Pharyngeal swab (Dacron or nylon flocced swabs) Cultured strains	Transport on ice or frozen Except for cultured strains at ambient temperature		Antimicrobial resistance testing by PCR	Project dependent	Only available for Surveillance purposes and research project. And special cases e.g. extensively-drug resistant strains (see above)	STI 011 555 0461 011 555 0468

Pathogen Species Name (disease/syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
SARS-CoV-2	(i) Pseudovirion assay (ii) Spike and RBD ELISA	Transport on dry ice Store at -70°C or cooler. Transport on dry ice or at 2-8°C. Depends on the cohort. Plasma/serum received for testing to be stored at 2-8°C if for immediate testing or -70°C or cooler for long term storage		Serum Serum/Plasma	2 months 2 months	All these tests are done for Research/Project purposes only.	Virology Research Laboratory Lab Manager L Nxumalo 011 386 6341
<i>Trichomonas vaginalis</i>	Urine Urethral swab Vaginal swab Endocervical swab	Transport on ice or frozen		NAAT PCR,TMA)PCR	10 days	Only available for Surveillance purposes and research project And special cases	STI 011 555 0461 011 555 0468

Centre for Emerging Zoonotic and Parasitic Diseases

Pathogen Species Name (disease/syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
<p>Arbovirus</p> <p>(Chikungunya, Dengue, West Nile, Rift Valley fever, Sindbis, Zika)</p> <p>ICD-10 code for unspecified arthropod-borne viral fever, A94</p>	<p>Clotted blood or serum, minimum of 1.0 ml</p> <p>Highly recommended to submit repeat specimens, preferably acute and convalescent specimens</p>	<p>Tubes in sealed in leak-proof containers.</p> <p>Label clearly as biohazardous. Adhere to national and international regulations for transportation of hazardous biological goods as required.</p> <p>Address to Centre for Emerging Zoonotic and Parasitic Diseases, Arbovirus Reference Laboratory, cold transport on ice packs Contact Laboratory or NICD Hotline for consult on urgent cases</p>	<p>Arbovirus</p> <p>(Routine arbo screen, include serology for Chikungunya, West Nile, Sindbis, Rift Valley fever)</p> <p>Dengue and Zika for patient with travel history.</p> <p>Zika virus neutralization test</p> <p>PCR/Isolation for Acute cases only</p> <p>Disease specific IgM or IgG ELISA</p>	<p><u>Serology:</u> (HAI is performed as initial screen; any positive results are followed up with IgM ELISA to confirm recent infection.</p> <p>Testing of acute and convalescent specimens to indicate seroconversion by four fold rise in IgG titre is highly recommended or IgM seroconversion) <i>Zika virus neutralization test is performed for confirmation of ELISA positive results.</i></p>	<p>HAI: 5 days</p> <p>HAI + IgM ELISA: 7 days</p> <p>IgM ELISA: 2-4 days</p> <p>Zika virus neutralization test: 21 days</p>	<p>Rash or arthralgia syndromes with fever which may be linked to travel history, insect bites. Some cases develop encephalitis.</p> <p>Refer to Arbovirus Diagnostic Guide or Outbreak specific guidelines available from www.nicd.ac.za</p> <p>Highly recommended to submit repeat / paired specimens, preferably acute and convalescent specimens.</p> <p>Submit Case investigation form (available from NICD website)</p>	<p>Arbovirus Reference Laboratory Dr O Hellferscee 011 386 6424/6353 082 908 8045</p> <p>NICD Hotline</p>

Pathogen Species Name (disease/syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
	<p>Alternative/additional specimens for PCR</p> <p>CSF (for encephalitic cases only, requires at least 500 µl)</p> <p>Liver biopsies (for post mortem confirmation only, contact Laboratory)</p>	See above	<p>Arbovirus PCR <i>(specify which arbovirus, for example "Dengue PCR")</i></p> <p>Arbovirus isolation</p>	<p>RT-PCR</p> <p>Virus isolation</p>	<p>24-48 hours</p> <p>18 days</p>	<p>See above.</p> <p>Should be requested in addition to the Arbovirus routine serology screen for acute cases only.</p> <p>Only useful for specimens collected within 7 days of onset of illness or acutely ill patients.</p> <p>Submit Case investigation form (available from NICD website)</p> <p><i>Note: Arbovirus isolation is no longer offered, until further notice.</i></p>	See above
<i>Bacillus anthracis</i> (anthrax)	<p><u>Cutaneous</u> Vesicular stage: Soak 2x sterile dry swabs in vesicular fluid from a previously unopened vesicle.</p> <p>Eschar stage: Rotate 2x sterile dry swabs for 2-3s beneath the edge of eschar without removing it.</p> <p>Biopsy of lesion: Fresh Tissue in PBS/saline <i>Please note:</i> Do not send preserved tissue</p> <p><u>Inhalation</u> Pleural fluid or sputum (>1 ml) collected in sterile container</p> <p>Blood culture <u>plus</u> 1 tube clotted blood and 1 tube whole blood (EDTA tube)</p>	<p>Transport at room temperature</p> <p>Transport at 2-8°C</p> <p>Transport blood culture at room temperature and blood tubes at 2-8°C</p>	<p>Anthrax (ANTHR)</p>	<p>Microscopy, culture and special identification</p>	<p>5 days</p>	<p>Safety precautions should be taken when handling/collecting samples</p> <p>Samples should be taken prior to antibiotic treatment and should reach laboratory as quickly as possible</p> <p><u>Important</u> Please notify laboratory prior to sending specimens</p>	<p>Special Bacterial Pathogens Reference Laboratory 011 555 0331 011 555 0306</p>

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
	<p><u>Gastrointestinal</u> Stool or rectal swab</p> <p>Blood culture <u>plus</u> 1 tube clotted blood and 1 tube whole blood (EDTA tube)</p>	<p>Transport at 2-8°C</p> <p>Transport blood culture at room temperature and blood tubes at 2-8°C</p>					
	<p><u>Meningitis</u> CSF in sterile container</p> <p>Blood culture <u>plus</u> 1 tube clotted blood and 1 tube whole blood (EDTA tube)</p>	<p>Transport at 2-8°C</p> <p>Transport blood culture at room temperature and blood tubes at 2-8°C</p>					
<i>Clostridium botulinum</i> (botulism)	20 to 30 ml clotted blood or 10 to 15 ml serum 25 to 50 g of stool Gastric washing Vomit Suspected food	Transport at 2-8°C	Botulinum	Mouse neutralization assay, Anaerobic culture and confirmation	4 weeks	<u>Important</u> Please notify laboratory prior to sending specimens. Samples should be taken as soon as possible after onset of illness.	Special Bacterial Pathogens Reference Laboratory 011 555 0331 011 555 0306
Monkeypox	<p><u>Prodromal phase:</u> Tonsillar tissue swab, nasopharyngeal swab, clotted blood, serum or whole blood. Minimum of 0.5ml serum</p> <p><u>Rash/lesion phase:</u> Biopsy, lesion fluid, scab, clotted blood, serum or whole blood. Minimum of 0.5ml serum</p>	<p>Specimen containers must be sealed Leak-proof containers</p> <p>Adhere to national and international transportation of biological goods. Address to Centre for Emerging Zoonotic and Parasitic Diseases, Special Viral Pathogens, cold transport on ice packs</p>	Monkeypox	Monkeypox PCR	24-48 hrs	Fever, headache, muscle aches, backache, chills, exhaustion, lymphadenopathy, lesions. Travel history to central and western Africa or other exposure events (i.e. contact with exotic pets)	Special Viral Pathogens Laboratory Dr J Weyer 011 386 6376 082 903 9131 NICD Hotline

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
Rabies	Clotted blood or serum, minimum of 0.5 ml	Specimen containers must be sealed in leak-proof containers. Adhere to national and international transportation of biological goods. Address to Centre for Emerging Zoonotic and Parasitic Diseases, Special Viral Pathogens, cold transport on ice packs.	Rabies immunity	Serology: Rabies IgG	24-48 hours	Only for post vaccinal immunity screening. Usually for high risk occupational groups such as veterinarians or animal handlers or laboratory workers in laboratories that handle rabies virus	Special Viral Pathogens Laboratory Dr J Weyer 011 386 6376 082 903 9131
Rabies (Rabies and rabies-related lyssaviruses) ICD-10 code is A82.9	Antemortem: Saliva, minimum of 0.5 ml CSF, minimum of 0.5 ml Nuchal biopsies, single biopsy collected with dermatological punch from nape of neck <i>Post mortem:</i> Brain. Brainstem is most sensitive specimen. Whole, half or cubes of representative regions of brain submitted in 50 % glycerol saline in plastic screw top containers. Brain biopsies collected via the supraorbital fissure (contact laboratory for instructions). Do not submit in formalin. If no saline available freeze fresh.	Specimen containers must be sealed in leak-proof containers. Adhere to national and international regulations for transportation of hazardous biological goods as required. Address to Centre for Emerging Zoonotic and Parasitic Diseases, Arbovirus, cold transport on ice packs Contact Laboratory or NICD Hotline for consult on urgent cases	Rabies	<i>Antemortem:</i> RT-PCR (performed on saliva, CSF and biopsy) Serology (Rabies IgG/IgM) (performed on CSF and blood/serum) <i>Post mortem:</i> Rabies fluorescent antibody test (DFA or FAT)	24-48 hrs 24-48 hours 24-48 hours	For antemortem investigation of suspected rabies cases. Encephalitis with rapid progression, hydrophobia, dementia, hallucinations, lucid periods usually present. Suggested that all specimens should be co-submitted to rule out rabies. It is highly recommended to submit at least 3 consecutively collected saliva samples (collected at least 3-6 hours apart). Serology is of limited value to investigate clinical cases, but may be performed as part of battery of laboratory investigations. Post- mortem investigation is most conclusive. Contact Laboratory/NICD Hotline for advice	Special Viral Pathogens Laboratory Dr J Weyer 011 386 6376 082 903 9131 NICD Hotline

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
Severe fever with thrombocytopenia syndrome (SFTS)	Clotted blood or serum, minimum of 0.5 ml	Specimen containers must be sealed leak-proof containers. Adhere to national and international transportation of biological goods. Address to Centre for Emerging Zoonotic and Parasitic Diseases, Special Viral Pathogens, cold transport on ice packs Contact Laboratory or NICD Hotline	SFTS	SFTS PCR	24-48 hrs	High fever, thrombocytopenia, leukocytopenia and gastrointestinal disorders. Travel history to endemic areas or other exposure events	Special Viral Pathogens Laboratory Dr J Weyer 011 386 6376 082 903 9131 NICD Hotline
<i>Leptospira</i> spp. (Leptospirosis)	1 Tube of clotted blood or serum <i>Please note:</i> Tests not done on urine, plasma, haemolysed, icteric or lipemic blood.	Transport at 2-8°C Sample should reach laboratory within 3 days.	Leptospirosis (LEPTO)	Serology: IgM ELISA	4 days		Special Bacterial Pathogens Reference Laboratory 011 555 0331 011 555 0306

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
<p>Viral haemorrhagic fevers</p> <p>(Crimean-Congo haemorrhagic fever, Ebola, Marburg, Lassa, Hanta and Yellow fever)</p> <p>ICD-10 code for unspecified viral haemorrhagic fever, A99</p>	Clotted blood or serum, minimum of 1.0 ml	<p>Tubes in sealed in leak-proof containers. Label clearly as biohazardous and suspected VHF. Adhere to national and international regulations for transportation of hazardous biological goods (Category A) as required.</p> <p>Address to Center for Emerging Zoonotic and Parasitic Diseases, Special pathogens, cold transport on ice packs</p> <p>Contact Laboratory and/or NICD Hotline</p>	VHF	<p>Serology (Fluorescent antibody test: IgG and IgM ELISA: IgG and IgM)</p> <p>RT-PCR</p> <p>Virus isolation</p>	<p>24-48 hrs</p> <p>24-48 hrs</p> <p>21 days</p>	<p>Fever, Rash Headache, Arthralgia, Myalgia, Haemorrhagic manifestations, Gastrointestinal symptoms, Pathology indicators, Travel history or other exposure events</p> <p>Contact Laboratory and/or NICD Hotline</p> <p>Submission of repeat specimen critical to confirm or exclude VHF</p> <p>Full battery of tests (serology, PCR and isolation) recommended for most conclusive testing)</p>	<p>Special Viral Pathogens Laboratory Dr J Weyer 011 386 6376 082 903 9131</p> <p>Contact NICD Hotline</p>
Virus screening (infections of unknown aetiology)	Cultured isolates, any fluid specimen (minimum 0.1 ml) in sterile container or dried between 2 microscope slides and sealed (lesion fluid, aspirate, serum, CSF, tears), scabs, crusts, biopsy/autopsy tissues (in saline and not larger than 10 ³ mm), clarified stool samples	<p>Specimen containers must be sealed in plastic bags.</p> <p>Adhere to national and international transportation of biological goods. Address to Centre for Emerging Zoonotic and Parasitic Diseases, Electron Microscopy Laboratory</p> <p>Transport at 2-8°C or room temperature</p>	<u>Viral screening</u>	<u>Transmission electron microscopy and negative staining</u>	<u>3 days</u>	<u>Please contact laboratory for specimen-specific requirements. Notify the laboratory when specimens are dispatched</u>	<p>Electron Microscope Laboratory</p> <p>011 386 6318</p>

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's			
<i>Yersinia pestis</i> (Plague)	<u>Bubonic</u> Bubo aspirate in PBS in sterile container plus 2x swabs (absorb a few drops of sample on a sterile swab and place into Cary Blair transport medium)	Transport at 2-8°C or room temperature.	Plague (PLAGE)	Microscopy (Gram, Wayson & DFA), culture and confirmation	5 days	Safety precautions should be taken when handling/collecting samples Samples should be taken prior to antibiotic treatment and should reach laboratory as quickly as possible <u>Important</u> Please notify laboratory prior to sending specimens	Special Bacterial Pathogens Reference Laboratory 011 555 0331 011 555 0306			
	<u>Pneumonic</u> Sputum in sterile container plus 2x swabs (rotate swab in sputum sample and place into Cary Blair transport medium)	Transport at 2-8°C or room temperature.								
	<u>Septicemic</u> Blood culture bottle	Transport at room temperature								
	1 tube clotted blood or serum <i>Please note:</i> Paired serum taken 2-3 weeks apart is required for confirmation	Transport at 2-8°C						Plague serology (PLAGE)	Serology: ELISA	2 days
	<u>Surveillance</u> Clotted animal blood e.g. rodent, dog	Transport at 2-8°C						Plague surveillance (RATS)	Serology	4 weeks
Parasitology										
<u>Parasite -Borrelia duttoni</u> or <i>B. recurrentis</i> (Relapsing fever)	Unclotted, EDTA blood or blood films. <i>Clotted blood is unsuitable.</i>	Transport at ambient temperature or on ice.	Relapsing fever borrelia	Staining and microscopy	1 day		Parasitology Reference Laboratory 011 555-0304 011 555-0311			
<u>Parasite -Babesia</u> species (Babesiosis)	Unclotted, EDTA blood or blood films. <i>Clotted blood is unsuitable.</i>	Transport at ambient temperature or on ice.	Babesia	Staining and microscopy	1 day		Parasitology Reference Laboratory 011 555-0304 011 555-0311			

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
Parasite - <i>Plasmodium</i> species (Malaria)	Unclotted, EDTA blood or blood films. <i>Clotted blood is unsuitable.</i>	Transport at ambient temperature or on ice.	Malaria investigation	Staining and microscopy	1 day		Parasitology Reference Laboratory 011 555-0304 011 555-0311
Parasite - <i>Plasmodium</i> species (Malaria)	Unclotted, EDTA blood. Blood spots or blood films may be used in exceptional circumstances. <i>Clotted blood is unsuitable.</i>	Transport at ambient temperature or on ice. Blood should be transported to the laboratory as quickly as possible.	Malaria PCR	Malaria PCR	2 days	Only request PCR test when: <ul style="list-style-type: none"> ■ the malaria microscopy and RDT (rapid diagnostic test) results do not correlate ■ the malaria microscopy and/or RDT results are negative and malaria is still suspected ■ malaria species confirmation is needed ■ the patient has already been treated for malaria but routine tests are either not done or negative 	Parasitology Reference Laboratory 011 555-0304 011 555-0311
Parasite - <i>Plasmodium falciparum</i> (Malaria)	Unclotted, EDTA blood. <i>Clotted blood is unsuitable.</i>	Transport at ambient temperature or on ice.	Antigen test for <i>Plasmodium falciparum</i>	Antigen test for <i>Plasmodium falciparum</i>	1 day		Parasitology Reference Laboratory 011 555-0304 011 555-0311
Parasite -Microfilaria species (<i>W. bancrofti</i> , <i>L. loa</i> , <i>M. perstans</i> etc)	Unclotted, EDTA blood or blood films. Blood sampling should be performed at the correct times depending on the suspected filarial species*	Transport at ambient temperature or on ice.	Microfilaria investigation	Staining and microscopy	1 day	*For <i>W. bancrofti</i> sample after 8pm, for <i>L. loa</i> at ~12pm and others at any time.	Parasitology Reference Laboratory 011 555-0304 011 555-0311

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
Parasite - <i>Trypanosoma</i> species (Sleeping sickness or trypanosomiasis)	Unclotted, EDTA blood, blood films or fresh CSF (1ml). <i>Clotted blood is unsuitable.</i>	Transport at ambient temperature or on ice. Ideally CSF specimens must reach laboratory within 30 minutes of sampling.	Trypanosome investigation	Staining and microscopy	1 day		Parasitology Reference Laboratory 011 555-0304 011 555-0311
Parasite -Human protozoa including: <i>Cryptosporidium</i> species, <i>Cystoisospora belli</i> , <i>Cyclospora cayetanensis</i> , <i>Giardia lamblia</i> , <i>Entamoeba coli</i> , <i>Blastocystis hominis</i>	Stool or duodenal aspirate/string test. Specimens, that may take longer than a day to reach the laboratory, should ideally be preserved in an equal quantity of 10% formalin.	Transport at ambient temperature or on ice.	Stool parasites (specify suspected parasite, if applicable)	Microscopic identification of all human protozoa	1 day		Parasitology Reference Laboratory 011 555-0304 011 555-0311
Parasite - <i>Entamoeba histolytica</i> (Amoebiasis)	Stool, liver abscess fluid or cyst fluid. Minimum volume is 1ml.	Transport at ambient temperature or on ice. Don't add preservative. Ideally specimen must reach laboratory within 30 minutes of sampling.	Amoebiasis	Microscopy	1 day		Parasitology Reference Laboratory 011 555-0304 011 555-0311
Parasite -Human nematodes including: <i>Ascaris lumbricoides</i> (common roundworm), hookworms (<i>Ancylostoma duodenale/ Necator americanus</i>), <i>Trichuris trichiura</i> (whipworm), <i>Strongyloides stercoralis</i>	Stool or worm/s. Specimens that may take longer than a day to reach the laboratory should ideally be preserved in an equal quantity of 10% formalin.	Transport at ambient temperature or on ice.	Stool parasites (specify suspected parasite, if applicable)	Macroscopic and/or microscopic identification.	1 day		Parasitology Reference Laboratory 011 555-0304 011 555-0311
Parasite - <i>Enterobius vermicularis</i> (pinworm)	Early morning sticky tape swab is the optimal specimen.	Transport at ambient temperature or on ice.	<i>Enterobius vermicularis/</i> (pinworm)	Microscopy	1 day	Eggs may be infective – handle with care.	Parasitology Reference Laboratory 011 555-0304 011 555-0311

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
<u>Parasite -Strongyloides stercoralis</u>	Sputum, urine or CSF. Minimum volume is 1ml. Larvae may be infective – handle with care.	Transport at ambient temperature or on ice.	<i>Strongyloides stercoralis</i>	Microscopy and/or culture for larvae of <i>Strongyloides stercoralis</i> .	1 day	For disseminated infections.	Parasitology Reference Laboratory 011 555-0304 011 555-0311
<u>Parasite -Human trematodes including: Schistosoma mansoni/ haematobium (Bilharzia, schistosomiasis)</u>	Stool. Specimens, that may take longer than a day to reach the laboratory, should ideally be preserved in an equal quantity of 10% formalin.	Transport at ambient temperature or on ice.	Stool parasites (specify suspected parasite, if applicable)	Microscopic identification of all human trematodes.	1 day		Parasitology Reference Laboratory 011 555-0304 011 555-0311
<u>Parasite -Schistosoma haematobium (Schistosomiasis, bilharzia)</u>	Urine, minimum volume is 5ml.	Transport at ambient temperature or on ice.	<i>Schistosoma haematobium</i>	Microscopy	1 day		Parasitology Reference Laboratory 011 555-0304 011 555-0311
<u>Parasite -Human cestodes (tapeworms) including: Taenia saginata, T. solium, H. nana, H. diminuta and D. latum</u>	Stool. Specimens, that may take longer than a day to reach the laboratory, should ideally be preserved in an equal quantity of 10% formalin.	Transport at ambient temperature or on ice.	Stool parasites (specify suspected parasite, if applicable)	Microscopic identification of all human cestodes/ tapeworms.	1 day		Parasitology Reference Laboratory 011 555-0304 011 555-0311
<u>Parasite -Tapeworm (adult) identification</u>	Worm or proglottid (tapeworm segment). Submit proglottids in saline.	Transport at ambient temperature or on ice.	Tapeworm identification	Macroscopic and/or microscopic identification.	1-2 days	Proglottids may be infective – handle with care.	Parasitology Reference Laboratory 011 555-0304 011 555-0311
<u>Parasite -Echinococcus species (Echinococcosis, hydatid disease)</u>	Cyst tissue, cyst fluid or sputum, minimum volume is 1ml.	Transport at ambient temperature or on ice, if transport is expected to take longer than 2 days it is best to transport at 4°C.	Echinococcus	Microscopic examination for hydatid hooklets and scolices of <i>Echinococcus</i> species.	1 day	May be infective – handle with care.	Parasitology Reference Laboratory 011 555-0304 011 555-0311

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
<u>Parasite -</u> <i>Acanthamoeba</i> spp. (<i>Acanthamoeba</i> keratitis)	Corneal scrapings or biopsy, or contact lenses, cases and solutions. Send tissue in saline.	Transport at ambient temperature or on ice.	Acanthamoeba	Culture, Staining	2 weeks (Prov. result sent day 2)		Parasitology Reference Laboratory 011 555-0304 011 555-0311
<u>Parasite -</u> <i>Leishmania</i> species (Leishmaniasis)	Bone marrow, liver biopsy, skin biopsy/impression smears. Submit skin biopsy in saline.	Transport at ambient temperature or on ice.	Leishmania investigation	Staining and microscopy	1 day		Parasitology Reference Laboratory 011 555-0304 011 555-0311
<u>Parasite -</u> <i>Pneumocystis jirovecii</i> (<i>Pneumocystis</i> pneumonia, PCP)	Respiratory specimens including: induced sputum, tracheal aspirates, bronchial washings, bronchoalveolar lavage (BAL) or gastric wash. Random sputum is not an optimal specimen. Minimum volume is 1ml.	Transport at ambient temperature or on ice.	<i>Pneumocystis jirovecii</i> PCR	PCR	2 days	Please send a respiratory specimen (induced or expectorated sputum, BAL etc.) for this test when PCP is suspected clinically/ radiologically, but the routine lab test (IFA) is negative.	Parasitology Reference Laboratory 011 555-0304 011 555-0311
Vector Control Reference Laboratory							

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
Arthropods/ Insects	<p>Adult mosquitos must be undamaged and preserved on silica. A piece of paper must separate the specimen from the silica crystals. There must be only 1 mosquito per silica tube.</p> <p>Mosquito larvae should preferentially be reared to adults and preserved on silica tubes as described above.</p> <p>Identifying larvae by PCR is not recommended; However, if rearing of the larvae is not possible, larvae must be preserved in 70% ethanol with 1 larva per tube.</p>	Transport at ambient temperature. Estimated time of arrival: 1 week	Morphological identification, PCR identification and/or ELISA where indicated	Morphological identification; PCR identification: <i>An. gambiae</i> and <i>An. funestus</i> identification, MS PCR and <i>kdr</i> PCR and/or ELISA	<p>1 Month for I and PCR identification from the time the samples were received at VCRL.</p> <p>6 weeks for morphologica I identification as well as PCR identification</p> <p>Please note: TAT does not apply for project specimens</p> <p>Samples for ELISA testing are done in a batch, unless ELISA testing of samples is urgent.</p>	<p>Please send undamaged samples and all information relating to each sample must be in document form.</p> <p>Samples must be clearly labelled on the individual tube caps. The labels must match the information sheet.</p> <p>Sample numbers should be listed in consecutive order on the information sheet. Two samples must not have the same number.</p> <p>GPS coordinates for all locations must be provided.</p>	<p>Vector Control Reference Laboratory</p> <p>011 386 6480</p>

Centre for Enteric Diseases

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
<i>Campylobacter spp</i>	Laboratory-confirmed isolate from any body site	Inoculated Transwab Dorset slopes incubated at 37°C overnight at source laboratory, prior to submission to CED-Bacteriology. Transwabs /Dorset slope to be kept at room temperature upon receipt at NICD.	Enteric surveillance and reference function	Phenotypic: Identification Genotypic: Real time PCR for detection of <i>Campylobacter jejuni</i> and <i>Campylobacter coli</i>	Project dependent and GERMS-SA Quarterly stats timelines	Attach stool isolate form obtainable from CED-Bacteriology or copy of laboratory report Surveillance isolates are tested in batches	CED Bacteriology 011 386 6235 011 555 0348 011 555 0334 011 555 0426

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
<p>Diarrhoeagenic <i>Escherichia coli</i></p> <p>Watery or bloody diarrhoea, abdominal cramps, with or without fever</p>	<p>Laboratory- confirmed isolate from stool or rectal swab only.</p> <p>Laboratory confirmed isolate from all body sites for suspect EHEC/<i>E. coli</i> O157.</p> <p>Environmental isolates in water or food-borne disease outbreaks only.</p>	<p>Inoculated Dorset slopes incubated at 37°C overnight at source laboratory, prior to submission to CED- Bacteriology. Dorset slope to be kept at room temperature upon receipt at NICD.</p>	<p>Diarrhoeagenic <i>Escherichia coli</i> characterization</p> <p>Enteric surveillance and reference function</p>	<p>Phenotypic: Identification</p> <p>Serotyping O-antigen</p> <p>Genotypic: Virulence gene detection by multiplex-PCR.</p> <p>DNA fingerprinting of strains by PFGE analysis</p> <p>Phenotypic and genotypic characterization of Diarrhoeagenic <i>Escherichia coli</i></p>	<p>3 Working days</p> <p>7 days</p> <p>3 days</p> <p>4 days</p> <p>Project dependent and GERMS-SA Quarterly stats timelines</p>	<p>Attach stool isolate form obtainable from CED-Bacteriology or copy of patient's laboratory report</p> <p>Provide contact Person's email address</p> <p>Attach stool isolate form obtainable from CED-Bacteriology or copy of laboratory report</p> <p>Surveillance isolates are tested in batches</p>	<p>CED Bacteriology 011 386 6235 011 555 0348 011 555 0334 011 555 0360/0426</p>

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
<p><i>Vibrio cholerae</i> O1 and non-O1 (non-cholera <i>Vibrio</i> species)</p> <p>Acute watery diarrhoea</p>	<p>Laboratory-confirmed cultured isolate from all body sites.</p> <p>Environmental isolates in water or food-borne disease outbreaks only.</p>	<p>Bacterial isolates sub-cultured onto Dorset transport medium.</p> <p>Inoculated dorsets incubated at 37°C overnight at source laboratory, prior to submission to CED Bacteriology.</p> <p>Dorset slope to be kept at room temperature upon receipt at NICD.</p>	<p>Enteric surveillance and reference function / <i>Vibrio</i> species characterization</p>	<p>Phenotypic:</p> <p>Identification</p> <p>Serotyping</p> <p>Antimicrobial susceptibility testing. Screening of ESBL</p> <p>Genotypic:</p> <p>Cholera enterotoxin detection by real-time PCR.</p> <p>DNA fingerprinting of strains by PFGE analysis</p>	<p>3 days</p> <p>3 days</p> <p>4 days</p> <p>1 day</p> <p>4 days</p>	<p>Attach stool isolate form obtainable from CED-Bacteriology or copy of patient's laboratory report</p> <p>Provide contact Person's email address</p>	<p>CED Bacteriology</p> <p>011 386 6235</p> <p>011 555 0348</p> <p>011 555 0334</p> <p>011 555 0360/0426</p>

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
<i>Salmonella</i> Typhi and non-typhoidal <i>Salmonella</i> species	Laboratory- confirmed cultured isolates from all body sites. Environmental isolates in water or food-borne disease outbreaks only.	Bacterial isolates sub- cultured onto Dorset transport medium. Inoculated dorsets incubated at 37°C overnight at source laboratory, prior to submission to CED Bacteriology.	Enteric surveillance and reference function / <i>Salmonella</i> species characterization	Phenotypic and genotypic characterization of <i>Salmonella</i> and <i>Shigella</i> spp	Project dependent GERMS-SA Quarterly stats timelines	Attach stool isolate form obtainable from CED-Bacteriology or copy of patient's laboratory report. Surveillance isolates are tested in batches	CED Bacteriology 011 386 6235 011 555 0348 011 555 0334 011 555 0426
<i>Listeria</i> <i>monocytogenes</i>	Laboratory- confirmed isolate from all body site	Bacterial isolates sub- cultured onto Dorset transport medium. Inoculated dorsets incubated at 37°C overnight at source laboratory, prior to submission to CED Bacteriology.	Characterization of <i>Listeria spp</i> / Enteric surveillance and reference function	Phenotypic: <i>Salmonella</i> serotyping <i>Shigella</i> serotyping Antimicrobial susceptibility testing. Screening of ESBL Genotypic: PFGE analysis: MLVA analysis:	12 Working days 5 days 4 days 4 days 3 days 7 days 2 weeks	Provide contact Person's email address Attach stool isolate form obtainable from CED-Bacteriology or copy of patient's laboratory report. Provide contact Person's email address	CED Bacteriology 011 386 6235 011 555 0348 011 555 0334 011 555 0426
Adenovirus Gastroenteritis: Diarrhoea, vomiting, fever	Specimen: Stool Unsuitable Specimen: Rectal swab in VTM	Maintenance of cold chain from collection to receipt at NICD. Store and transport at 4°C.	Enteric Adenovirus detection	Real-time PCR (Enteric adenovirus)	5 days	Specimen containers closed tightly and sealed in plastic bag accompanied with date of specimen collection and date of birth	CED Virology 011 555 0370

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
Astrovirus Gastroenteritis: Diarrhoea, vomiting, fever	Specimen: Stool Unsuitable Specimen: Rectal swab in VTM	Maintenance of cold chain from collection to receipt at NICD. Store and transport at 4°C.	Astrovirus detection	Real-time PCR	5 days	Specimen containers closed tightly and sealed in plastic bag accompanied with date of specimen collection and date of birth	CED Virology 011 555 0370
Sapovirus Gastroenteritis: Diarrhoea, vomiting, fever	Specimen: Stool Unsuitable Specimen: Rectal swab in VTM	Maintenance of cold chain from collection to receipt at NICD. Store and transport at 4°C.	Sapovirus detection	Real-time PCR	5 days	Specimen containers closed tightly and sealed in plastic bag accompanied with date of specimen collection and date of birth	CED Virology 011 555 0370
Norovirus Gastroenteritis: Diarrhoea, vomiting, fever	Specimen: Stool Unsuitable Specimen: Rectal swab in VTM	Maintenance of cold chain from collection to receipt at NICD. Store and transport at 4°C.	Norovirus detection	Real-time PCR	5 days	Specimen containers closed tightly and sealed in plastic bag accompanied with date of specimen collection and date of birth	CED Virology 011 555 0370
Rotavirus Gastroenteritis: Diarrhoea, vomiting, fever	Specimen: Stool Unsuitable Specimen: Rectal swab in VTM	Maintenance of cold chain from collection to receipt at NICD. Store and transport at 4°C.	Rotavirus detection	ELISA	5 days	Specimen containers closed tightly and sealed in plastic bag accompanied with date of specimen collection and date of birth as well as vaccination history (where possible)	CED Virology 011 555 0370
			Rotavirus genotyping	RT-PCR genotyping			

Centre for Healthcare-Associated Infections, Antimicrobial Resistance and Mycoses

The Centre for Healthcare-Associated Infections, Antimicrobial Resistance and Mycoses (CHARM) aims to prevent and control opportunistic, healthcare-associated infections, to combat AMR (including antifungal and antibacterial resistance) and to provide expertise in mycology in South Africa

Pathogen Species Name (disease/syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
CHARM - Mycology							
Fungal pathogen from clinical sources which cannot be identified by academic/ referral laboratories	Unidentified fungal pathogen from clinical source should be inoculated into an agar slope (screwtop) bottle such as Dorset transport medium or potato dextrose medium	Room temperature; inform Mycology Reference Laboratory if suspected dimorphic fungus which needs to be processed in a BSL-3 facility	Identification of fungal pathogen; Antifungal susceptibility testing	Phenotypic and genotypic identification (where required) of unidentified fungal pathogens from clinical sources; Antifungal susceptibility testing	10 days (unless very slow-growing organism)	Complete case report form (can be obtained from the Mycology Reference Laboratory)	CHARM Mycology Reference Laboratory 011 555 0323 011 555 0325 011 555 0353
Moulds and dimorphic fungi from normally-sterile sites (surveillance)	Inoculated into an agar slope (screwtop) bottle such as Dorset transport medium or potato dextrose medium	Room temperature; inform Mycology Reference Laboratory if suspected dimorphic fungus which needs to be processed in a BSL-3 facility	Identification of fungal pathogen; antifungal susceptibility testing; genotyping	Phenotypic and genotypic identification (where required) of unidentified fungal pathogens from clinical sources; Antifungal susceptibility testing	10 days (unless very slow-growing organism)	Laboratory report	CHARM Mycology Reference Laboratory 011 555 0323 011 555 0325 011 555 0353
Cryptococcus (surveillance)	Cryptococcal isolate should be inoculated into an agar slope (screwtop) bottle such as Dorset transport medium	Room temperature	Cryptococcal surveillance	Identification by phenotypic and genotypic methods; antifungal susceptibility testing	N/A	Complete GERMS case report or attach final lab report; Submission requested from GERMS-SA enhanced surveillance sites; private labs and NHLS labs in KZN	CHARM Mycology Reference Laboratory 011 555 0325 011 555 0381 011 555 0323

Pathogen Species Name (disease/syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
Cryptococcus Meningitis	Whole blood ,Plasma, serum and CSF	Transport at ambient temperature or on ice	Identification of Cryptococcal antigen (CrAg)	CrAg Lateral Flow Assay (LFA) CrAg Enzyme Immunoassay (EIA)	72hrs- 1 week	Laboratory Report and worksheet	CHARM Mycology Reference Laboratory 011 555 0381 011 555 0323 011 555 0325
Candida (surveillance)	Candida isolate should be inoculated into an agar slope (screwtop) bottle such as Dorset transport medium	Room temperature	Candida surveillance	Identification by phenotypic and genotypic methods; antifungal susceptibility testing	N/A	Complete GERMS case report or attach final lab report; Submission requested from participating GERMS-SA labs	CHARM Mycology Reference Laboratory 011 555 0325 011 555 0381 011 555 0323
Histoplasmosis	Urine	Transport in cooler box with ice packs	Identification of Histoplasma galactomannan antigen	Histoplasma Enzyme Immunoassay (EIA)	1 week	Clinical history and case investigation form	CHARM Mycology Reference Laboratory 011 555 0325 011 555 0381 011 555 0323
CHARM Antimicrobial Resistance Laboratory and Culture Collection(AMRL-CC)							

Pathogen Species Name (disease/syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
<i>Staphylococcus aureus</i>	Blood culture isolates, pus aspirate, sputum and other significant isolates sub-cultured on Dorset slopes	Transport at ambient temperature	AMR and virulence factor detection/confirmation	ID, MIC and molecular characterization of resistance mechanisms <i>mecA</i> and <i>mecC</i> ; cfr-linezolid resistant gene, Virulence factor – <i>pvl</i> ; vancomycin AST	14 days from receipt of culture	Referral isolate case Report forms	CHARM AMRL-CC 011 555 0344 011 555 0342
Gram-negative organisms ESKAPE organisms (Carbapenem resistant enterobacteriaceae and <i>Acinetobacter baumannii</i>)	Blood culture isolates, sub-cultured on Dorset slopes	Transport at ambient temperature Cultures on Dorset slopes viable for up to one month at room temperature	Surveillance	ID, MIC and molecular characterization of resistance mechanisms	30-60 days from receipt of culture	Complete GERMS case report or attach final lab report; Submission requested from participating GERMS-SA labs *	CHARM AMRL-CC 011555 0344 011555 0342
Multi-drug resistant (MDROs) Enterobacteriaceae and other organisms (*As per current GERMS-SA case definition)	Cultures on Dorset slopes or other suitable media to sustain viability during transport	Transport at ambient temperature	AMR –CPE; colistin and others detection/	ID, MIC and molecular characterization of resistance mechanisms	2 weeks from receipt of culture	*Case Report forms Highly resistant nosocomial infections	CHARM AMRL-CC 011555 0344 011555 0342
<i>Enterococcus faecium</i> and <i>faecalis</i>	Cultures on Dorset slopes or other suitable media to sustain viability during transport	Transport at ambient temperature Cultures on Dorset slopes viable for up to one month at room temperature	AMR detection/confirmation	<i>Van A, B and C</i> genes	14 days from receipt of culture	Referral isolate case Report forms	CHARM AMRL-CC 011 555 0344 011 555 0342
MALDI-TOF identification	Cultures on Dorset slopes or other suitable media to sustain viability during transport	Transport at ambient temperature	Identification	Identification on MALDI-TOF	1 week (organism dependent)	Submit isolates with final lab report and/or lab working card	CHARM AMRL-CC 011555 0344 011555 0342

Centre for Respiratory Diseases and Meningitis

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
Severe Respiratory Illness: <ul style="list-style-type: none"> • Adenovirus • Influenza A virus {H1N1pdm09 and H3N2} • Influenza B virus • Parainfluenza virus types 1,2,3,4 [PIV] • Respiratory Syncytial virus [RSV] • Rhinovirus • Human Metapneumovirus [hMPV] • Human Boca virus • Human Corona viruses (229E, OC43, NL63, HKU1) 	Any respiratory sample (e.g. Nasopharyngeal or tracheal aspirates, nasal, nasopharyngeal and/or oropharyngeal swabs, sputum, biopsies, autopsies etc.) Swabs should be placed in universal / viral transport medium (UTM/VTM). Nasopharyngeal- and oropharyngeal swabs should be combined in same UTM/VTM *Flocked swabs recommended, alternatively dacron or rayon swabs.	Transport in cooler box with ice packs. Sample should reach laboratory within 72hrs	Respiratory Panel (viral) Please note that this will only be done as part of comprehensive diagnosis to investigate cause of severe respiratory illness or outbreaks of unknown cause	Real-Time PCR Note: routine diagnostic testing available at NHLS or private pathology laboratories	3 working days	Specimens should be submitted with CRDM specimen submission form available on NICD website	CRDM Results; 011-386 6404 Queries; 011 555 6390 011 386 6373 Prof Anne von Gottberg 011 555 0316

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
Meningitis/Encephalitis Screen (viral/bacterial): <i>Escherichia coli</i> K1 <i>Haemophilus influenzae</i> <i>Listeria monocytogenes</i> <i>Neisseria meningitidis</i> <i>Streptococcus agalactiae</i> <i>Streptococcus pneumoniae</i> Cytomegalovirus (CMV) Enterovirus (EV) Herpes simplex virus 1 (HSV-1) Herpes simplex virus 2 (HSV-2) Human herpesvirus 6 (HHV-6) Human parechovirus (HPeV) Varicella zoster virus (VZV) Cryptococcus <u>neoformans/gattii</u>	CSF (Cerebrospinal fluid)	Transport in cooler box with ice packs. Sample should reach laboratory as soon as possible	Viral	Real-Time PCR	24 hours	Specimens should be submitted with CRDM specimen submission form available on NICD website	CRDM 011 555 0315/7 (micro lab) 011 555 0387 (Dr Mignon du Plessis) 011 555 0352 (Dr Nicole Wolter) 011 555 0356 (Molecular)

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
SARS-CoV-2 COVID-19 (Coronavirus disease 2019)	Any respiratory sample e.g. Nasopharyngeal or tracheal aspirates Nasal, nasal mid-turbinate, nasopharyngeal and/or oropharyngeal swabs placed in UTM/VTM Sputum BAL Biopsies Other samples – discuss with Dr on call Swabs should be placed in universal / viral transport medium (UTM/VTM). Nasopharyngeal- and oropharyngeal swabs should be combined in same UTM/VTM Flocked swabs recommended. Alternatively, dacron or rayon swabs	Transport in cooler box with ice packs. Sample should reach laboratory within 72hrs	SARS-CoV-2 COVID-19	Real-Time PCR Note: routine diagnostic testing available at NHLS or private pathology laboratories	2 working days	Specimens should be submitted with CRDM specimen submission form available on NICD website Contact line list should be completed for contact tracing COVID-19 is classified as a Category 1 notifiable medical condition (NMC).	NICD Hotline 082 883 9920 CRDM Results: 011 386 6373/0356/6390 Queries: 011 555 6390/0356/0387/0352 Prof Anne von Gottberg 011 555 0316
Avian Influenza (H5, H7, H9)	Any respiratory sample (e.g. Nasopharyngeal or tracheal aspirates, nasal, nasopharyngeal and/or oropharyngeal swabs, sputum, biopsies, autopsies etc.) Swabs should be placed in universal / viral transport medium (UTM/VTM). Nasopharyngeal- and oropharyngeal swabs can be combined in same UTM/VTM Flocked swabs recommended. Alternatively, dacron or rayon swabs	Transport in cooler box with ice packs. Sample should reach laboratory within 72hrs Transported as Infectious substance (packaging instruction 602 of IATA).	Avian influenza Contact NICD hotline on 082 883 9920 for all suspected avian influenza requests before submitting samples	Real-Time PCR	24 hours	Clinical features consistent with Avian Influenza infection PLUS meets epidemiological criteria as outlined in AI Case Investigation Document (Contact Epidemiologist) Specimens should be submitted with CRDM specimen submission form available on NICD website	NICD Hotline 082 883 9920 CRDM 011 555 6390 (Dr Jinal Bhiman) 011 555 0316 (Prof Anne von Gottberg)

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
MERS-Coronavirus (MERS-CoV)	<p>Induced sputum, lung aspirates, combined nasopharyngeal and oropharyngeal swabs or nasopharyngeal aspirates in UTM/VTM, BAL and biopsies of respiratory tract tissues. Other samples - discuss with Dr on call.</p> <p>Swabs should be placed in universal / viral transport medium (UTM/VTM).</p> <p>Nasopharyngeal and oropharyngeal swabs should be combined in same UTM/VTM</p> <p>Flocked swabs recommended. Alternatively, dacron or rayon swabs</p>	<p>Transport in cooler box with ice packs.</p> <p>Sample should reach laboratory within 72hrs</p> <p>Transport as infectious substance (packaging instruction 602 of IATA Category A).</p>	<p>MERS-CoV</p> <p>Contact NICD hotline on 082 883 9920 for all suspected MERS coronavirus test requests</p> <p>Please note that this test will only be done if case definition was met and discussed with NICD Dr on call (see NICD hotline number)</p>	Real-Time PCR	24 hours	<p>Clinical features consistent with Avian Influenza infection PLUS meets epidemiological criteria as outlined in AI Case Investigation Document (Contact Epidemiologist)</p> <p>Specimens should be submitted with CRDM specimen submission form available on NICD website</p>	<p>NICD Hotline 082 883 9920</p> <p>CRDM 011 555 6390 (Dr Jinal Bhiman) 011 555 0316 (Prof Anne von Gottberg)</p>
Severe respiratory illness of unknown cause	<p>Any respiratory sample (e.g. Nasopharyngeal or tracheal aspirates, nasal, nasopharyngeal and/or oropharyngeal swabs, sputum, biopsies, autopsies etc.)</p> <p>Swabs should be placed in universal / viral transport medium (UTM/VTM).</p> <p>Nasopharyngeal- and oropharyngeal swabs should be combined in same UTM/VTM</p> <p>Flocked swabs recommended. Alternatively, dacron or rayon swabs</p>	<p>Transport in cooler box with ice packs.</p> <p>Sample should reach laboratory within 72hrs</p>	<p>Respiratory panel (bacterial and viral)</p> <p>Contact NICD hotline on 082 883 9920</p>	Real-time PCR	24hours	Specimens should be submitted with CRDM specimen submission form available on NICD website	

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
<p><i>Neisseria</i> species, <i>Haemophilus</i> species and <i>Streptococcus</i> species (GERMS surveillance)</p> <p>Bacterial meningitis screen - diagnostic (<i>N. meningitidis</i>, <i>H. influenzae</i>, <i>S. pneumoniae</i> - routinely done),(Group B <i>Streptococcus</i>*, <i>E. coli</i>*, <i>S. aureus</i>*, <i>L. monocytogenes</i>*)</p> <p>*on request</p>	<p>Isolates inoculated onto Dorset transport medium and incubated overnight at 37°C in 5% CO₂</p> <p>Normally sterile site specimen (blood or cerebrospinal fluid [CSF] or other fluids e.g. pleural, peritoneal, synovial)</p> <p>Isolates inoculated onto Dorset transport medium and incubated overnight at 37°C in 5% CO₂</p> <p>Clinical samples: minimum volume of 200µl, EDTA blood</p>	<p>Submit isolates on Dorset transport media after overnight incubation.</p> <p>Inoculate as per NIC0184.</p> <p>Do not refrigerate. Do not batch for longer than 1 week as isolates will lose viability.</p> <p>Clinical specimens at cool ambient temperature (e.g. in a cooler box)</p>	<p>Identification of <i>Neisseria</i> species, <i>Haemophilus</i> species or <i>Streptococcus</i> species for GERMS surveillance (GERMS-SA)</p> <p>Bacterial meningitis screen (PCR and culture)</p>	<p>PCR identification of <i>S. pneumoniae</i>, <i>N. meningitidis</i>, <i>H. influenzae</i> Group B <i>Streptococcus</i>, <i>E. coli</i>, <i>S. aureus</i>, <i>L. monocytogenes</i></p> <p>Serotyping/ Serogrouping of <i>S. pn</i>, Nm, Hi (phenotypic and molecular)</p> <p>Antimicrobial susceptibility testing (disc and minimum inhibitory concentration - MIC)</p>	<p>PCR urgent 1-2 days</p> <p>Surveillance 1-2 months</p>	<p>Isolates submitted to NICD for national surveillance (GERMS-SA). Submit with sterile isolate form or LIS report.</p> <p>Specimens should be submitted with CRDM specimen submission form available on NICD website</p>	<p>NICD Hotline 082-883-9920</p> <p>CRDM 011 555 0315 (micro) 011 555 0317 (micro) 011 555 0356 (molecular) 011 555 0387 (Dr Mignon du Plessis) 011 555 0352 (Dr Nicole Wolter)</p>
<p>Atypical pneumonia-causing pathogens (<i>Mycoplasma pneumoniae</i>, <i>Chlamydia pneumoniae</i>, <i>Legionella</i> spp., <i>Bordetella</i> spp.)</p>	<p>Sputum (expectorated or induced) in a sterile container, and/or nasopharyngeal- oropharyngeal aspirate or combined flocked swabs preferably in universal transport medium (UTM) or Primestore.</p> <p>For <i>Bordetella pertussis</i> culture, swabs should be placed in Regan Lowe (RL)</p> <p>Urine (for <i>L. pneumophila</i> serogroup 1). Minimum volume 5ml in a sterile leak-proof container</p>	<p>Sputum to be frozen immediately and transported on dry ice</p> <p>UTM and urine to be stored and transported refrigerated. Primestore – ambient temperature</p> <p>RL to be stored and transported at room temperature</p>	<p>Atypical pneumonia PCR</p> <p><i>Bordetella pertussis</i></p> <p><i>Legionella</i> spp.</p>	<p>PCR for identification of atypical pneumonia-causing bacteria (<i>M. pneumoniae</i>, <i>C. pneumoniae</i>, and <i>Legionella</i> spp)</p> <p>PCR for <i>B. pertussis</i> and other <i>Bordetella</i> spp.</p> <p>Culture for <i>Legionella</i> spp., <i>B. pertussis</i> and <i>Bordetella</i> spp.</p> <p>Binax NOW for <i>Legionella pneumophila</i> (serogroup 1)</p>	<p>PCR urgent 1 day/24 hours</p> <p>Non-urgent 2-4 days</p> <p>Culture 7-10 days</p>	<p>Specimens should be submitted with CRDM specimen submission form available on NICD website</p>	<p>CRDM 011 555 0315 011 555 0317 011 555 0352 (Dr Nicole Wolter) 011 555 0316 (Prof Anne von Gottberg)</p>

Pathogen Species Name (disease/ syndrome, ICD-code)	Samples Collection (E.g. Sample source, type amount, specimen container, transport medium, type of swab)	Specimen Transportation Requirements (Temperature requirements, Stability, Minimum time to reach laboratory)	Tests To Request	Available Tests (Methodology)	Turn-around time	Special Instructions (Case investigations forms to be completed, guidelines, other relevant info)	Department and Contact telephone no's
<i>Corynebacterium diphtheriae</i> (Diphtheria)	Respiratory: Throat swab, pseudomembrane scraping or tissue Cutaneous: skin scrapings/lesions	Isolates on Hoyles or Dorset transport medium for identification Clinical samples (swabs) in Amies transport medium (for culture + PCR) Stored and transported at cool ambient temperature.	<i>Corynebacterium diphtheriae</i> identification Elek and PCR for toxin production	Phenotypic identification (MALDI-ToF) Elek for toxin production and PCR for presence of toxin gene	PCR urgent 1 day/24 hours Non-urgent 2-4 days Culture 3-5 days	Specimens should be submitted with CRDM specimen submission form available on NICD website	CRDM 011 555 0315 011 555 0317 011 555 0327 (Linda de Gouveia) 011 555 0316 (Prof Anne von Gottberg)
<i>Streptococcus pyogenes</i> (Group A Streptococcus)	Respiratory: Throat or nasopharyngeal swab Cutaneous: skin scrapings/lesions Normally sterile site specimen (blood or cerebrospinal fluid [CSF] or other fluids e.g. pleural, peritoneal, synovial)	Isolates on blood agar or Dorset transport medium for identification Clinical samples (swabs) in Amies transport medium (for culture + PCR). Stored and transported at cool ambient temperature.	<i>Streptococcus pyogenes</i> (Group A Streptococcus)	Phenotypic identification (MALDI-ToF) PCR identification of <i>S.pyogenes</i> Antimicrobial susceptibility testing (disc and minimum inhibitory concentration - MIC)	PCR urgent 1 day/24 hours Non-urgent 2-4 days Culture 3-5 days	Specimens should be submitted with CRDM specimen submission form available on NICD website	CRDM 011 555 0315 011 555 0317 011 555 0327 (Linda de Gouveia) 011 555 0316 (Prof Anne von Gottberg)
<i>Streptococcus agalactiae</i> (Group B Streptococcus)	Respiratory: Normally sterile site specimen (blood or cerebrospinal fluid [CSF] or other fluids e.g. pleural, peritoneal, synovial)	Isolates on blood agar or Dorset transport medium for identification Clinical samples (swabs) in Amies transport medium (for culture + PCR). Stored and transported at cool ambient temperature.	<i>Streptococcus agalactiae</i> (Group B Streptococcus)	Phenotypic identification (MALDI-ToF) PCR identification of <i>S.agalactiae</i> Serotyping/ Serogrouping of isolates Antimicrobial susceptibility testing (disc and minimum inhibitory concentration - MIC)	PCR urgent 1 day/24 hours Non-urgent 2-4 days Culture 3-5 days	Specimens should be submitted with CRDM specimen submission form available on NICD website	CRDM 011 555 0315 011 555 0317 011 555 0327 (Linda de Gouveia) 011 555 0316 (Prof Anne von Gottberg)

Public Health, Surveillance and Response

The Public Health Surveillance and Response Division includes the Outbreak Unit, the GERMS-SA surveillance programme, Travel Health. The division facilitates communication and data sharing between the national and provincial health departments and the NICD and the public and provides epidemiological input to other NICD units through collaborative projects and support of surveillance and epidemiological activities and outbreak responses.

Outbreak Response Unit

The Outbreak Response Unit (ORU) provides technical support for all aspects of communicable disease outbreaks and control in South Africa. Through close collaboration with provincial and national health departments and other stakeholders, together with systems for early detection and improved reporting of epidemic-prone communicable diseases, the ORU functions as a source of intelligence for outbreak detection and facilitates comprehensive outbreak response activities. In addition, close partnerships with NHLS diagnostic laboratories and NICD centres provide appropriate laboratory diagnostic services during outbreaks and specialised diagnostic testing as required.

Public Health Services

The ORU's role in outbreaks may include, but is not limited to, the following: outbreak detection and reporting, field investigation, development of clinical and laboratory guidelines, management of laboratory data and interpretation of results, and recommendations for prevention and control.

Contact Details

Dr Kerrigan Mc Carthy	+27 11 555 0542	079 8717278
24/7 Doctor on call/Outbreak Hotline	082 883 9920	

Please refer to the NICD website for further contact information:

<http://www.nicd.ac.za/index.php/centres/division-of-public-health-surveillance-and-response/>

GERMS-SA

Surveillance/Diagnostic Services

The GERMS-SA laboratory-based surveillance programme for diseases of public health importance is co-ordinated by a core team within the Division of Public Health Surveillance and Response and spans most of the centres at the NICD. Laboratory work is done through the respective Centre. The laboratory surveillance pathogens routinely include: *Streptococcus pneumoniae*, *Haemophilus influenzae*, *Neisseria meningitidis*, *Streptococcus pyogenes*, *Streptococcus agalactiae*, *Salmonella* Typhi, *Salmonella enterica* serotype Paratyphi (A, B and C) and *Vibrio cholerae*, *Cryptococcus* spp., carbapenem-resistant Enterobacteriaceae (CRE) and for outbreak-related *Salmonella* non-Typhi, *Shigella* spp, non-cholera *Vibrio*, *Campylobacter* spp, Diarrheoagenic *E.coli* and *Listeria* spp. GERMS-SA is an active surveillance programme and relies not only on participating laboratories to submit isolates,

but also makes use of the NHLS Corporate Data Warehouse to ensure that all cases that meet the case definition are included in the various surveillance databases.

The aim of GERMS-SA is to use the data to inform and guide public health policymakers in their decisions. The objectives include estimating the burden of both community- and hospital-acquired infectious diseases under surveillance; monitoring antimicrobial susceptibility trends; monitoring HIV-associated opportunistic infections; and evaluating the impact of vaccines included in the Expanded Programme on Immunisation (EPI). GERMS-SA's work is funded through the NICD/DoH.

GERMS-SA clinic-based surveillance (STI, HIV and TB)

GERMS-SA has expanded to include clinic-based surveillance. Sites have been initiated in all provinces. Clinic-based surveillance includes TB and HIV surveillance to describe the epidemiology of drug resistance among HIV-infected persons initiating ART and/or TB treatment at the selected sites, as well as undertake STI surveillance. The STI component includes surveillance of STI syndrome aetiologies, gonococcal antimicrobial resistance and HPV genotypes among patients attending the clinic's STI.

The Acute Febrile Illness Surveillance Project continues to be incorporated into clinic-based syndromic surveillance at one clinic site in rural Mpumalanga. This surveillance is a One-Health project and takes place in collaboration with veterinary practitioners and researchers from the University of Pretoria Veterinary Faculty. The aim is to describe the prevalence of zoonotic infections in adult patients presenting with acute febrile illness and for whom the clinic sisters would do a malaria test. Laboratory testing includes PCR and serology for brucellosis, bartonella infections, leptospirosis, Q-fever, tick bite fever, West Nile virus, Sindbis, Rift Valley fever and chikungunya virus infections.

Travel Health:

Travel Health was established in 2008 with the aim of being a centre for travel health-related activities and consulting on international health matters. This working group provides pre- and post-travel health advice and consultations for travel health practitioners, as well as for staff of the NHLS and NICD for work related field activities. Regular expert consultations are provided both locally and internationally to international focal points, institutes and health practitioners for South African travellers presenting with infectious diseases after travel within southern Africa and further abroad. These include diseases in travellers such as trypanosomiasis, severe malaria and rickettsial disease (tick bite fever). Consultation is also provided for the elimination of malaria in South Africa and other countries within southern Africa.

NICD Sequencing Core Facility

The NICD Sequencing Core Facility was established in January 2016 to promote and expedite research and surveillance activities at the NICD aimed at providing accurate, high quality and cost-effective next generation sequencing (NGS) solutions. The Sequencing Core Facility currently supports all centres at the NICD in terms of NGS and bioinformatics needs, thus acts as an extension of every centre with regards to NGS capacity. Since its inception, the core facility has continually engaged in several projects from design to completion.

Some key focus areas involve whole genome sequencing (de novo and re-sequencing), custom amplicon sequencing and metagenomics (viral and bacterial). The NICD Sequencing Core Facility currently supports three Illumina MiSeq sequencers. In addition to NGS, the core facility has a dedicated server (genomics server) for data analysis. The genomics server offers secure, powerful, and flexible bioinformatic computing accessible to all NICD scientists. CLC Genomics Server Core aims to provide a unique and stable software architecture core that makes it possible to apply a range of bioinformatics analysis-solutions on high-throughput sequencing data.

Important Considerations:

Sample Requirements: For NGS, accepted sample types include genomic DNA (gDNA), complementary DNA (cDNA), PCR products and amplicons. No original clinical specimens (e.g. swabs, blood, urine, stools, etc.) are accepted. Nucleic acid material (DNA or RNA) can be prepared from any sample source. DNA must be eluted in Tris-Cl buffer, pH 8.5 (not Tris-EDTA buffer). RNA must be eluted in nuclease-free ultra-pure water (not Tris-EDTA buffer) followed by conversion into cDNA. PCR products and cDNA preparations must be purified (recommended) using Agencourt® AMPure® beads or Zymo DNA Clean & Concentrator™-5 kit. **DNA Input Recommendations:** > 100 ng of DNA, non-degraded and free of particulate matter, double-stranded DNA > 70 bp; OD_{260/280} > 1.8, minimum concentration 10 ng/μL in 10mM Tris-Cl, pH 8.5. **RNA Input Recommendations:** 1-4 μg of total RNA, minimum concentration 20 ng/μL in nuclease-free ultra-pure water (do not exceed 50 μL) non-degraded with a Bioanalyzer RIN value of 8. When submitting samples, a Sample Request Form needs to be completed in full and emailed to the laboratory.

Sample Packaging and Transportation: Samples should be submitted to the laboratory in sterile tubes or reaction plates, and sample leak during transportation should be avoided. It is recommended that the cold chain be maintained during sample shipment in order to ensure that nucleic acid integrity is not compromised. Samples should be dispatched in a sealed container and marked appropriately. Courier service must be employed for sample shipment. Dispatching of samples on Fridays and weekends should be avoided so as to keep transportation time to the absolute minimum.

Expected Turnaround Time: Samples have an expected turnaround time of up to two weeks (project-dependent), and this is calculated from sample reception in the laboratory until the sequencing data is sent to the client. Urgent samples (i.e. samples from suspected outbreak cases) are prioritized, and will therefore have a shortened turnaround time. It is imperative that the laboratory is notified of priority samples before they are dispatched.

NGS Applications: The core facility engages in various NGS projects from design to completion, and the key focus areas include whole genome sequencing (de novo and re-sequencing), custom amplicon sequencing, whole transcriptome sequencing, RNA sequencing and metagenomics (viral and bacterial).

Sample Rejection Criteria: All incoming samples are recorded in the Sample Receiving Log Sheet and verification is done to ensure that each sample meets all acceptance criteria. Samples are rejected if they fall in any of the following rejection criteria: unlabelled samples; leaking samples; incorrect sample type (e.g. blood, sputum, urine, swab, etc.); samples not properly labelled; information on sample tube and Sample Request Form not corresponding; insufficient sample volume; samples with very low concentrations; samples received without completed Sample Request Form; sample tubes with illegible handwriting; and samples submitted for tests not performed in the laboratory.

Contact Details (Sequencing Core Facility):

E-mail address : sequencing@nicd.ac.za

Telephone no. : 011 386 6459 / 011 386 6322 / 011 555 0450

Shipping address : ATT: Dr Arshad Ismail

Sequencing Core Facility

National Institute for Communicable Diseases

1 Modderfontein Road, Sandringham, 2192

Key Contact Staff – National Institute for Communicable Diseases

Department	Contact person	Contact numbers	Email address
24/7 NICD Emergency/Assistance cell phone		082 883 9920	
Executive Director	Prof L Morris	011 386-6332 Fax:011 386-6453	lynnm@nicd.ac.za
Deputy Director	Dr N Mayet	011 386 6038	nataliem@nicd.ac.za
Senior Administrator	Ms. C Singh	011 386-6058	charmains@nicd.ac.za
NICD Quality Assurance			
Quality Assurance Manager	Mr. H Julius	011 386 6095	henryj@nicd.ac.za
Centre for Vaccines and Immunology:			
Centre for Vaccines and Immunology: Head	Dr M Suchard	011 386 6387 083 671 3909	melindas@nicd.ac.za
Laboratory Manager Hepatitis	Dr N Prabdial Sing	011 386-6387	niships@nicd.ac.za
Laboratory Manager Polio Molecular	Mrs. S Moonsamy	011 555-0504	shelinam@nicd.ac.za
Laboratory Manager Measles	Mrs. S Smit	011 386 6343	sheilaghs@nicd.ac.za
Poliovirus Serology and Isolation	Mrs. S Moonsamy	011 386-6422	shelinam@nicd.ac.za
Centre for Tuberculosis:			
Centre for Tuberculosis: Head		011- 885 5321	
Pathologist	Dr F Ismail	011-885 5323	farzanai@nicd.ac.za

Department	Contact person	Contact numbers	Email address
Head Scientist	Dr S V Omar	011-885 5343	shaheedvo@nicd.ac.za
Laboratory Manager	Mrs. C De Abreu	011 885 5316 /5317	ceceliad@nicd.ac.za
Centre for HIV and STI			
Centre for HIV and STI : Head	Prof A Puren	011 386-6328 082 908 8048	adrianp@nicd.ac.za
HIV Molecular and Serology Diagnostic Laboratories: Head	Prof A Puren	011 386-6328 082 908 8048	adrianp@nicd.ac.za
Secretary	Mrs. M van Rensburg	011 386 6462	monicavr@nicd.ac.za
Laboratory Supervisor HIV Molecular Diagnostics	Mrs. E Cutler	011 386-6439	ewaldec@nicd.ac.za
Laboratory Supervisor HIV Serology Diagnostics	Mrs. B Singh	011 386- 6437 011 386 6435	beverleys@nicd.ac.za
Virology Laboratory	Prof L Morris	011 386-6332 Fax:011 386-6453	lynnm@nicd.ac.za
Personal Assistant	Mrs. C Kriel	011 386-6362	carinak@nicd.ac.za
Laboratory Manager	Mrs. L Nxumalo	011 386 6341	Livhuwanin@nicd.ac.za
Cell Biology Unit	Prof C Tiemessen	011 386-6366 082 883 6663	carolinet@nicd.ac.za
Sexually Transmitted Infections Reference Section: Head	Dr R Kularatne	011 555 0468 082 442 1348	ranminik@nicd.ac.za
Laboratory Manager	Ms. V Maseko	011 555 0461 083 406 4848	vanessam@nicd.ac.za

Department	Contact person	Contact numbers	Email address
Centre for Emerging, Zoonotic and Parasitic Diseases			
Centre for Emerging, Zoonotic and Parasitic Diseases: Head	Prof J Paweska	011 386-6336 082 908 8046	januszp@nicd.ac.za
Secretary	Ms. N Mphula	011 386 6382	nondumisom@nicd.ac.za
Vector Control Reference Laboratory	Prof B Brooke	011 386-6480 Fax:011 386 6481	basilb@nicd.ac.za
Parasitology Reference Laboratory	Prof J Freaan	011 555 0308 082 906 2856	johnf@nicd.ac.za
Laboratory Manager	Mrs. B Moodley	011 555 0311	bhavanim@nicd.ac.za
Special Viral Pathogens Laboratory			
Special Viral Pathogens Reference Laboratory			
Head Scientist	Dr J Weyer	011 386 6376 011 386 6339 082 903 9131	jacquelinew@nicd.ac.za
Head Scientist	Dr O Hellferscee	011 386 6424/6353 082 908 8045	orienkah@nicd.ac.za
Special Bacterial Pathogens Laboratory			
Head Scientist	Dr J Rossouw	011 555 0331	jennyr@nicd.ac.za
Electron Microscope Laboratory			
Head Scientist	Dr M Birkhead	011 386 6318	monicab@nicd.ac.za
Centre for Respiratory Diseases and Meningitis			

Department	Contact person	Contact numbers	Email address
CRDM : Head	Prof C Cohen	011 386-6593 082 803 8093	cherylc@nicd.ac.za
Clinical microbiologist	Prof A von Gottberg	011 555-0316 082 572 0057	annev@nicd.ac.za
Laboratory Manager	Mrs. L De Gouveia	011 555 0327	lindad@nicd.ac.za
National Influenza Centre	Dr. J Bhiman	011 386 6390	jinalb@nicd.ac.za
Principal Scientist	Dr M du Plessis	011 555 0387	mignond@nicd.ac.za
Virus Isolation / Serology	Ms. C Fourie	011-386-6412	cardiaf@nicd.ac.za
	Mrs. A Buys	011 386-6373	ameliab@nicd.ac.za
Centre for Enteric Diseases			
Centre for Enteric Diseases: Head	Dr J Thomas	0113866452 /0731708874	junot@nicd.ac.za
Virology Laboratory	Prof N Page	011 555-0370 082 447 2745	nicolap@nicd.ac.za
Bacteriology Laboratory		011 386-6269 082 809 5667	
Laboratory Manager	Ms H Ngomane	011 386 6235	mimmyn@nicd.ac.za
Centre for Healthcare-Associated Infections, Antimicrobial Resistance and Mycoses (CHARM)			
CHARM: Head	Prof N Govender	Fax:011 555 0353	neleshg@nicd.ac.za

Department	Contact person	Contact numbers	Email address
Mycology Laboratory	Prof N Govender	Fax:011 555 0353	neleshg@nicd.ac.za
Laboratory Manager Mycology	Ms. R Mpenbe	011 555 0325	ruthm@nicd.ac.za
Antimicrobial Resistance Laboratory and Culture Collection	Prof O Perovic	011 386 6278	olgap@nicd.ac.za
Laboratory Manager	Mrs. M Smith	011 555 0342	marshagnes@nicd.ac.za
National Stock Culture Collection	Mrs. M Smith	011 555 0342	marshagnes@nicd.ac.za
Division of Public Health, Surveillance and Response			
Public Health Surveillance and Response: Head	Dr K Mc Carthy	0798717278 0115550542	kerriganm@nicd.ac.za
Public Health Surveillance and Response	Dr V Quan	011 386 6012	vanessaq@nicd.ac.za
Notifiable Medical Conditions (NMC)	Dr P Mutevedzi	072 621 3805	portiamu@nicd.ac.za
Sequencing Core Facility			
Sequencing Core Facility	Dr Arshad Ismail	011 386 6459 072 98 5275	arshadi@nicd.ac.za
Specimen Receiving Office			
NICD Specimen receiving office	Ms. B Masengemi	011 386 6314	BusisiweM@nicd.ac.za

Acknowledgement of Reading Form

Document number: _____ Version Number: _____

Title: _____

My signature confirms that I have read and understood the content of this document and relevant kit insert (where applicable).

Name	Signature	Date

Note to the Quality Rep: - This form must be filed for 5 years to provide audit traceability.