

SOUTH AFRICA WEEK 4 2021

OVERVIEW OF REPORT

This report summarises national laboratory testing for SARS-CoV-2, the virus causing COVID-19, in South Africa. This report is based on data for specimens reported up to 30 January 2021 (Week 4 of 2021).

HIGHLIGHTS

- In the period 1 March 2020 through 30 January 2021, 8,234,743 laboratory tests for SARS-CoV-2 have been performed nationally.
- The number of tests performed in week 4 of 2021 (n=230,376) was lower than performed in the previous few weeks.
- KwaZulu-Natal (513 per 100,000 persons), Gauteng (476 per 100,000 persons), Northern Cape (467 per 100,000 persons) and Western Cape (452 per 100,000 persons) provinces had the highest testing rates in week 4 of 2021.
- In week 4 of 2021 the percentage testing positive was 14.0%, lower than has been observed in the previous few weeks.
- Percentage testing positive in week 4 of 2021 was highest in Limpopo (25.6%) and Mpumalanga (20.9%) provinces. Percentage testing positive was 10%-20% in the Western Cape, Northern Cape, KwaZulu-Natal, North West, Free State and Gauteng, and was <10% in the Eastern Cape.
- In week 4 of 2021, compared to the previous week, the percentage testing positive decreased in all provinces.
- Mean laboratory turnaround time in week 4 of 2021 was 2.0 days; 3.2 days in the public sector and <1 day in the private sector.
- Integration of antigen test data from the different settings where these tests are being used into the national database is ongoing, leading to some underestimation of testing volumes.

SOUTH AFRICA | WEEK 4 2021

Methods

Testing for SARS-CoV-2 began on 28 January 2020 at the NICD and after the first case was confirmed on 5th March 2020, testing was expanded to a larger network of private and NHLS laboratories. Laboratory testing was conducted for people meeting the case definition for persons under investigation (PUI). This definition was updated several times over the reporting period but at different times included (i) symptomatic individuals seeking testing, (ii) hospitalised individuals for whom testing was done, (iii) individuals in high-risk occupations, (iv) individuals in outbreak settings, and (v) individuals identified through community screening and testing (CST) programmes which were implemented in April 2020 and was discontinued from the week beginning 17th May. CST was implemented differently in different provinces, and ranged from mass screening approaches (including asymptomatic individuals) to screening of individuals in contact with a confirmed case to targeted testing of clusters of cases. Respiratory specimens were submitted to testing laboratories. Testing was performed using reverse transcriptase real-time PCR, which detects SARS-CoV-2 viral genetic material. Laboratories used any one of several in-house and commercial PCR assays to test for the presence of SARS-CoV-2 RNA. Testing for SARS-CoV-2 using rapid antigen-based tests was implemented towards the end of October 2020. Results of reported rapid antigen-based tests are included in this report, however data are incomplete and efforts are ongoing to improve data completeness.

Test results were automatically fed into a data warehouse after result authorisation. We excluded specimens collected outside South Africa and duplicate entries of the same test for an individual. From week 48 of 2020 onwards, test data were reported from the Notifiable Medical Conditions Surveillance System (NMCSS). Date of specimen receipt in the laboratory was used when date of specimen collection was missing. Proportion testing positive (PTP) was calculated as the number of positive tests/total number of tests and presented as percentage by multiplying with 100. We used 2020 mid-year population estimates from Statistics South

Africa to calculate the testing rate, expressed as tests per 100,000 persons. Patient admission status for public and private sector tests was determined based on the reported patient facility. Laboratory turnaround times were calculated for PCR-based tests and were calculated as the mean number of days between specimen collection and reporting of the result. Categorical variables were compared using the chi-squared test, and continuous variables with the students t-test, with a P-value<0.05 considered statistically significant.


Health district and sub-district (in the metros) level results were mapped based on geo-locatable public and private sector testing facilities. Estimates of overall prevalence were derived using regression techniques. Estimates were adjusted to produce district-specific positive test prevalences based on the national average age and sex profile of testing for that week. This adjustment allows more accurate comparison of the proportion testing positive across districts.

The report includes tests conducted between 1 March 2020 (week 10 of 2020), the week when the first case of COVID-19 was confirmed, and 30 January 2021 (week 4 of 2021).

Testing volumes and proportion testing positive

From 1 March 2020 through 30 January 2021, 8,234,743 laboratory tests (PCR and antigen tests) for SARS-CoV-2 were performed. The number of tests performed increased weekly from week 10 of 2020, with the highest number of tests performed during the first wave occurring in week 28 of 2020 (n=307,900), and subsequently decreased. Weekly testing volumes increased again from week 47 (beginning 15 November 2020), with the highest weekly number of tests performed since the start of the pandemic performed in week 1 of 2021 (n=492,346). In week 4 of 2021, 230,376 tests were performed, lower than the previous few weeks. All tests for samples collected in the previous week may not yet be reflected. Reduced testing volumes were observed over weekends and public holidays (Figure 1).

SOUTH AFRICA | WEEK 4 2021



DATE OF SPECIMEN COLLECTION

Figure 1. Number laboratory tests conducted by date of specimen collection, South Africa, 1 March 2020 – 30 January 2021. Blue dotted line shows the 7-day moving average of the number of tests conducted. Grey bars highlight weekend days and public holidays.

The overall percentage testing positive from week 10 of 2020 through week 4 of 2021 was 18.6% (Table 1). During the first wave of infections, the percentage testing positive increased week on week from week 18 to a peak of 29.7% in week 29 of 2020, and subsequently decreased to 8.4% in week 44 of 2020. During the second wave of infections the percentage testing positive started increasing from week 46 of 2020, to a peak of 34.9% in week 53 of 2020. The percentage testing positive in week 4 of 2021 was 14.0%, lower than has been observed in the previous few weeks (Figure 2).

Table 1. Weekly number of tests conducted and positive tests, South Africa, 1 March 2020 – 30 January 2021

34/ I I	Maria da la casta astrono	No 6 to - to (0/)	No. of a california and	B
Week number	Week beginning	No. of tests n (%)	No. of positive tests	Percentage testing positive (%)
10	01-Mar-20	453 (0.0)	13	2.9
11	08-Mar-20	2380 (0.0)	103	4.3
12	15-Mar-20	21567 (0.3)	897	4.2
13	22-Mar-20	17541 (0.2)	543	3.1
14	29-Mar-20	18245 (0.2)	520	2.9
15	05-Apr-20	26298 (0.3)	796	3.0
16	12-Apr-20	43748 (0.5)	1295	3.0
17	19-Apr-20	79174 (1.0)	2177	2.7
18	26-Apr-20	93807 (1.1)	3203	3.4
19	03-May-20	142700 (1.7)	6017	4.2
20	10-May-20	165368 (2.0)	8090	4.9
21	17-May-20	166534 (2.0)	11379	6.8
22	24-May-20	156133 (1.9)	12967	8.3
23	31-May-20	153562 (1.9)	15079	9.8
24	07-Jun-20	173890 (2.1)	22358	12.9
25	14-Jun-20	186044 (2.3)	32638	17.5
26	21-Jun-20	252082 (3.1)	55044	21.8
27	28-Jun-20	302693 (3.7)	75306	24.9
	05-Jul-20	307900 (3.7)	86032	27.9
	12-Jul-20	285586 (3.5)		29.7
<u></u>	19-Jul-20	270880 (3.3)		29.0
<u></u>	26-Jul-20	216373 (2.6)		27.0
32	02-Aug-20	179560 (2.2)	40991	22.8
<u></u>	09-Aug-20	141072 (1.7)	26262	
<u></u>	16-Aug-20	135004 (1.6)	21374	15.8
3 <u>5</u>	23-Aug-20	123327 (1.5)	16330	
35 36		112755 (1.4)	12790	
36 37		116991 (1.4)	11952	10.2
	13-Sep-20	120706 (1.5)	12011	10.2 10.0
	20-Sep-20			10.2
39		98813 (1.2)	10098	
40	27-Sep-20	123047 (1.5)	11006	8.9
41	04-Oct-20	131026 (1.6)	11777	9.0
42	11-Oct-20	137949 (1.7)	12072	8.8
43	18-Oct-20	142144 (1.7)	12066	8.5
44	25-Oct-20	135826 (1.6)	11477	8.4
45	01-Nov-20	138785 (1.7)	12132	8.7
<u>46</u>	08-Nov-20	146952 (1.8)	14838	10.1
47	15-Nov-20	160608 (2.0)	18760	11.7
48	22-Nov-20	175641 (2.1)	22042	12.5
49	29-Nov-20	202939 (2.5)	30761	15.2
50	06-Dec-20	266846 (3.2)	53288	20.0
51	13-Dec-20	293234 (3.6)	68529	23.4
52	20-Dec-20	283004 (3.4)	81858	28.9
53	27-Dec-20	330571 (4.0)	115331	34.9
11	03-Jan-21	492346 (6.0)	150008	30.5
2	10-Jan-21	409923 (5.0)	104091	25.4
3	17-Jan-21	322340 (3.9)	62784	19.5
4	24-Jan-21	230376 (2.8)	32304	14.0
	Total	8234743 (100.0)	1533337	18.6

SOUTH AFRICA | WEEK 4 2021




Figure 2. Percentage of laboratory tests positive for SARS-CoV-2 by date of specimen collection, South Africa, 1 March 2020 – 30 January 2021. Blue dotted line shows the 7-day moving average of the percentage testing positive. Grey bars highlight weekend days and public holidays.

Testing in private and public sectors

From 1 March 2020 through 30 January 2021, 3,570,171 laboratory tests were conducted in public sector laboratories, with 19.3% testing positive. Over this same period, private sector laboratories conducted 4,664,572 tests, with 18.1% testing positive (Table 2). Overall the public sector has conducted 43.4% of tests and accounted for 45.0% of positive tests. In the first wave of infections the peak percentage testing positive was observed in week 30 of 2020 in the public sector (28.8%), and in week 29 of 2020 in the private sector (30.6%). In the second wave of infections the highest percentage testing positive was observed in week 53 of 2020 in both the public sector (35.5%) and private sector (34.4%). From week 3 to week 4 of 2021, the percentage testing positive

decreased by 5.7% in the public sector (20.8% to 15.1%, P<0.001), and decreased by 5.0% (18.0% to 13.0%, P<0.001) in the private sector. In week 4 of 2021 the percentage testing positive was higher in the public sector (15.1%) compared to the private sector (13.0%) (P<0.001).

The mean turnaround time for PCR tests conducted in week 4 of 2021 was 2.0 days. Turnaround time decreased slightly in the public sector (3.2 days) and remained consistent in the private sector (0.8 days) (Figure 3). Turnaround times for public sector tests were highest in KwaZulu-Natal (5.2 days), Mpumalanga (5.0 days), North West (4.1 days) and Limpopo (3.6 days) provinces in week 4 (Figure 4). Fourteen of the 28 (50%) NHLS laboratories performing PCR testing for SARS-CoV-2 had turnaround times ≤2 days (Figure 5).

SOUTH AFRICA | WEEK 4 2021

Table 2. Weekly number of tests conducted and positive tests, by healthcare sector, South Africa, 1 March 2020 – 30 January 2021

		Publi	Public sector Private sector		Public sector percentage of		Ratio	
Week	Week	Tests	Cases	Tests	Positive tests	Tests (%)	Positive tests	of PTP ^a
number_	beginning	rests	n (%)	Tests	n (%)		(%)	<i>) 16</i> []
10	01-Mar-20	293	10 (3.4)	160	3 (1.9)	64.7	76.9	1.820
11	08-Mar-20	401	27 (6.7)	1979	76 (3.8)	16.8	26.2	1.753
12	15-Mar-20	1442	81 (5.6)	20125	816 (4.1)	6.7	9.0	1.385
13	22-Mar-20	3477	149 (4.3)	14064	394 (2.8)	19.8	27.4	1.530
14	29-Mar-20	5868	194 (3.3)	12377	326 (2.6)	32.2	37.3	1.255
15_	05-Apr-20	11735	417 (3.6)	14563	379 (2.6)	44.6	52.4	1.365
16	12-Apr-20	24166	672 (2.8)	19582	623 (3.2)	55.2	51.9	0.874
17	19-Apr-20	55110	1595 (2.9)	24064	582 (2.4)	69.6	73.3	1.197
18	26-Apr-20	67468	2453 (3.6)	26339	750 (2.8)	71.9	76.6	1.277
19	03-May-20	94336	4506 (4.8)	48364	1511 (3.1)	66.1	74.9	1.529
20	10-May-20	107997	5443 (5.0)	57371	2647 (4.6)	65.3	67.3	1.092
21	17-May-20	98647	7031 (7.1)	67887	4348 (6.4)	59.2	61.8	1.113
22	24-May-20	77596	6411 (8.3)	78537	6556 (8.3)	49.7	49.4	0.990
23	31-May-20	63943	6626 (10.4)	89619	8453 (9.4)	41.6	43.9	1.099
24	07-Jun-20	64653	8038 (12.4)	109237	14320 (13.1)	37.2	36.0	0.948
25	14-Jun-20	61147	11982 (19.6)	124897	20656 (16.5)	32.9	36.7	1.185
26	21-Jun-20	90452	20425 (22.6)	161630	34619 (21.4)	35.9	37.1	1.054
27	28-Jun-20	106366	27244 (25.6)	196327	48062 (24.5)	35.1	36.2	1.046
28	05-Jul-20	117722	32238 (27.4)	190178	53794 (28.3)	38.2	37.5	0.968
29	12-Jul-20	110658	31383 (28.4)	174928	53541 (30.6)	38.7	37.0	0.927
30	19-Jul-20	105206	30319 (28.8)	165674	48314 (29.2)	38.8	38.6	0.988
31	26-Jul-20	81234	22782 (28.0)	135139	35609 (26.3)	37.5	39.0	1.064
32	02-Aug-20	70566	16996 (24.1)	108994	23995 (22.0)	39.3	41.5	1.094
33	09-Aug-20	58660	11172 (19.0)	82412	15090 (18.3)	41.6	42.5	1.040
34	16-Aug-20	56136	9621 (17.1)	78868	11753 (14.9)	41.6	45.0	1.150
35	23-Aug-20	50317	7790 (15.5)	73010	8540 (11.7)	40.8	47.7	1.324
36	30-Aug-20	45419	6096 (13.4)	67336	6694 (9.9)	40.3	47.7	1.350
37	06-Sep-20	51054	6421 (12.6)	65937	5531 (8.4)	43.6	53.7	1.499
38	13-Sep-20	53704	6547 (12.2)	67002	5464 (8.2)	44.5	54.5	1.495
39	20-Sep-20	44839	5530 (12.3)	53974	4568 (8.5)	45.4	54.8	1.457
40	27-Sep-20	48626	5567 (11.4)	74421	5439 (7.3)	39.5	50.6	1.566
41	04-Oct-20	50428	5688 (11.3)	80598	6089 (7.6)	38.5	48.3	1.493
42	11-Oct-20	53444	5702 (10.7)	84505	6370 (7.5)	38.7	47.2	1.415
43	18-Oct-20	56120	6044 (10.8)	86024	6022 (7.0)	39.5	50.1	1.538
44	25-Oct-20	51281	5721 (11.2)	84545	5756 (6.8)	37.8	49.8	1.639
45	01-Nov-20	53017	6067 (11.4)	85768	6065 (7.1)	38.2	50.0	1.618
46	08-Nov-20	58992	8110 (13.7)	87960	6728 (7.6)	40.1	54.7	1.797
47	15-Nov-20	67795	10702 (15.8)	92813	8058 (8.7)	42.2	57.0	1.818
48	22-Nov-20	74781	12243 (16.4)	100860	9799 (9.7)	42.6	55.5	1.685
49	29-Nov-20	81348	15833 (19.5)	121591	14928 (12.3)	40.1	51.5	1.585
50	06-Dec-20	107170	24734 (23.1)	159676	28554 (17.9)	40.2	46.4	1.291
51	13-Dec-20	116489	29862 (25.6)	176745	38667 (21.9)	39.7	43.6	1.172
52	20-Dec-20	109208	34168 (31.3)	173796	47690 (27.4)	38.6	41.7	1.140
53	27-Dec-20	148717	52757 (35.5)	181854	62574 (34.4)	45.0	45.7	1.031
	03-Jan-21	230103	70487 (30.6)	262243	79521 (30.3)	46.7	47.0	1.010
2	10-Jan-21	201928	53628 (26.6)	207995	50463 (24.3)	49.3	51.5	1.095
3	17-Jan-21	166963	34762 (20.8)	155377	28022 (18.0)	51.8	55.4	1.154
4	24-Jan-21	113149	17068 (15.1)	117227	15236 (13.0)	49.1	52.8	1.161
	Total	3570171	689342 (19.3)	4664572	843995 (18.1)	43.4	45.0	1.067

^aRatio of percentage testing positive (PTP) in the public sector to the private sector calculated as (no. of cases/total tests in public sector)/ (no. of cases/total tests in private sector)




Figure 3. Mean number of days between date of specimen collection and date of test result, by week of test result, South Africa, 3 – 30 January 2021




Figure 4. Mean number of days between date of specimen collection and date of test result, by week of test result and province, public sector, South Africa, 3 – 30 January 2021. WC, Western Cape; EC, Eastern Cape; FS, Free State; KZN, KwaZulu-Natal; GT, Gauteng; NC, Northern Cape; NW, North West; MP, Mpumalanga; LP, Limpopo

SOUTH AFRICA

WEEK 4 2021



Figure 5. Mean number of days between date of specimen collection and date of test result, by public sector laboratory, 10 – 30 January 2021. The horizontal black line indicates 48-hour turnaround time (TAT).

Testing by province

Gauteng (32.0%) performed the largest number of tests in week 4 of 2021, followed by KwaZulu-Natal (25.7%) and Western Cape (13.7%) provinces (Table 3). KwaZulu-Natal (513 per 100,000 persons), Gauteng (476 per 100,000 persons), Northern Cape (467 per 100,000 persons) and Western Cape (452 per 100,000 persons) provinces had the highest testing rates in week 4 of 2021 (Figure 6). Testing rates have decreased in all provinces over the past few weeks.

The percentage testing positive in week 4 of 2021 was highest in Limpopo (25.6%) and Mpumalanga (20.9%). Percentage testing positive was 10-20% in the Western Cape, Northern Cape, KwaZulu-Natal, North West, Free State and Gauteng, and was <10% in the Eastern Cape in week 4 of 2021 (Figure 7 and Table 3). Compared to the previous week, the percentage testing positive decreased in week 4 in all provinces (P≤0.001). The percentage testing positive was higher than the national average, not weighted for population size, in the Northern Cape, Free State, KwaZulu-Natal, North West, Mpumalanga and Limpopo (Figure 7).




Figure 6. Testing rate per 100,000 persons by province and week of specimen collection, South Africa, 1 March 2020 – 30 January 2021

SOUTH AFRICA WEEK 4 2021

Table 3. Weekly number of tests performed and positive tests, by province, South Africa, 10 - 30 January 2021

		10 –	l6 Jan 21	17 – 3	23 Jan 21	24 -	30 Jan 21	ď	
Province	Population ^a	No. of tests	No. positive tests (%)	No. of tests	No. positive tests (%)	No. of tests	No. positive tests (%)	Tests per 100,000 persons	Change in percentage positive ^b
Western Cape	7005741	50155	14311 (28.5)	38837	7469 (19.2)	31638	4356 (13.8)	452	-5.5%
Eastern Cape	6734001	27471	5205 (18.9)	21320	2785 (13.1)	15493	1365 (8.8)	230	-4.3%
Northern Cape	1292786	8648	2096 (24.2)	7483	1397 (18.7)	6039	1019 (16.9)	467	-1.8%
Free State	2928903	17967	3983 (22.2)	15122	2897 (19.2)	11090	1765 (15.9)	379	-3.2%
KwaZulu-Natal	11531628	108277	27503 (25.4)	88205	17555 (19.9)	59106	8423 (14.3)	513	-5.7%
North West	4108816	13215	3800 (28.8)	11810	2566 (21.7)	8503	1445 (17.0)	207	-4.7%
Gauteng	15488137	132601	29230 (22.0)	101508	17154 (16.9)	73746	8368 (11.3)	476	-5.6%
Mpumalanga	4679786	31451	9805 (31.2)	24306	6349 (26.1)	16229	3391 (20.9)	347	-5.2%
Limpopo	5852553	19942	8149 (40.9)	13628	4606 (33.8)	8464	2167 (25.6)	145	-8.2%
Unknown		196	9 (4.6)	121	6 (5.0)	68	5 (7.4)		
Total	59622350	482275	147878 (30.7)	404069	103052 (25.5)	290614	55134 (19.0)	487	-6.5%

a 2020 Mid-year population Statistics SA

b Current week compared to previous week




Figure 7. Weekly percentage testing positive, by province, South Africa, 10 – 30 January 2021. The horizontal blue line shows the national mean for week 4, beginning 24 January 2021.

Testing in the public sector

In the public sector, the percentage testing positive decreased in the past week (20.8% in week 3 to 15.1% in week 4 of 2021, P<0.001) (Table 4). The percentage testing positive in week 4 of 2021 was highest in

Limpopo (31.2%) and Mpumalanga (21.7%). The percentage testing positive in the public sector remains higher than the national average, not weighted for population size, in the Western Cape, Northern Cape, Free State, North West, Mpumalanga and Limpopo provinces (Figure 8).

SOUTH AFRICA WEEK 4 2021

Table 4. Weekly number of tests conducted and positive tests in the public sector, by province, South Africa, 10 – 30 January 2021

10-16 Jar		an 2021	17-23 i		24-30 J	an 2021
Province	No. of tests	No. positive tests (%)	No. of tests	No. positive tests (%)	No. of tests	No. positive tests (%)
Western Cape	22273	6906 (31.0)	16349	3610 (22.1)	12282	2189 (17.8)
Eastern Cape	20043	3693 (18.4)	15789	2062 (13.1)	10493	880 (8.4)
Northern Cape	5654	1396 (24.7)	4870	983 (20.2)	3977	736 (18.5)
Free State	11654	2727 (23.4)	9337	1971 (21.1)	5967	1104 (18.5)
KwaZulu-Natal	66010	16670 (25.3)	58994	12044 (20.4)	39389	5677 (14.4)
North West	6760	1999 (29.6)	6642	1517 (22.8)	4278	733 (17.1)
Gauteng	46859	11330 (24.2)	37974	7213 (19.0)	26684	3250 (12.2)
Mpumalanga	14415	5108 (35.4)	11336	3205 (28.3)	6805	1479 (21.7)
Limpopo	8258	3799 (46.0)	5672	2157 (38.0)	3274	1020 (31.2)
Unknown	2	0 (0.0)	0	0 (0.0)	0	0 (0.0)
Total	201928	53628 (26.6)	166963	34762 (20.8)	113149	17068 (15.1)




Figure 8. Weekly percentage testing positive in the public sector, by province, South Africa, 10 – 30 January 2021. The horizontal blue line shows the national mean for week 4 of 2021, beginning 24 January 2021.

Facilities with high proportions testing positive

Table 5.1 shows the 25 public sector clinics, hospitals and testing laboratories (where specimens were not tied to a particular facility), that had 25 or more

specimens tested and at least five positive results in the week of 24-30 January 2021, with the highest proportion testing positive nationally. This week's list is dominated by facilities in Limpopo (9) and KwaZulu-Natal (8).

SOUTH AFRICA WEEK 4 2021

Table 5.1 Public sector healthcare facilities with a high proportion testing positive, 24-30 January 2021

Facility Name	Province	Tests	PTP (95% CI)
Facility 1	Limpopo	27	0.667 (0.489;0.844)
Facility 2	Limpopo	40	0.625 (0.475;0.775)
Facility 3	KwaZulu-Natal	31	0.613 (0.441;0.784)
Facility 4	KwaZulu-Natal	28	0.607 (0.426;0.788)
Facility 5	Free State	46	0.565 (0.422;0.708)
Facility 6	Limpopo	25	0.560 (0.365;0.755)
Facility 7	KwaZulu-Natal	45	0.533 (0.388;0.679)
Facility 8	Limpopo	40	0.525 (0.370;0.680)
Facility 9	Limpopo	27	0.519 (0.330;0.707)
Facility 10	KwaZulu-Natal	35	0.514 (0.349;0.680)
Facility 11	Eastern Cape	55	0.509 (0.377;0.641)
Facility 12	North West	76	0.500 (0.388;0.612)
Facility 13	KwaZulu-Natal	88	0.500 (0.396;0.604)
Facility 14	Free State	53	0.491 (0.356;0.625)
Facility 15	North West	33	0.485 (0.314;0.655)
Facility 16	Mpumalanga	44	0.477 (0.330;0.625)
Facility 17	Limpopo	40	0.475 (0.320;0.630)
Facility 18	Limpopo	38	0.474 (0.315;0.632)
Facility 19	KwaZulu-Natal	28	0.464 (0.280;0.649)
Facility 20	KwaZulu-Natal	81	0.457 (0.348;0.565)
Facility 21	Free State	48	0.438 (0.297;0.578)
Facility 22	KwaZulu-Natal	53	0.434 (0.301;0.567)
Facility 23	Limpopo	83	0.434 (0.327;0.540)
Facility 24	Limpopo	79	0.430 (0.321;0.540)
Facility 25	Mpumalanga	28	0.429 (0.245;0.612)

Table 5.2 shows the 25 private sector clinics, hospitals and testing laboratories (where specimens were not tied to a particular facility), that had 25 or more specimens tested and at least five positive results in the week of 24-30 January 2021, with the highest proportion testing positive nationally. Private-sector facilities with high proportions testing positive are concentrated in KwaZulu-Natal (6), Mpumalanga (5) and Gauteng (4).

SOUTH AFRICA | WEEK 4 2021

Table 5.2 Private sector healthcare facilities with a high proportion testing positive, 24-30 January 2021

Facility Name	Province	Tests	PTP (95% CI)
Facility 1	KwaZulu-Natal	43	0.395 (0.249;0.541)
Facility 2	Western Cape	107	0.374 (0.282;0.466)
Facility 3	Gauteng	31	0.355 (0.186;0.523)
Facility 4	Gauteng	110	0.345 (0.257;0.434)
Facility 5	KwaZulu-Natal	27	0.333 (0.156;0.511)
Facility 6	Limpopo	284	0.331 (0.276;0.386)
Facility 7	Free State	65	0.323 (0.209;0.437)
Facility 8	Free State	112	0.321 (0.235;0.408)
Facility 9	Limpopo	73	0.315 (0.209;0.422)
Facility 10	Mpumalanga	64	0.313 (0.199;0.426)
Facility 11	Eastern Cape	29	0.310 (0.142;0.479)
Facility 12	North West	26	0.308 (0.130;0.485)
Facility 13	Limpopo	33	0.303 (0.146;0.460)
Facility 14	Northern Cape	83	0.301 (0.203;0.400)
Facility 15	Gauteng	37	0.297 (0.150;0.445)
Facility 16	KwaZulu-Natal	87	0.287 (0.192;0.382)
Facility 17	KwaZulu-Natal	60	0.283 (0.169;0.397)
Facility 18	Mpumalanga	469	0.281 (0.241;0.322)
Facility 19	KwaZulu-Natal	175	0.280 (0.213;0.347)
Facility 20	Mpumalanga	61	0.279 (0.166;0.391)
Facility 21	Mpumalanga	313	0.272 (0.222;0.321)
Facility 22	KwaZulu-Natal	82	0.268 (0.172;0.364)
Facility 23	Mpumalanga	159	0.264 (0.196;0.333)
Facility 24	Western Cape	38	0.263 (0.123;0.403)
Facility 25	Gauteng	73	0.260 (0.160;0.361)

95% CI: 95% confidence interval; PTP: positive test proportion

Health district-level results

The data from geo-locatable public testing (almost every public sector facility in the country) and private testing (approximately 83% of private testing facilities) in the week from 24-30 January 2021 have been located within the spatial framework of the health districts and health sub-districts (in the metros). The results, for the 25 municipalities and metropolitan health sub-districts showing the greatest proportions testing positive (PTP) are shown in Table 6. Districts showing the greatest proportions testing positive are concentrated in KwaZulu-Natal (7 districts), the Western Cape and Limpopo (5 districts each).

No districts showed an adjusted proportion testing positive greater than 50% this week, and only two exceeded 40%. The proportion of tests returned positive fell significantly in three districts (2 in KwaZulu-Natal, and one in Limpopo).


WEEK 4 2021 SOUTH AFRICA

Table 6. Health sub-districts with the highest proportion testing positive based on public and private sector data for the week of

Health district or sub-district	Province	PTP (95% CI)	Previous week
Ezingoleni	KwaZulu-Natal	0.465 (0.384-0.546)	0.545 (0.467-0.624)
Greater Letaba	Limpopo	0.446 (0.318-0.574)	0.454 (0.361-0.546)
Maquassi Hills	NorthWest	0.387 (0.311-0.463)	0.369 (0.313-0.426)
Cederberg	Western Cape	0.373 (0.265-0.481)	0.434 (0.345-0.523)
Ratlou	NorthWest	0.368 (0.232-0.505)	
Umzumbe	KwaZulu-Natal	0.367 (0.312-0.423)	0.514 (0.465-0.563)
Mtubatuba	KwaZulu-Natal	0.364 (0.296-0.432)	0.465 (0.418-0.511)
Matzikama	Western Cape	0.359 (0.292-0.425)	0.351 (0.300-0.402)
Dannhauser	KwaZulu-Natal		0.517 (0.427-0.606)
Ndwedwe	KwaZulu-Natal	0.344 (0.300-0.387)	0.388 (0.349-0.426)
Makhuduthamaga	Limpopo		0.507 (0.439-0.576)
Richtersveld	Northern Cape	0.322 (0.222-0.423)	0.291 (0.218-0.364)
Albert Luthuli	Mpumalanga	0.319 (0.266-0.372)	0.274 (0.232-0.316)
Makhado	Limpopo	0.319 (0.281-0.357)	0.372 (0.342-0.402)
Mafube	Free State	0.312 (0.228-0.396)	0.367 (0.277-0.458)
Pixley Ka Seme	Mpumalanga	0.299 (0.249-0.350)	0.345 (0.304-0.386)
Blouberg	Limpopo	0.298 (0.185-0.412)	0.359 (0.273-0.445)
Mandeni	KwaZulu-Natal	0.290 (0.247-0.333)	0.339 (0.306-0.372)
Langeberg	Western Cape	0.288 (0.202-0.374)	0.298 (0.234-0.361)
Bergrivier	Western Cape	0.288 (0.230-0.345)	0.276 (0.230-0.322)
Nketoana	Free State	0.286 (0.231-0.340)	0.322 (0.270-0.373)
Ubuhlebezwe	KwaZulu-Natal	0.285 (0.231-0.339)	0.340 (0.294-0.386)
Kannaland	Western Cape	0.283 (0.139-0.427)	0.350 (0.228-0.473)
Thaba Chweu	Mpumalanga	0.283 (0.238-0.327)	0.339 (0.305-0.373)
Modimolle	Limpopo	0.280 (0.144-0.415)	0.255 (0.192-0.318)

testing positive that are higher than, and CIs that do not overlap with, the previous week proportions and CIs. Elements have current week proportions testing positive that are previous week proportions and Cls.


The data for every district with a non-zero proportion testing positive or where the range of confidence interval is not more than 30% (15% either side of the point estimate) for the current week is presented graphically below.



HEALTH SUB-DISTRICT

Figure 9.1 Proportions testing positive by health sub-district in the Western Cape, Eastern Cape, Gauteng and KwaZulu-Natal provinces based on public and private sector data for the week of 24-30 January 2021.

SOUTH AFRICA | WEEK 4 2021



HEALTH SUB-DISTRICT

Figure 9.2 Proportions testing positive by health sub-district in the North West, Free State, Limpopo and Mpumalanga provinces based on public and private sector data for the week of 24-30 January 2021.

SOUTH AFRICA WEEK 4 2021

PROPORTION TESTING POSITIVE



HEALTH SUB-DISTRICT

Figure 9.3 Proportions testing positive by health sub-districts in the Northern Cape Province based on public and private sector data for the week of 24-30 January 2021.

The spatial pattern of adjusted proportions testing positive, including both public and private sector data, by health district and sub-district are shown for South Africa (Figure 10), Western Cape (Figure 11), Eastern Cape (Figure 12), Northern Cape (Figure 13), Free State (Figure 14), KwaZulu-Natal (Figure 15), North West (Figure 16), Gauteng (Figure 17), Mpumalanga (Figure 18) and Limpopo (Figure 19).




Figure 10. Proportion testing positive by health sub-district in South Africa for the week of 24-30 January 2021. Areas shaded white represent districts in which either (i) no tests were conducted, (ii) all tests were negative, or (iii) the confidence interval exceeded 30%.




Figure 11. Proportion testing positive by health sub-district in the Western Cape province for the week of 24-30 January 2021. Areas shaded white represent districts in which either (i) no tests were conducted, (ii) all tests were negative, or (iii) the confidence interval exceeded 30%




Figure 12. Proportion testing positive by health sub-district in the Eastern Cape province for the week of 24-30 January 2021. Areas shaded white represent districts in which either (i) no tests were conducted, (ii) all tests were negative, or (iii) the confidence interval exceeded 30%.




Figure 13. Proportion testing positive by health sub-district in Northern Cape Province for the week of 24-30 January 2021. Areas shaded white represent districts in which either (i) no tests were conducted, (ii) all tests were negative, or (iii) the confidence interval exceeded 30%.




Figure 14. Proportion testing positive by health sub-district in Free State Province for the week of 24-30 January 2021. Areas shaded white represent districts in which either (i) no tests were conducted, (ii) all tests were negative, or (iii) the confidence interval exceeded 30%.




Figure 15. Proportion testing positive by health sub-district in KwaZulu-Natal Province for the week of 24-30 January 2021. Areas shaded white represent districts in which either (i) no tests were conducted, (ii) all tests were negative, or (iii) the confidence interval exceeded 30%.




Figure 16. Proportion testing positive by health sub-district in North West Province for the week of 24-30 January 2021. Areas shaded white represent districts in which either (i) no tests were conducted, (ii) all tests were negative, or (iii) the confidence interval exceeded 30%.




Figure 17. Proportion testing positive by health sub-district in Gauteng Province for the week of 24-30 January 2021. Areas shaded white represent districts in which either (i) no tests were conducted, (ii) all tests were negative, or (iii) the confidence interval exceeded 30%.




Figure 18. Proportion testing positive by health sub-district in Mpumalanga Province for the week of 24-30 January 2021. Areas shaded white represent districts in which either (i) no tests were conducted, (ii) all tests were negative, or (iii) the confidence interval exceeded 30%.




Figure 19. Proportion testing positive by health sub-district in Limpopo Province for the week of 24-30 January 2021. Areas shaded white represent districts in which either (i) no tests were conducted, (ii) all tests were negative, or (iii) the confidence interval exceeded 30%.

SOUTH AFRICA WEEK 4 2021

Testing by patient admission status

In week 4 of 2021, 30.2% of tests were performed for hospitalised patients; 39.0% in the public sector and 23.6% in the private sector (Figure 20). The percentage testing positive decreased among both

inpatients and outpatients in the past week, and in week 4 was similar among outpatients (14.6%) and inpatients (15.3%) (Figure 21). In week 4 of 2021 the mean laboratory turnaround time in the public sector continued to be lower for inpatients (3.4 days) compared to outpatients (3.7 days) (Figure 22).



Figure 20. Percentage of inpatient tests performed by health sector, 1 March 2020 - 30 January 2021



Figure 21. Percentage testing positive by patient admission status, 6 December 2020 – 30 January 2021

SOUTH AFRICA | WEEK 4 2021




Figure 22. Mean number of days between date of specimen collection and date of test result in the public sector by patient admission status, 3 – 30 January 2021

Testing by age and sex

The mean age of individuals tested in week 4 of 2021 was 40.1 years, and was similar among females (40.2 years) and males (40.1 years). As in the previous few weeks, the majority of tests were performed in individuals in the 25-44 years' age groups (Figure 23).

In week 4, the testing rate in males was 355 per 100,000 persons and in females was 399 per 100,000 persons (Figure 24). The highest testing rates were observed in individuals ≥80 years of age (761 per 100,000 persons) in week 4. The percentage testing positive increased with increasing age, with a peak in the 10-14 year age group, and was higher in females compared to males (Figure 24).




Figure 23. Proportion of tests by age group and sex, South Africa, 24 - 30 January 2021

SOUTH AFRICA WEEK 4 2021




Figure 24. Testing rates per 100,000 persons and percentage testing positive by age group and sex, South Africa, week 4, 24 - 30 January 2021

Testing by test type

Testing for SARS-CoV-2 using rapid antigen tests was initiated towards the end of October 2020. Up to the end of week 4 of 2021, 306,455 antigen tests had been performed (38,430 in week 4). The percentage of antigen tests of all tests conducted has increased over recent weeks from 0.1% in week 44 of 2020 to 16.6% in week 4 of 2021. The majority of antigen tests have been performed in KwaZulu-Natal (n=143,907, 47.0%) and Eastern Cape (n=53,601, 17.5%) provinces

(Figure 25). Over recent weeks the percentage testing positive was higher for PCR tests compared to antigen tests, likely due to the lower sensitivity of antigen tests and the setting in which they were used (Figure 26). Not all antigen tests are included in this report, efforts are ongoing to improve completeness.




Figure 25. Number of antigen tests by province, and overall percentage antigen tests, South Africa, 1 November 2020 – 30 January 2021. WC, Western Cape; EC, Eastern Cape; FS, Free State; KZN, KwaZulu-Natal; GT, Gauteng; NC, Northern Cape; NW, North West; MP, Mpumalanga; LP, Limpopo




Figure 26. Percentage of laboratory tests positive for SARS-CoV-2 by test type and date of specimen collection, South Africa, 1 November 2020 – 30 January 2021

SOUTH AFRICA | WEEK 4 2021

Limitations

- A backlog in testing of samples by laboratories affects the reported numbers of tests performed. As a result, numbers tested during this period may change in subsequent reports.
- If higher-priority specimens were tested preferentially, this would likely result in an inflated proportion testing positive.
- Different and changing testing strategies (targeted vs. mass testing and PCR vs. antigenbased tests) used by different provinces makes percentage testing positive difficult to interpret and compare.
- Health district and sub-district level were mapped based on the testing facility and not place of residence.
- Patient admission status was categorised based on the reported patient facility and may not reflect whether the patient was actually admitted to hospital.
- Antigen tests may be underestimated as they are used in a number of different settings and results may not be reported.

CONCLUSIONS

Weekly testing volumes were highest in the first wave of infections in week 28 of 2020 (n=307,900), and in the second wave of infections in week 1 of 2021 (n=492,346). The number of tests performed in week 4 of 2021 (n=230,376) was lower than the previous few weeks. Gauteng (32.0%) performed the largest number of tests in week 4 of 2021, followed by KwaZulu-Natal (25.7%) and Western Cape (13.7%) provinces. KwaZulu-Natal (513 per 100,000 persons), Gauteng (476 per 100,000 persons), Northern Cape (467 per 100,000 persons) and Western Cape (452 per 100,000 persons) provinces had the highest testing rates in week 4 of 2021. The percentage of antigen tests of all tests performed has increased since antigen testing was implemented towards the end of October 2020, and accounted for 16.6% of all tests in week 4. The overall laboratory turnaround time for PCR tests in week 4 of 2021 was 2.0 days; 3.2 days in the public sector and <1 day in the private sector.

In the first wave of infections the percentage testing positive peaked at 29.7% in week 29 of 2020, and in the second wave of infections the percentage testing positive peaked at 34.9% in week 53 of 2020. In week 4 of 2021 the percentage testing positive was 14.0%, lower than has been observed in the previous few weeks. The percentage testing positive in week 4 of 2021 was highest in Limpopo (25.6%) and Mpumalanga (20.9%). Percentage testing positive was 10-20% in the Western Cape, Northern Cape, KwaZulu-Natal, North West, Free State and Gauteng, and was <10% in the Eastern Cape in week 4 of 2021. Compared to the previous week, in week 4 the percentage testing positive decreased in all provinces. Districts showing the greatest proportions testing positive in week 4 were concentrated in KwaZulu-Natal (7 districts), the Western Cape and Limpopo (5 districts each).