

Wastewater-based epidemiology for SARS-CoV-2 surveillance in South Africa

Report date: 15 October 2021

Sample collection dates up to 12 October 2021 (epidemiological week 41)

Centre for Vaccines and Immunology
National Institute for Communicable Diseases









Why test wastewater for SARS-CoV-2?



SARS-CoV-2 virus is excreted in stool by persons with active and recovering COVID-19 and can be found in wastewater

SARS-CoV-2 is not transmitted by faeco-oral route. Wastewater with SARS-CoV-2 is **not infectious**



Levels of SARS-CoV-2 in wastewater reflect population changes in case load and geographical distribution of cases



SARS-CoV-2 can be detected in wastewater before clinical cases appear

Increases in SARS-CoV-2 levels will appear before increases in clinical case load



Monitoring changes in SARS-CoV-2 levels can support public health preparedness and response activities









How is wastewater tested for SARS-CoV-2?





One litre of influent wastewater is collected in the morning (a 'grab' sample) and transported on ice to the lab



The one litre sample is concentrated



Viral RNA is extracted and PCR for SARS-CoV-2 is done





Results are visualised in a graph

SARS-CoV-2 is not transmitted by faeco-oral route. Wastewater with SARS-CoV-2 is not infectious











What does wastewater testing for SARS-CoV-2 mean?



How can we use wastewater testing for strengthening public health responses




- What should we do if levels go from negative to positive?
 - Inform public
 - Strengthen
 health
 promotion
 messages
 regarding NPIs
 and vaccination
 - Strengthen testing in affected areas



- What should we do if levels are increasing?
 - Inform public
 - Identify where NPIs are not being adhered to and target these areas for strengthened health promotion messages regarding NPIs and vaccination
 - Strengthen testing in affected areas
 - Prepare hospitals for overflow



- What should we do if levels are decreasing?
 - Inform and congratulate public
 - Reinforce public health messaging regarding NPIs
 - Strengthen
 vaccination to
 ensure population
 is protected for the
 next wave




- What should we do if levels go from positive to negative?
 - Inform public
 - Redirect resources from NPIs to vaccination to ensure population is protected for the next wave





Results: Gauteng 1



Changes in SARS-Cov-2 viral load (data points and coloured lines) in in-flowing untreated wastewater from plants in City of Tshwane, compared with <u>laboratory-confirmed cases resident in City of Tshwane</u> (grey bars), by epidemiological week, 2021










Results: Gauteng 2



Changes in SARS-Cov-2 viral load (data points and coloured lines) in in-flowing untreated wastewater from plants in City of Johannesburg, compared with <u>laboratory-confirmed cases resident in City of Johannesburg</u> (grey bars), by epidemiological week, 2021










Results: Gauteng 3



Changes in SARS-Cov-2 viral load (data points and coloured lines) in in-flowing untreated wastewater from plants in Ekurhuleni, compared with <u>laboratory-confirmed cases resident in Ekurhuleni</u> (grey bars), by epidemiological week, 2021










Results: KwaZulu-Natal 1



Changes in SARS-Cov-2 viral load (data points and coloured lines) in in-flowing untreated wastewater from plants in eThekwini, compared with <u>laboratory-confirmed cases resident in eThekwini</u> (grey bars), by epidemiological week, 2021










Results: KwaZulu-Natal 2



Changes in SARS-Cov-2 viral load (data points and coloured lines) in in-flowing untreated wastewater from plants in Mgungundlovu, compared with <u>laboratory-confirmed cases resident in Umgungundlovu</u> (grey bars), by epidemiological week, 2021










Results: Free State



Changes in SARS-Cov-2 viral load (data points and coloured lines) in in-flowing untreated wastewater from plants in Mangaung, compared with <u>laboratory-confirmed cases in Mangaung</u> (grey bars), by epidemiological week, 2021










Results: Eastern Cape 1



Changes in SARS-Cov-2 viral load (data points and coloured lines) in in-flowing untreated wastewater from plants in Nelson Mandela Metro, compared with <u>laboratory-confirmed cases resident in Nelson Mandela Metro</u> (grey bars), by epidemiological week, 2021










Results: Eastern Cape 2



Changes in SARS-Cov-2 viral load (data points and coloured lines) in in-flowing untreated wastewater from plants in Buffalo City Metro, compared with <u>laboratory-confirmed cases resident in Buffalo City</u> (grey bars), by epidemiological week, 2021










Results: Western Cape



Changes in SARS-Cov-2 viral load (data points and coloured lines) in in-flowing untreated wastewater from plants in Western Cape Province, compared with laboratory-confirmed cases resident in City of Cape Town (grey bars), by epidemiological week, 2021










Results: Northern Cape 1



Changes in SARS-Cov-2 viral load (data points and coloured lines) in in-flowing untreated wastewater from a plant in the Northern Cape Province, compared <u>with laboratory-confirmed cases resident in Namakwa</u> (grey bars), by epidemiological week, 2021










Results: Northern Cape 2



Changes in SARS-Cov-2 viral load (data points and coloured lines) in in-flowing untreated wastewater from a plant in the Northern Cape Province, compared with laboratory-confirmed cases resident in Frances Baard (grey bars), by epidemiological week, 2021










Results: Limpopo



Changes in SARS-Cov-2 viral load (data points and coloured lines) in in-flowing untreated wastewater from a plant in the Limpopo Province, compared <u>with laboratory-confirmed cases resident in Polokwane</u> (grey bars), by epidemiological week, 2021










Results: Mpumalanga 1



Changes in SARS-Cov-2 viral load (data points and coloured lines) in in-flowing untreated wastewater from a plant in the Mpumalanga Province, compared <u>with laboratory-confirmed cases resident in Mbombela</u> (grey bars), by epidemiological week, 2021










Results: Mpumalanga 2



Changes in SARS-Cov-2 viral load (data points and coloured lines) in in-flowing untreated wastewater from a plant in the Mpumalanga Province, compared <u>with laboratory-confirmed cases resident in Emalahleni</u> (grey bars), by epidemiological week, 2021










Results: North West 1



Changes in SARS-Cov-2 viral load (data points and coloured lines) in in-flowing untreated wastewater from a plant in the North West Province, compared <u>with laboratory-confirmed cases resident in Matlosana</u> (grey bars), by epidemiological week, 2021










Results: North West 2



Changes in SARS-Cov-2 viral load (data points and coloured lines) in in-flowing untreated wastewater from a plant in the North West Province, compared <u>with laboratory-confirmed cases resident in JB Marks</u> (grey bars), by epidemiological week, 2021










Results: North West 3



Changes in SARS-Cov-2 viral load (data points and coloured lines) in in-flowing untreated wastewater from a plant in the North West Province, compared <u>with laboratory-confirmed cases resident in Bojanala</u> (grey bars), by epidemiological week, 2021







