


SOUTH AFRICA WEEK 45 2021

CONTENTS

Surveillance programme	2
description and comments	
Data from virologic surveillance to	4
aid interpretation	
•Influenza detections per week	
• Respiratory syncytial virus	
(RSV) detections per week	
• SARS-CoV-2 detections per	
week	
Number of consultations per week	5
Private consultation thresholds by	6
age group: All Ages	
<5 years	7
5-19 years	8
20-49 years	9
≥50 years	10
Private consultation thresholds by	10
province: Eastern Cape	

HIGHLIGHTS: WEEK 45

- Total number of respiratory hospitalisations have generally been decreasing in recent weeks to below/low threshold in most provinces and age groups with slight increases in the 0-4-year age group in Eastern Cape and KwaZulu Natal provinces.
- The percentage of respiratory general practitioner visits have been decreasing in recent weeks reaching below/low threshold in all age groups and provinces.
- The percentage of respiratory emergency department visits have been decreasing in recent weeks reaching below/low threshold in most age groups and provinces with slight increases in the 0-4-year age group and in Eastern Cape province.



Gauteng

KwaZulu-Natal

Western Cape

12

13

14

WEEK **45** 2021

PROGRAMME DESCRIPTIONS

Inpatient data from a large national private hospital group and outpatient data from a general practitioner network linked to the same hospital group were received for the last week. Data were obtained from eight provinces (Eastern Cape, Free State, Gauteng, Limpopo, KwaZulu-Natal, Mpumalanga, North West, Western Cape). Sufficient numbers for province-level reporting were available for four of these (bold). Consultations and admissions were coded based on discharge diagnosis using the International Classification of Diseases and Related Health Problems, 10th revision (ICD-10). Data were analysed using the indicator: All respiratory and confirmed or suspected COVID-19 (J00-J99 & U07.1 & U07.2)/Total consultations. Data on the indicator Pneumonia and Influenza (J10-J18)/Total consultations are available on request but were not included in this report.

Data were categorised in the following age groups: All ages, <5 years, 5-19 years, 20-49 years, ≥50 years

Epidemic Threshold

Thresholds were calculated using the Moving Epidemic Method (MEM), a sequential analysis using the R Language, available from: http://CRAN.R-project.org/web/package=mem, designed to calculate the duration, start and end of the annual influenza epidemic. MEM uses the 40th, 90th and 97.5th percentiles established from historical data (2015-2019 for inpatients, 2016-2019 for outpatients) to calculate thresholds of activity, defined as follows:

- · Epidemic threshold: Median of weekly values for all baseline years
- · Low activity: Between epidemic threshold including 40th percentile
- · Moderate activity: Between 40th and 90th percentile
- · High activity: Between 90th and 97.5th percentile
- · Very high activity: 97.5th percentile and above

Hospitalization data for recent weeks are adjusted for delayed reporting (diagnosis codes assigned on discharge delayed for prolonged hospitalisations). Adjustment accounts for the probability of being admitted, but not yet discharged at the time of data drawdown using the age- and syndrome-specific probability distribution of duration of admission obtained from all hospitalizations that occurred during 2015-2019 and applied to weeks reported.

WEEK **45** 2021

INTERPRETATION OF DATA PRESENTED IN THIS REPORT

Total respiratory admissions peaked in the first COVID-19 wave in week 29 of 2020 reaching approximately double the level before the lockdown after which numbers decreased. The second wave numbers peaked in week 1 of 2021 while the third wave numbers peaked in week 20 of 2021. Numbers have been decreasing since then.

The proportion of admissions coded as confirmed COVID-19 (among the suspected cases) in the first COVID-19 wave peaked in week 31 of 2020 at 79%, in the second wave in week 1 of 2021 at 95%, and in the third wave in week 27 of 2021 at 97%. It has since decreased to 17%.

Total COVID-19 and respiratory outpatient (general practitioner) consultations peaked in week 28 of 2020 for the first COVID-19 wave, in week 1 of 2021 for the second COVID-19 wave and week 25 of 2021 in the third COVID-19 wave and have since decreased. The proportion of general practitioner consultations coded as confirmed COVID-19 (among the suspected cases) peaked in week 33 of 2020 at 34% for the first wave, in week 2 of 2021 at 34% for the second wave and again in week 28 at 50.1% for the third wave. Currently, the proportion is at 10%.

Total respiratory emergency department consultations coded as confirmed COVID-19 (among the suspected cases) peaked in week 30 of 2020 at 84% for the first wave, in week 2 of 2021 at 96% for the second wave and in week 25 at 98.4% for the third wave. The current proportion has decreased to 50%.

Proportion of respiratory or suspected COVID-19 admissions overall peaked in the very high level in weeks 29 of 2020, 53 of 2020 and 28 of 2021 for the first, second and third waves respectively. Currently, admissions are decreasing to low threshold. By age group, percent respiratory or suspected COVID admissions for 0-4 years, remained below threshold throughout the three waves, but have been increasing over the past weeks and is currently low threshold. Among individuals aged 5-19 years, percent admissions mainly remained below threshold with a brief increase to low/moderate activity during weeks 27 to 32 2020. Among individuals aged 20-

49 years and ≥50 years, percent respiratory admissions peaked in the very high level around week 29 of 2020, week 53 of 2020 and week 27 of 2021. Currently, the admissions for children aged <5 years slightly increased but are in low threshold and for the remaining age groups the admissions have decreased to below/low threshold.

Proportion of outpatient (general practitioner) respiratory or suspected COVID-19 consultations peaked in week 28 of 2020, week 53 of 2020 and week 26 of 2021, after which it has recently declined to low threshold. In recent weeks, the overall level declined to below/low threshold in all age groups.

Proportion of emergency department respiratory or suspected COVID-19 consultations peaked in week 28 of 2020, week 53 of 2020 and week 26 of 2021, and recently decreasing to low threshold. By age group, percent emergency department visits showed similar trends. Currently, emergency visits are below/low threshold in all age groups but have marginally increased in the 0-4-year age group.

Trends in proportion of admissions, outpatient and emergency department respiratory or COVID consultations varied by province. The first wave peaked between weeks 27 and 30 of 2020 in the four provinces evaluated, the second wave in weeks 49 for Eastern Cape and 53 of 2020 for the remaining three provinces and the third wave in weeks 26 for Gauteng and 28 of 2021 for the remaining three provinces. The second wave peak exceeded the first wave peak in Gauteng. Kwazulu-Natal, and the Western Cape, and the third wave peak exceeded the second wave peak in Gauteng only. The proportion of respiratory inpatients slightly increased in Eastern Cape and KwaZulu Natal province but remaining below/low thresholds. The proportion of outpatients/general practitioner consultations remained low and below/low thresholds in all provinces. The proportion of

emergency department consultations slightly increased in Eastern Cape province but also remain currently below/low thresholds.


WEEK **45** 2021


Limitations

Thresholds are established based on the proportion of respiratory consultations. If the numbers of non-respiratory consultations drop substantially because of changes in health-seeking behaviour, this could lead to elevated respiratory proportions. Delays in the coding of consultations may lead to changes in data from previous weeks. Differences by province and age group should be interpreted with caution due to low numbers in some groups.

WEEK **45** 2021


DATA FROM VIROLOGIC SURVEILLANCE PROGRAMMES TO AID IN INTERPRETATION OF CONSULTATION TRENDS





WEEK **45** 2021


Number of consultations - all respiratory including confirmed or suspected COVID-19 and other consultations by week



WEEK 45 2021


ALL AGES

All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators - Inpatients




All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Outpatients

General Practitioners



Emergency Department




www.nicd.ac.za

TOLL-FREE NUMBER 0800 029 999

WEEK 45 2021


0-4 YEARS OF AGE

All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators - Inpatients



All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Outpatients

General Practitioners



Emergency Department



www.nicd.ac.za

TOLL-FREE NUMBER **0800 029 999**


WEEK 45 2021


5-19 YEARS OF AGE

All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators - Inpatients



All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators - Outpatients






WEEK 45 2021


20-49 YEARS OF AGE


All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators - Inpatients



All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Outpatients

General Practitioners






www.nicd.ac.za

TOLL-FREE NUMBER 0800 029 999

WEEK **45** 2021


≥50 YEARS OF AGE

All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators - Inpatients




All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Outpatients

General Practitioners



Emergency Department




www.nicd.ac.za

TOLL-FREE NUMBER 0800 029 999

WEEK 45 2021


EASTERN CAPE PROVINCE

All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Inpatients




All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Outpatients

General Practitioners



Emergency Department




www.nicd.ac.za

TOLL-FREE NUMBER 0800 029 999

WEEK 45 2021


GAUTENG

All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators - Inpatients




All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Outpatients

General Practitioners



Emergency Department




Mean % 2015- 2019 (Inpatients/Casualty) 2016-2019 (General Practitioner)

WEEK **45** 2021


KWAZULU-NATAL

All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators - Inpatients




All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Outpatients

General Practitioners








www.nicd.ac.za

TOLL-FREE NUMBER 0800 029 999

WEEK 45 2021


WESTERN CAPE PROVINCE

All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators - Inpatients




All Respiratory including confirmed or suspected COVID-19 (J00-J99 & U07) indicators – Outpatients

General Practitioners



Emergency Department



www.nicd.ac.za

TOLL-FREE NUMBER 0800 029 999

WEEK **45** 2021

ACKNOWLEDGEMENT

We would like to acknowledge the contribution of the following individuals:

Dr Anchen Laubscher, Group Medical Director Netcare

Dr Caroline Maslo, Head of Infection Control Netcare

Mande Toubkin, General Manager Emergency, Trauma, Transplant CSI and Disaster Management Netcare

