

COVID-19 WEEKLY EPIDEMIOLOGY BRIEF




NATIONAL INSTITUTE FOR
COMMUNICABLE DISEASES

Division of the National Health Laboratory Service

SOUTH AFRICA WEEK 16 2022

CUMULATIVE DATA FROM



CASES

3 762 911
IN TOTAL

20 813
THIS WEEK**




PERSONS

6 256,6
INCIDENCE RISK*

38
MEDIAN AGE

PROVINCES AT A GLANCE



* Incidence risk - cases per 100 000 persons
** based on samples collected/received in current reporting week

SUMMARY

Overview of report

Disease surveillance is a core function of the National Institute for Communicable Diseases (NICD), a division of the National Health Laboratory Service (NHLS). This report summarises data from a national laboratory-based surveillance system that is used to monitor the coronavirus disease 2019 (COVID-19) pandemic, caused by the SARS-CoV-2 virus, in South Africa. This report is based on data collected up to 23 April 2022 (week 16 of 2022). Note: Trends in numbers of new cases by province and age group may be affected by changes in testing practice and delays in testing of specimens and numbers are updated weekly as new data become available. The methods and data sources can be found at the end of the report.

Highlights

- As of 23 April, a total of 3 762 911 laboratory-confirmed COVID-19 cases had been detected in South Africa. Of these, 21 681 were cases reported since the last report (week 15 of 2022). There was a 123.3% increase in the number of new cases detected in week 16 of 2022 (20 813) compared to the number of new cases detected in week 15 of 2022 (9 322).
- In the past week, the Gauteng Province reported the highest number of cases detected (10 433/20 813, 50.1%), followed by the KwaZulu-Natal Province (4 741/20 813, 22.8%), and Western Cape Province (2 970/20 813, 14.3%), with other provinces reporting $\leq 5\%$ each.
- In the past week, Gauteng Province reported the highest weekly incidence risk (66.0 cases per 100 000 persons), followed by the Western Cape Province (41.7 cases per 100 000 persons), and the KwaZulu-Natal Province (41.2 cases per 100 000 persons). The other provinces reported weekly incidence below 20 cases per 100 000 persons.
- In the past week, all provinces reported an increase in weekly incidence risk, compared to the previous week. The increase in weekly incidence risk ranged from 1.8 cases per 100 000 persons (141.9% increase) in the Limpopo Province to 36.6 cases per 100 000 persons (124.8% increase) in the Gauteng Province.
- All the districts in Gauteng Province reported an increase in weekly incidence risk in the past week, the increase ranged from 17.5 cases per 100 000 persons (117.3% increase) in the West Rand District to 33.5 cases per 100 000 persons (113.0% increase) in the City of Johannesburg Metro.
- The highest weekly incidence risk among cases detected in week 16 of 2022 was reported in the ≥ 80 -year age group (64.8 cases per 100 000 persons), and the lowest weekly incidence risk was in the 0-4-year age group (9.4 cases per 100 000 persons).

INCIDENCE
RISK FOR
CURRENT WEEK

34,6

CASES PER
100 000
PERSONS

50,1%

OF CASES
REPORTED IN
GAUTENG IN
CURRENT WEEK

IN CURRENT
WEEK, THE
HIGHEST
WEEKLY
INCIDENCE
RISK WAS IN
CASES AGED
80+ YEARS (64,8
CASES PER 100
000 PERSONS)

COVID-19 WEEKLY EPIDEMIOLOGY BRIEF

WEEK 16 2022




Figure 1. Number and cumulative number of laboratory-confirmed cases of COVID-19 by date of specimen collection, South Africa, 3 March 2020 – 23 April 2022 (n = 3 762 911)

Table 1. Number and cumulative/weekly incidence risk of laboratory-confirmed cases of COVID-19 and testing per 100 000 persons by province, South Africa, 3 March 2020 – 23 April (n = 3 762 911)

Province	Cumulative cases (n) (percentage, n/ total cases in South Africa)	New cases ¹ detected in week 16 of 2022 (17-23 Apr), n (percentage ² , n/total)	Population in mid-2021 ³ , n	Cumulative incidence risk (cases per 100 000 persons)	Incidence risk of new cases detected in week 16 of 2022 (cases/100 000 persons)	Tests ⁴ per 100 000 persons, , 17-23 Apr 2022
Eastern Cape	347 190 (9.2)	864 (4.2)	6 676 590	5 200.1	12.9	105.8
Free State	203 514 (5.4)	573 (2.8)	2 932 441	6 940.1	19.5	218.4
Gauteng	1 230 111 (32.7)	10 433 (50.1)	15 810 388	7 780.4	66.0	347.3
KwaZulu-Natal	669 861 (17.8)	4 741 (22.8)	11 513 575	5 818.0	41.2	222.1
Limpopo	156 271 (4.2)	179 (0.9)	5 926 724	2 636.7	3.0	31.1
Mpumalanga	194 328 (5.2)	464 (2.2)	4 743 584	4 096.6	9.8	105.9
North West	193 902 (5.2)	398 (1.9)	4 122 854	4 703.1	9.7	112.3
Northern Cape	109 293 (2.9)	191 (0.9)	1 303 047	8 387.5	14.7	170.5
Western Cape	658 441 (17.5)	2 970 (14.3)	7 113 776	9 255.9	41.7	252.4
Unknown						
Total	3 762 911	20 813	60 142 978	6 256.6	34.6	210.5

¹New cases refer to cases whose samples were collected or received in the current reporting week; ²Percentage=n/total number of new cases (specimen collected or received in current reporting week); ³2021 Mid-year population Statistics South Africa ⁴Data on number of tests conducted sourced from COVID-19 weekly testing report of the same reporting week

COVID-19 WEEKLY EPIDEMIOLOGY BRIEF

WEEK 16 2022




Figure 2. Cumulative incidence risk of laboratory-confirmed cases of COVID-19 by province and epidemiologic week South Africa 3 March 2020 – 23 April 2022 (n = 3 762 911)




Figure 3. Weekly incidence risk of laboratory-confirmed cases of COVID-19 by province and epidemiologic week South Africa 3 March 2020 – 23 April 2022 (n = 3 762 911)

COVID-19 WEEKLY EPIDEMIOLOGY BRIEF

WEEK 16 2022

Characteristics of COVID-19 cases in South Africa by age and sex




Figure 4. Weekly incidence risk of laboratory-confirmed cases of COVID-19 by age group in years and epidemiologic week South Africa 3 March 2020 – 23 April 2022 (n = 3 727 671, 35 240 missing age)




Figure 5. Weekly incidence risk of laboratory-confirmed cases of COVID-19 by sex and epidemiologic week South Africa 3 March 2020 – 23 April 2022 (n = 3 723 251, sex missing for 39 660)

COVID-19 WEEKLY EPIDEMIOLOGY BRIEF

WEEK 16 2022

Table 2. Number of laboratory-confirmed cases of COVID-19 and cumulative/weekly incidence risk by age group South Africa 3 March 2020 – 23 April 2022 n = 3 727 671, 35 240 missing age)

Age group (years)	Cumulative cases (n) (percentage, n/total cases in South Africa)	New cases ¹ detected in week 16 of 2022 (17-23 Apr) n (percentage ² , n/total)	Population in mid-2021 ³ , n	Cumulative incidence risk (cases per 100 000 persons)	Incidence risk of new cases detected in week 16 of 2022 (cases/100 000 persons)
0-4	58 701 (1.6)	539 (2.6)	5 708 956	1 028.2	9.4
5-9	80 299 (2.2)	770 (3.7)	5 663 296	1 417.9	13.6
10-14	144 584 (3.9)	991 (4.8)	5 671 023	2 549.5	17.5
15-19	208 955 (5.6)	1 038 (5.0)	4 909 941	4 255.8	21.1
20-24	247 766 (6.6)	1 248 (6.0)	4 739 305	5 227.9	26.3
25-29	363 019 (9.7)	1 913 (9.2)	5 324 134	6 818.4	35.9
30-34	418 516 (11.2)	2 225 (10.7)	5 630 643	7 432.8	39.5
35-39	424 607 (11.4)	2 364 (11.4)	4 985 251	8 517.3	47.4
40-44	359 810 (9.7)	1 836 (8.9)	3 881 731	9 269.3	47.3
45-49	342 547 (9.2)	1 838 (8.9)	3 254 138	10 526.5	56.5
50-54	307 384 (8.2)	1 672 (8.1)	2 625 390	11 708.1	63.7
55-59	255 127 (6.8)	1 309 (6.3)	2 243 823	11 370.2	58.3
60-64	178 585 (4.8)	986 (4.8)	1 815 810	9 835.0	54.3
65-69	122 878 (3.3)	661 (3.2)	1 422 604	8 637.5	46.5
70-74	89 601 (2.4)	575 (2.8)	1 024 345	8 747.2	56.1
75-79	58 029 (1.6)	348 (1.7)	647 265	8 965.3	53.8
≥80	67 263 (1.8)	386 (1.9)	595 323	11 298.6	64.8
Unknown	35 240 (0.0)	114 (0.0)			
Total	3 762 911 (100.0)	20 813 (100.0)	60 142 978	6 256.6	34.6

¹New cases refer to cases whose samples were collected or received in the current reporting week; ²Percentage=n/total number of new cases (specimen collected or received in current reporting week); ³2021 Mid-year population Statistics South Africa

COVID-19 WEEKLY EPIDEMIOLOGY BRIEF

WEEK 16 2022

Provincial trends of COVID-19 cases

Eastern Cape Province




Figure 6. Weekly incidence risk of laboratory-confirmed cases of COVID-19 by district and epidemiologic week Eastern Cape Province 3 March 2020 – 23 April 2022 (n = 292 303, 54 887 missing district)




Figure 7. Weekly incidence risk of laboratory-confirmed cases of COVID-19 by age group in years and epidemiologic week Eastern Cape Province 3 March 2020 – 23 April 2022 (n = 343 533, 3 657 missing age)

COVID-19 WEEKLY EPIDEMIOLOGY BRIEF

WEEK 16 2022

Western Cape Province




Figure 8. Weekly incidence risk of laboratory-confirmed cases of COVID-19 by district and epidemiologic week Western Cape Province 3 March 2020 – 23 April 2022 (n = 605 193, 53 248 missing district)




Figure 9. Weekly incidence risk of laboratory-confirmed cases of COVID-19 by age group in years and epidemiologic week Western Cape Province 3 March 2020 – 23 April 2022 (n = 656 670, 1 771 missing age)

COVID-19 WEEKLY EPIDEMIOLOGY BRIEF

WEEK 16 2022

Gauteng Province




Figure 10. Weekly incidence risk of laboratory-confirmed cases of COVID-19 by district and epidemiologic week Gauteng Province 3 March 2020 – 23 April 2022 (n = 875 421, 35 469 missing district)




Figure 11. Weekly incidence risk of laboratory-confirmed cases of COVID-19 by age group in years and epidemiologic week Gauteng Province 3 March 2020 – 23 April 2022 (n = 1 217 691, 12 420 missing age)

COVID-19 WEEKLY EPIDEMIOLOGY BRIEF

WEEK 16 2022

KwaZulu-Natal Province




Figure 12. Weekly incidence risk of laboratory-confirmed cases of COVID-19 by district and epidemiologic week KwaZulu-Natal Province 3 March 2020 – 23 April 2022 (n = 413 423, 256 438 missing district)




Figure 13. Weekly incidence risk of laboratory-confirmed cases of COVID-19 by age group in years and epidemiologic week KwaZulu-Natal Province 3 March 2020 – 23 April 2022 (n = 661 280, 8 581 missing age)

COVID-19 WEEKLY EPIDEMIOLOGY BRIEF

WEEK 16 2022

Free State Province




Figure 14. Weekly incidence risk of laboratory-confirmed cases of COVID-19 by district and epidemiologic week Free State Province 3 March 2020 – 23 April 2022 (n = 173 641, 29 873 missing district)




Figure 15. Weekly incidence risk of laboratory-confirmed cases of COVID-19 by age group in years and epidemiologic week Free State Province 3 March 2020 – 23 April 2022 (n = 202 695, 819 missing age)

COVID-19 WEEKLY EPIDEMIOLOGY BRIEF

WEEK 16 2022

Limpopo Province




Figure 16. Weekly incidence risk of laboratory-confirmed cases of COVID-19 by district and epidemiologic week Limpopo Province 3 March 2020 – 23 April 2022 (n = 116 202, 40 069 missing district)




Figure 17. Weekly incidence risk of laboratory-confirmed cases of COVID-19 by age group in years and epidemiologic week Limpopo Province 3 March 2020 – 23 April 2022 (n = 155 561, 710 missing age)

COVID-19 WEEKLY EPIDEMIOLOGY BRIEF

WEEK 16 2022

Mpumalanga Province




Figure 18. Weekly incidence risk of laboratory-confirmed cases of COVID-19 by district and epidemiologic week Mpumalanga Province 3 March 2020 – 23 April 2022 (n = 127 006, 67 322 missing district)




Figure 19. Weekly incidence risk of laboratory-confirmed cases of COVID-19 by age group and epidemiologic week Mpumalanga Province 3 March 2020 – 23 April 2022 (n = 190 253, 4 075 missing age)

COVID-19 WEEKLY EPIDEMIOLOGY BRIEF

WEEK 16 2022

North West Province




Figure 20. Weekly incidence risk of laboratory-confirmed cases of COVID-19 by district and epidemiologic week North West Province 3 March 2020 – 23 April 2022 (n = 120 495, 73 407 missing district)




Figure 21. Weekly incidence risk of laboratory-confirmed cases of COVID-19 by age group in years and epidemiologic week North West Province 3 March 2020 – 23 April 2022 (n = 191 414, 2 488 missing age)

COVID-19 WEEKLY EPIDEMIOLOGY BRIEF

WEEK 16 2022

Northern Cape Province




Figure 22. Weekly incidence risk of laboratory-confirmed cases of COVID-19 by district and epidemiologic week Northern Cape Province 3 March 2020 – 23 April 2022 (n = 83 688, 25 605 missing district)




Figure 23 Weekly incidence risk of laboratory-confirmed cases of COVID-19 by age group in years and epidemiologic week Northern Cape Province 3 March 2020 – 23 April 2022 (n = 108 574, 719 missing age)

Methods

Testing for SARS-CoV-2 began on 28 January 2020 at the NICD and after the first case was confirmed in early March 2020, testing was expanded to a larger network of private and NHLS laboratories. Respiratory specimens were submitted from persons under investigation (PUI). Initially, tested individuals were those who had travelled to countries with COVID-19 transmission but the PUI definition was changed over time. Community symptom screening and referral for PCR testing was implemented in April 2020 but the strategy was changed to a more targeted approach in May 2020. Community screening was largely discontinued and testing efforts then focussed on areas identified as hot spots and on investigating clusters. Contacts of cases were traced and tested if symptomatic. In some provinces and certain circumstances (e.g. closed settings, workplaces), asymptomatic contacts were tested. In recent weeks, testing has been prioritised for healthcare workers and hospitalised patients. Laboratories used any one of several in-house and commercial PCR assays to test for the presence of SARS-CoV-2 RNA. Testing for SARS-CoV-2 using rapid antigen-based tests was implemented during November 2020. We excluded specimens collected outside South Africa. Date of specimen receipt in the laboratory was used when date of specimen collection was missing. A case of COVID-19 was defined as any person, resident in South Africa, with a single positive SARS-CoV-2 PCR or antigen test. For reports published from week 2 of 2022 onwards, we used 2021 mid-year population estimates to calculate incidence risk (cumulative and weekly incidence). For historical reports published from week 41 of 2020 to week 1 of 2022, 2020 mid-year population estimates were used, and reports published from epidemiologic week 10 (during the start of COVID-19 epidemic in South Africa) to week 40 of 2020, 2019 mid-year population estimates were used. Data on number of tests conducted in the past week as reported in the simultaneously-published COVID-19 weekly testing report was used to calculate tests conducted per 100 000 persons. Data on province and district allocation was based on geocoding algorithm using in order of priority (i) completeness of patient data, (ii) submitting doctor's address, (iii) registering doctor's address and as final option, (iv) the guarantor's address data. The geocoding algorithm used the most complete

data for assigning data on province and district where adequate information was provided on the lab request form at the time of sample collection. Data on district allocation may lag resulting in number of cases in recent weeks missing district allocation. Prevalence and incidence risk by districts should be interpreted with caution.

Until the week 29 of 2020 report, new cases were defined as all cases reported since the last report, irrespective of when the sample was collected. Subsequent to the week 29 of 2020 report, new cases are now defined as cases detected in the past epidemiologic week based on date of sample collection or sample receipt. It is therefore possible for numbers reported as new cases for the current reporting week not to tally with total additional cases reported since the last report. This will be the case when there was a delay in reporting of cases.

Limitations

This report is based on laboratory-based surveillance of laboratory-confirmed cases. The number of reported cases is heavily dependent on testing practices. Although trends over time and comparisons by geographic area are presented in this report, changes in testing practices over time or differences by region may partially explain the results. Differences in health-seeking behaviour by age group and sex could also contribute to the observed differences in case numbers between groups. Delays in reporting may result in incomplete data for recent weeks, leading to an apparent reduction in number of cases. Changes in testing strategy during the different times of the epidemic may also affect the number of cases reported, leading to a decrease in number of positive cases if testing is only conducted for severe cases or certain risk groups.