


SARS-CoV-2 Sequencing Update 27 May 2022



Supported by the DSI and the SA MRC

Msomi N, Mlisana K, et al. Lancet Microbe 2020

The genomic data presented here are based on South African SARS-CoV-2 sequence data downloaded from GISAID (www.gisaid.org) on 27 May 2022 at 08h00














Data license: <https://www.gisaid.org/registration/terms-of-use/>

Elbe, S., and Buckland-Merrett, G. (2017) Data, disease and diplomacy: GISAID's innovative contribution to global health. *Global Challenges*, 1:33-46. DOI: 10.1002/gch2.1018 PMCID: 31565258

Shu, Y., McCauley, J. (2017) GISAID: Global initiative on sharing all influenza data – from vision to reality. *EuroSurveillance*, 22(13) DOI: 10.2807/1560-7917.ES.2017.22.13.30494 PMCID: PMC5388101

SARS-CoV-2 GENOMIC SURVEILLANCE

epiweeks 39 (2021) - 20 (2022)



966 genomes added since the previous report


Bar graphs represent genomes sequenced per epiweek, with lines representing cases by collection date (weeks 39 [2021] – 20 [2022])

Genomes and cases presented as provincial total (percentage of national total) for epiweeks 39 (2021) – 20 (2022)

PTP: percentage testing positive in week 20 (15 May 2022 – 21 May 2022); arrow indicates direction of change since previous week (8 May 2022 – 14 May 2022) if change was significant ($P < 0.05$)

Number of South African genomes deposited on GISAID, by specimen collection week, 2020 – 2022


(N=40 993*)




*This represents the cleaned, de-duplicated dataset of unique **National and Pneumonia Surveillance** sequences. This dataset will be used for all further figures.

GISAID genomes vs total cases, 2020 – 2022 (N=40 993)

Provincial breakdown of genomes deposited into GISAID




Proportion of total cases versus proportion of total genomes



All provinces, apart from GP, KZN and WC, have comparable percentages of overall cases and sequenced genomes. All provinces have contributed sequences for April and May.


Number and percentage of clades by epiweek in South Africa, 2021 – 2022 (34 252*)



Delta dominated in South Africa until October at >80%. Omicron has dominated from November onwards.


Prevalence of Variants of Concern (VOC) and Variants of Interest (VOI) in March – May 2022

March (N=1462)




Total Omicron in Mar: 1438 (98.4%)

April (N=2269)



Total Omicron in Apr: 2262 (99.7%)

May (N=742)




Total Omicron in May: 739 (99.6%)

Omicron (21K/BA.1.*)	Omicron (21M/BA.3)	Omicron (22B/BA.5)	Delta (21I)	C.1.2 (20D)	Beta (20H, V2)	Kappa (21B)	Eta (21D)	Other	unassigned
Omicron (21L/BA.2.*)	Omicron (22A/BA.4)	Delta (21A)	Delta (21J)	Alpha (20I, V1)					

Omicron dominated in March (98.4%, 1438/1462), April (99.7%, 2262/2269) and May (99.6%, 739/742). BA.2 was dominant in March. BA.4 and BA.5 together were dominant in April and are dominant in May.

Detection Rates: Beta, Delta, C.1.2 and Omicron



Detection rates of variants being monitored in South Africa



*Bars represent percentage prevalence of variant for the month; total number sequences collected for the month are given below the bar

Omicron has been dominant since November (>85% in November, >98% in December – May).
BA.2 made up 43% of genomes in January, 86% in February, 78% in March and 26% in April. BA.4 and BA.5 together dominated in April at 72%, and in May at 92%.


South Africa, 2021-2022, n = 34 252*



Clade key (bar graph)




Weekly percentage testing positive key (line graph)




*Excludes sequences missing collection dates, as well as those collected January 1st and 2nd 2021 as they are part of epiweek 53 of 2020.

Eastern Cape Province, 2021-2022, n = 2743

EC number of genomes, cases, and PTP per epiweek (N=2743)




EC percentage of genomes, number of cases, and PTP per epiweek (N=2743)



Clade key (bar graph)




Weekly percentage testing positive key (line graph)




Free State Province, 2021-2022, n = 1484

FS number of genomes, cases, and PTP per epiweek (N=1484)




FS percentage of genomes, number of cases, and PTP per epiweek (N=1484)



Clade key (bar graph)




Weekly percentage testing positive key (line graph)




Gauteng Province, 2021-2022, n = 9724

GP number of genomes, cases, and PTP per epiweek (N=9724)




GP percentage of genomes, number of cases, and PTP per epiweek (N=9724)





Clade key (bar graph)



Weekly percentage testing positive key (line graph)




KwaZulu-Natal Province, 2021-2022, n = 3784



Clade key (bar graph)




Weekly percentage testing positive key (line graph)




Limpopo Province, 2021-2022, n = 2447

LP number of genomes, cases, and PTP per epiweek (N=2447)




LP percentage of genomes, number of cases, and PTP per epiweek (N=2447)



Clade key (bar graph)




Weekly percentage testing positive key (line graph)




Mpumalanga Province, 2021-2022, n = 2482

MP number of genomes, cases, and PTP per epiweek (N=2482)




MP percentage of genomes, number of cases, and PTP per epiweek (N=2482)





Clade key (bar graph)



Weekly percentage testing positive key (line graph)




Northern Cape Province, 2021-2022, n = 1773





Clade key (bar graph)



Weekly percentage testing positive key (line graph)




North West Province, 2021-2022, n = 2064





Clade key (bar graph)



Weekly percentage testing positive key (line graph)




Western Cape Province, 2021-2022, n = 7746



Clade key (bar graph)



Weekly percentage testing positive key (line graph)




Summary

- **Variant of Concern Omicron in South Africa**
 - Dominates 2022 sequencing data at >98% of genomes.
 - While BA.1 (and sub-lineages) was the predominant lineage in January (55%), BA.2 dominated in February (86%) and March (78%).
 - Omicron lineages BA.4 and BA.5 increased in prevalence in March (16%), and together are dominant in April (72%) and May (92%).
- NGS-SA teams are monitoring sequencing data for recombinants.
- Low frequency of previously circulating variants such as Delta still detected in recent data.


¹ <https://github.com/cov-lineages/pango-designation/releases/tag/v1.3>

BA.4 and BA.5 spike mutations


SP
NTD
RBD
RBM
S1
S2
HR1
HR2




BA.4 whole genome mutation prevalence over time



BA.5 whole genome mutation prevalence over time



Omicron spike mutations compared to other VOC/VOIs




Only lineage-defining mutations are pictured here. Low prevalence mutations can be seen on the following slide.

Mutation impact key

- Unknown or unconfirmed impact
- Known/predicted immune escape
- Enhanced infectivity

- Multiple changes within the two immunogenic regions in S1 (NTD and RBD)
 - including a three amino acid insertion
- Accumulation of mutations surrounding the furin cleavage site
 - Including combination of N679K and P681H
- Effect of most spike S2 subunit changes have not been defined, but may be linked to immune escape



Supported by the DSI and the SA MRC



UNIVERSITEIT VAN PRETORIA
UNIVERSITY OF PRETORIA
YUNIBESITHI YA PRETORIA



UNIVERSITEIT
YUNIBESITHI
STELLENBOSCH
UNIVERSITY



UNIVERSITY OF CAPE TOWN
IYUNIBESITHI YASEKAPA • UNIVERSITEIT VAN KAAPSTAD



science & innovation
Department:
Science and Innovation
REPUBLIC OF SOUTH AFRICA



WITS
UNIVERSITY



UNIVERSITY OF
KWAZULU-NATAL
INYUVESI
YAKWAZULU-NATALI



EDCTP

This project (RIA2020EF-3030) is part of the EDCTP2 programme supported by the European Union



NATIONAL HEALTH
LABORATORY SERVICE

University of Stellenbosch & NHLS Tygerberg Virology



Susan Engelbrecht
Wolfgang Preiser
Gert van Zyl
Tongai Maponga
Bronwyn Kleinhans
Shannon Wilson
Karabo Phadu
Tania Stander
Kamela Mahlakwane
Mathilda Claassen
Diagnostic laboratory staff

UKZN-Inkosi Albert Luthuli Central Hospital



Dr Khanyi Msomi
Dr Kerusha Govender
Dr Pravi Moodley
Dr Aabida Khan
Dr Lili Gounder
Dr Kerri Francois
Dr Cherise Naicker
Dr Joedene Chetty

Dr Neli Ngcaba
Dr Tshepiso Mosito
Mr Malcolm Ellapen
Mr Kubendran Reddy
The COVID-19 Bench team

University of KwaZulu-Natal & Africa Health Research Institute



KRISP at UKZN:
Tulio de Oliveira
Richard Lessels
Houriiyah Tegally
Eduan Wilkinson
Jennifer Giandhari
Sureshnee Pillay
Emmanuel James San



AHRI
Alex Sigal
Sandile Cele
Willem Hanekom

University of Cape Town, NHLS & Western Cape Government



NHLS-UCT
Carolyn Williamson
Nei-yuan Hsiao
Diana Hardie
Kruger Marais
Stephen Korsman
Ziyaad Valley-Omar

WCG-UCT
Mary-Anne Davies
Hannah Hussey
Andrew Boule
Masudah Paleker
Theuns Jacobs
Erna Morden

NHLS Greenpoint
Annabel Enoch

Zoonotic arbo and respiratory virus program Centre for Viral Zoonoses Department Medical Virology/ NHLS Tshwane Academic division University of Pretoria



ZARV research program/UP
Marietjie Venter (Head: ZARV)
Adriano Mendes (Postdoc)
Amy Strydom (Postdoc)
Michaela Davis (MSc, intern medical scientist)
Carien van Niekerk

NHLS Tshwane
Prof Simnikiwe Mayaphi (HOD)

Funders:
GIZ/BMBF: African Network for Improved
diagnostics and epidemiology of common and
emerging infectious agents (ANDEMIA)
G7 Global Health fund, Robert Koch Institute, Dr
Fabian Leendertz

National Institute for Communicable Diseases



Centre for Respiratory Diseases & Meningitis

Anne von Gottberg
Thabo Mohale
Daniel Amoako
Josie Everatt
Boitshoko Mahlangu
Noxolo Ntuli
Anele Mnguni
Amelia Buys
Cardia Fourie
Noluthando Duma
Linda de Gouveia
Jackie Kleynhans
Nicole Wolter
Sibongile Walaza
Mignon du Plessis
Stefano Tempia
Mvuyo Makhasi
Cheryl Cohen

Centre for HIV and STIs

Jinal Bhiman
Cathrine Scheepers
Constantinos Kurt Wibmer
Thandeka Moyo
Tandile Hermanus
Frances Ayres
Zanele Molaudzi
Bronwen Lambson
Tandile Hermanus
Mashudu Madzivhandila
Prudence Kgagudi
Brent Oosthuysen
Penny Moore
Lynn Morris

NICD Groups

NICD COVID-19 response team
NICD SARS-CoV-2 Sequencing
Group

Sequencing Core Facility

Zamantungwa Khumalo
Annie Chan
Morne du Plessis
Stanford Kwenda
Phillip Senzo Mtshali
Mushal Allam
Florah Mnyameni
Arshad Ismail



University of the Free State



UFS
Dominique Goedhals
Armand Bester
Martin Myaga
Peter Mwangi
Emmanuel Ogunbayo
Milton Mogotsi
Makgotso Maotoana
Lutfiyya Mohamed



NHLS Division of Virology
Sabeehah Vawda
Felicity Burt
Thokozani Mkhize
Diagnostic laboratory staff



Additional support and collaborators



NHLS

Koeleka Mlisana
Zinhle Makatini
Eugene Elliot
Florette K. Treurnicht
Kathleen Subramoney
Oluwakemi Laguda-Akingba
Shareef Abrahams
Greta Hoyland
Gloria Selabe
Elias Bereda
Jeannette Wadula

Hyrax Biosciences

Simon Travers

Cape Town HVTN Laboratory

Erica Anderson-Nissen
Anneta Naidoo

Ndlovu Research

Hugo Tempelman
CJ Umunnakwe

Lancet

Allison J. Glass
Raquel Viana

Ampath

Terry Marshall
Cindy van Deventer
Eddie Silberbauer

Pathcare Vermaak

Andries Dreyer
Howard Newman
Riaan Writes
Marianne Wolfaardt
Warren Lowman

Bridge-the-Gap

Raymond Rott

Cytespace Africa Laboratories

Christa Viljoen

ARC-OVI

Lia Rotherham

CAPRISA

Salim Abdool Karim
Nigel Garret

UKZN - Big Data

Francesco Pettruccione
Ilya Sinayskiy

University of Oxford

José Lourenço

FioCruz, Brazil

Vagner Fonseca
Marta Giovanetti
Luiz Carlos Junior Alcantara

Africa CDC

John Nkengasong
Sofonias Tessema

Netcare:

Richard Friedland
Craig Murphy
Caroline Maslo
Liza Sitharam

DSI


Glaudina Loots

SA MRC

Glenda Gray



South African genomes submitted per submitting lab, 2020 - 2022 (N=40 993)



NGS-SA Labs

CERi: Centre for Epidemic Response and Innovation

KRISP: KZN Research Innovation and Sequencing Platform

NDLOVU: Ndlovu Research Laboratories

NICD: National Institute for Communicable Diseases

NHLS: National Health Laboratory Service

SU: Stellenbosch University

UCT: University of Cape Town

UFS: University of the Free State

UP: University of Pretoria

Multiple labs from NGS-SA and collaborating public and private laboratories are contributing to sequencing, both as originating and as submitting (pictured here) laboratories.

Currently circulating Variants of Concern (VOC)

WHO label	Pango lineage●	GISAID clade	Nextstrain clade	Additional amino acid changes monitored°	Earliest documented samples	Date of designation
Delta	B.1.617.2	G/478K.V1	21A, 21I, 21J	+S:K417N +S:K484K	India, Oct-2020	VOI: 4-Apr-2021 VOC: 11-May-2021
Omicron*	B.1.1.529	GR/484A	21K	+S:R346K	Multiple countries, Nov-2021	VUM: 24-Nov-2021 VOC: 26-Nov-2021

<https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/> accessed 18 March 2022

●Includes all descendant lineages. See the cov-lineages.org and the Pango network websites for further details.

° Only found in a subset of sequences

Previously circulating Variants of Concern

WHO label	Pango lineage•	GISAID clade	Nextstrain clade	Earliest documented samples	Date of designation
Alpha	B.1.1.7	GRY	20I (V1)	United Kingdom, Sep-2020	VOC: 18-Dec-2020 Previous VOC: 09-Mar-2022
Beta	B.1.351	GH/501Y.V2	20H (V2)	South Africa, May-2020	VOC: 18-Dec-2020 Previous VOC: 09-Mar-2022
Gamma	P.1	GR/501Y.V3	20J (V3)	Brazil, Nov-2020	VOC: 11-Jan-2021 Previous VOC: 09-Mar-2022

<https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/> accessed 18 March 2022

- Includes all descendant lineages. See the cov-lineages.org and the Pango network websites for further details.

Submission of routine specimens for sequencing

- representative of multiple geographic regions (provinces/districts/health facilities) from individuals of
 - all ages
 - over as many time periods during the SARS-CoV-2 epidemic in South Africa
- requested that testing laboratories in both the private and public sectors, submit respiratory samples to their closest NGS-SA sequencing laboratory on a routine basis (ideally every week) as follows, depending on the capacity of the testing laboratory:
 - All positives samples should be sent every week (NGS-SA laboratory will perform random sampling as described below) **OR**
 - A weekly selection of approximately 10%-20% of randomly selected positive samples should be sent every week. Number of selected samples will depend on the size of laboratory and how many other laboratories are drained by the submitting laboratory.

Submission of special interest specimens for sequencing

In addition to routine samples mentioned above, please send specimens separately to above and clearly marked if:

- Suspected vaccine breakthrough (≥ 14 days after vaccine), especially if hospitalised and clinically severe
- Suspected re-infection (≥ 90 days after previous episode), especially if hospitalised and clinically severe
- Prolonged shedding with high SARS-CoV-2 viral loads (i.e. Ct values less than 30 for more than 1 month post-primary diagnosis) in immunocompromised individuals
- Possible animal-to-human transmission
- Suspected cases of importation from another country, especially countries known to harbour SARS-CoV-2 variants of concern or countries with little available information
- Clusters of “unusual” cases (e.g., in terms of disease presentation, patient groups affected, etc.)