



COVID-19 Weekly Testing Summary

Week 25 of 2022

This report summarises national laboratory testing for SARS-CoV-2, the virus causing COVID-19, in South Africa. This report is based on data for specimens reported up to 25 June 2022 (Week 25 of 2022).

Highlights:

- The number of tests reported in week 25 of 2022 (85,028: 59,835 PCR and 25,193 antigen tests) was 5.0% lower than the number of tests reported in the previous week (n=89,549).
- In week 25, the testing rate was 149 per 100,000 persons; highest in Gauteng (207 per 100,000 persons) and lowest in Limpopo (29 per 100,000 persons).
- In week 25, the percentage testing positive was 4.8%, which was 1.6% lower than the previous week.
- In week 25, compared to the previous week, the percentage testing positive decreased in all provinces, except in Mpumalanga where it was unchanged.
- In week 25, the percentage testing positive was highest in the Western Cape (7.9%) and was <7.0% in all other provinces.
- In week 25, the percentage testing positive was highest in the ≥80 years age group (10.1%).

Executive Summary:

- In the period 1 March 2020 through 25 June 2022, 25,424,238 tests for SARS-CoV-2 have been reported nationally: 20,739,119 PCR and 4,685,119 antigen tests.
- The number of tests reported in week 25 of 2022 (n=85 028: 59,835 PCR and 25,193 antigen tests) was 5.0% lower than the number of tests reported in the previous week (n=89,549 in week 24).
- Gauteng reported the largest proportion of tests (38.5%), followed by KwaZulu-Natal (22.5%) and Western Cape (11.0%).
- The overall testing rate decreased slightly from the previous week (149 per 100,000 persons in week 24 to 141 per 100,000 persons in week 25).
- In week 25, testing rates decreased slightly in the Western Cape, Mpumalanga and Northern Cape provinces and were similar to the previous week in all other provinces. The testing rate was highest in Gauteng (207 per 100,000 persons) and lowest in Limpopo (29 per 100,000 persons).
- The testing rate in week 25 was highest in the ≥80 years age group (332 per 100,000 persons).
- In week 25, the percentage testing positive was 4.8%, which was 1.6% lower than the previous week (6.4% in week 24 to 4.8% in week 25, $P<0.001$).
- In the past week, the percentage testing positive decreased by 1.3% in the public sector (4.9% in week 24 to 3.6% in week 25, $P<0.001$) and by 1.9% in the private sector (7.7% in week 24 to 5.8% in week 25, $P<0.001$).
- In week 25, compared to the previous week, the percentage testing positive decreased in all provinces, except in Mpumalanga where it was unchanged.

- The percentage testing positive in week 25 was highest in the Western Cape (7.9%) and was <7.0% in all other provinces.
- In week 25, health sub-districts showing the highest percentage testing positive were concentrated in the Western Cape (n=12), followed by Limpopo (n=3) and North West (n=3).
- In week 25, the percentage testing positive was highest in the ≥80 years age group (10.1%).
- Antigen tests accounted for 29.6% (25,193/85,028) of tests reported in week 25, however the number of antigen tests is likely underestimated due to under-reporting and delayed reporting of antigen tests.
- In week 25 the public sector accounted for 59.0% (14,855/25,193) of antigen tests reported and a decrease in the number of antigen tests reported was observed in all provinces except in KwaZulu-Natal in the past week.
- The mean turnaround time for PCR tests reported in week 25 was 0.8 days; 1.0 day in the public sector and 0.6 days in the private sector. Turnaround times for public sector PCR tests increased in Limpopo and the Eastern Cape and were <2 days in all provinces.
- The mean turnaround time for antigen tests reported in week 25 was 16.6 days in the public sector and 0.2 days in the private sector.




Figure 1. Number of SARS-CoV-2 tests reported by date of specimen collection, South Africa, 4 October 2020 – 25 June 2022. Blue line shows the 7-day moving average of the number of tests reported. Grey bars highlight weekend days and public holidays



Table 1. Weekly number of SARS-CoV-2 tests and positive tests reported, South Africa, 3 January 2021 – 25 June 2022

Week number	Week beginning	No. of tests n (%)	No. of positive tests	Percentage testing positive (%)
1	03-Jan-21	501386 (2.0)	151074	30.1
2	10-Jan-21	418301 (1.6)	104825	25.1
3	17-Jan-21	327534 (1.3)	63283	19.3
4	24-Jan-21	249623 (1.0)	34652	13.9
5	31-Jan-21	203797 (0.8)	22380	11.0
6	07-Feb-21	193340 (0.8)	16476	8.5
7	14-Feb-21	190714 (0.8)	12192	6.4
8	21-Feb-21	184732 (0.7)	10390	5.6
9	28-Feb-21	189731 (0.7)	8695	4.6
10	07-Mar-21	193454 (0.8)	8341	4.3
11	14-Mar-21	185527 (0.7)	8156	4.4
12	21-Mar-21	173275 (0.7)	7356	4.2
13	28-Mar-21	163976 (0.6)	7063	4.3
14	04-Apr-21	180875 (0.7)	7292	4.0
15	11-Apr-21	185350 (0.7)	8847	4.8
16	18-Apr-21	184922 (0.7)	9471	5.1
17	25-Apr-21	160025 (0.6)	9183	5.7
18	02-May-21	193978 (0.8)	13464	6.9
19	09-May-21	240329 (0.9)	19939	8.3
20	16-May-21	248497 (1.0)	24212	9.7
21	23-May-21	262638 (1.0)	29778	11.3
22	30-May-21	270322 (1.1)	36111	13.4
23	06-Jun-21	337915 (1.3)	59453	17.6
24	13-Jun-21	370992 (1.5)	88089	23.7
25	20-Jun-21	432643 (1.7)	118657	27.4
26	27-Jun-21	490277 (1.9)	146650	29.9
27	04-Jul-21	444110 (1.7)	141489	31.9
28	11-Jul-21	320844 (1.3)	100996	31.5
29	18-Jul-21	313377 (1.2)	88476	28.2
30	25-Jul-21	350819 (1.4)	88389	25.2
31	01-Aug-21	372347 (1.5)	88144	23.7
32	08-Aug-21	359664 (1.4)	83390	23.2
33	15-Aug-21	421122 (1.7)	95461	22.7
34	22-Aug-21	392820 (1.5)	78238	19.9
35	29-Aug-21	346236 (1.4)	55109	15.9
36	05-Sep-21	300620 (1.2)	38864	12.9
37	12-Sep-21	260731 (1.0)	24019	9.2

38	19-Sep-21	209230 (0.8)	14018	6.7
39	26-Sep-21	207957 (0.8)	9492	4.6
40	03-Oct-21	198063 (0.8)	6457	3.3
41	10-Oct-21	191812 (0.8)	5045	2.6
42	17-Oct-21	185619 (0.7)	3412	1.8
43	24-Oct-21	177233 (0.7)	2566	1.4
44	31-Oct-21	182981 (0.7)	2106	1.2
45	07-Nov-21	196868 (0.8)	2323	1.2
46	14-Nov-21	197956 (0.8)	4813	2.4
47	21-Nov-21	225551 (0.9)	18977	8.4
48	28-Nov-21	382164 (1.5)	98456	25.8
49	05-Dec-21	493430 (1.9)	175170	35.5
50	12-Dec-21	424671 (1.7)	154982	36.5
51	19-Dec-21	337285 (1.3)	117751	34.9
52	20-Dec-21	216654 (0.9)	66149	30.5
1	02-Jan-22	272609 (1.1)	61154	22.4
2	09-Jan-22	234513 (0.9)	35156	15.0
3	16-Jan-22	208583 (0.8)	24099	11.6
4	23-Jan-22	212696 (0.8)	25840	12.1
5	24-Jan-22	210697 (0.8)	22978	10.9
6	06-Feb-22	203536 (0.8)	20432	10.0
7	13-Feb-22	191313 (0.8)	19103	10.0
8	14-Feb-22	181720 (0.7)	16317	9.0
9	27-Feb-22	172743 (0.7)	13137	7.6
10	06-Mar-22	155535 (0.6)	10688	6.9
11	13-Mar-22	163676 (0.6)	9877	6.0
12	20-Mar-22	146387 (0.6)	9457	6.5
13	27-Mar-22	157550 (0.6)	10258	6.5
14	03-Apr-22	138158 (0.5)	10614	7.7
15	10-Apr-22	120084 (0.5)	12273	10.2
16	17-Apr-22	138871 (0.5)	24142	17.4
17	24-Apr-22	156132 (0.6)	33843	21.7
18	01-May-22	185143 (0.7)	47586	25.7
19	08-May-22	195601 (0.8)	48540	24.8
20	15-May-22	177540 (0.7)	38392	21.6
21	22-May-22	149060 (0.6)	24647	16.5
22	29-May-22	125522 (0.5)	14675	11.7
23	05-Jun-22	118600 (0.5)	10188	8.6
24	12-Jun-22	89549 (0.4)	5719	6.4
25	19-Jun-22	85028 (0.3)	4067	4.8
Total		25,424,238 (100.0)	4,328,450	




Figure 2. Percentage of tests positive for SARS-CoV-2 by date of specimen collection South Africa 1 March 2020 – 25 June 2022. Blue line shows the 7-day moving average of the percentage testing positive. Grey bars highlight weekend days and public holidays.




Figure 3. Testing rate per 100,000 persons by province and week of specimen collection, South Africa, 21 March 2021 – 25 June 2022

Table 2. Weekly number of tests and positive tests reported by province South Africa 5-25 June 2022

Province	Population ^a	5-11 Jun 2022		12-18 Jun 2022		19-25 Jun 2022		Testing rate per 100000	Change in percentage positive from previous week ^b
		No. of tests	No. positive tests (%)	No. of tests	No. positive tests (%)	No. of tests	No. positive tests (%)		
Western Cape	7113776	15862	2267 (14.3)	10614	1098 (10.3)	9383	743 (7.9)	132	-2.4%
Eastern Cape	6676590	9111	901 (9.9)	7355	500 (6.8)	6530	287 (4.4)	98	-2.4%
Northern Cape	1303047	3017	362 (12.0)	1933	168 (8.7)	1718	81 (4.7)	132	-4.0%
Free State	2932441	6929	616 (8.9)	4963	323 (6.5)	4900	188 (3.8)	167	-2.7%
KwaZulu-Natal	11513575	23403	1348 (5.8)	18907	763 (4.0)	19094	569 (3.0)	166	-1.1%
North West	4122854	5928	522 (8.8)	4251	331 (7.8)	4054	237 (5.8)	98	-1.9%
Gauteng	15810388	44505	3420 (7.7)	34133	2103 (6.2)	32722	1578 (4.8)	207	-1.3%
Mpumalanga	4743584	7007	511 (7.3)	5392	307 (5.7)	4535	276 (6.1)	96	0.4%
Limpopo	5926724	2127	184 (8.7)	1679	114 (6.8)	1732	89 (5.1)	29	-1.7%
Unknown		711	57 (8.0)	322	12 (3.7)	360	19 (5.3)		
Total	60142978	118600	10188 (8.6)	89549	5719 (6.4)	85028	4067 (4.8)	141	-1.6%

^a 2021 Mid-year population Statistics SA

^b Current week compared to previous week




Figure 4. Weekly percentage testing positive by province, South Africa, 5-25 June 2022. The horizontal blue line shows the national mean for week 25, beginning 19 June 2022




Figure 5. Testing rates per 100,000 persons and percentage testing positive by age group and sex, South Africa, week 25, 19-25 June 2022




Figure 6. Percentage testing positive by age group and week of specimen collection, South Africa, 3 October 2021 – 25 June 2022



Table 3. Health sub-districts with the highest proportion testing positive based on public and private sector data for the week of 19-25 June 2022

Health district or sub-district	Province	PTP (95% CI)	Previous week
Lephalale	Limpopo	0.144 (0.082-0.205)	0.070 (0.032-0.108)
Umjindi	Mpumalanga	0.139 (0.012-0.266)	0.046 (0.000-0.107)
Maruleng	Limpopo	0.135 (0.042-0.228)	0.120 (0.046-0.194)
Randfontein	Gauteng	0.134 (0.097-0.171)	0.327 (0.287-0.368)
George	Western Cape	0.130 (0.091-0.168)	0.227 (0.192-0.262)
Mthonjaneni	KwaZulu-Natal	0.125 (0.000-0.260)	...
Overstrand	Western Cape	0.120 (0.065-0.174)	0.178 (0.131-0.226)
Tswelopele	Free State	0.108 (0.000-0.224)	0.152 (0.087-0.217)
uMlalazi	KwaZulu-Natal	0.107 (0.026-0.189)	0.108 (0.017-0.198)
CT Khayelitsha	Western Cape	0.106 (0.044-0.168)	0.027 (0.000-0.058)
Nkonkobe	Eastern Cape	0.104 (0.000-0.217)	0.066 (0.003-0.129)
Langeberg	Western Cape	0.102 (0.007-0.198)	0.205 (0.091-0.318)
Witzenberg	Western Cape	0.102 (0.007-0.197)	0.192 (0.114-0.270)
Ditsobotla	North West	0.102 (0.000-0.212)	0.124 (0.031-0.217)
Greater Tubatse	Limpopo	0.101 (0.006-0.196)	0.140 (0.066-0.213)
Breede Valley	Western Cape	0.100 (0.057-0.143)	0.128 (0.091-0.166)
Kagisano/Molopo	North West	0.100 (0.006-0.194)	0.031 (0.000-0.073)
CT Northern	Western Cape	0.098 (0.077-0.119)	0.174 (0.152-0.195)
Oudtshoorn	Western Cape	0.098 (0.029-0.168)	0.183 (0.121-0.245)
Saldanha Bay	Western Cape	0.097 (0.053-0.141)	0.220 (0.169-0.271)
Drakenstein	Western Cape	0.096 (0.063-0.129)	0.117 (0.088-0.145)
Mafikeng	North West	0.093 (0.063-0.123)	0.070 (0.046-0.093)
Mossel Bay	Western Cape	0.091 (0.053-0.128)	0.167 (0.125-0.208)
CT Western	Western Cape	0.089 (0.076-0.101)	0.174 (0.161-0.188)
Lukanji	Eastern Cape	0.088 (0.047-0.130)	0.118 (0.074-0.161)

95% CI: 95% confidence interval; PTP: adjusted positive test proportion; Elements marked in **red** have current week proportions testing positive that are **higher** than and CIs that do not overlap with the previous week proportions and CIs. Elements marked in **blue** have current week proportions testing positive that are **lower** than and CIs that do not overlap with the previous week proportions and CIs




Figure 7. Proportion testing positive by health sub-district in South Africa for the week of 19-25 June 2022. Areas shaded white represent districts in which either (i) no tests were reported (ii) all tests were negative or (iii) the confidence interval exceeded 30%.




Figure 8. Proportion testing positive by health sub-district in the Western Cape Province for the week of 19-25 June 2022. Areas shaded white represent districts in which either (i) no tests were reported (ii) all tests were negative or (iii) the confidence interval exceeded 30%

BUF	Buffalo City	EC122	Mnquma	EC135	Intsika Yethu	EC155	Nyandeni
EC101	Camdeboo	EC123	Great Kei	EC136	Emalahleni	EC156	Mhlontlo
EC102	Blue Crane Route	EC124	Amahlathi	EC137	Engcobo	EC157	King Sabata Dalindyebo
EC103	Ikwezi	EC126	Ngqushwa	EC138	Sakhisizwe	EC441	Mataliele
EC104	Makana	EC127	Nkonkobe	EC141	Elundini	EC442	Umtzimvubu
EC105	Nclambe	EC128	Nxuba	EC142	Sengu	EC443	Mbizana
EC106	Sundays River Valley	EC131	Inxuba Yethemba	EC143	Maletswai	EC444	Ntabankulu
EC107	Baviaans	EC132	Tsolwana	EC144	Gariep	NMAA	Nelson Mandela Bay A
EC108	Kouga	EC133	Inkwanca	EC153	Ngqiza Hill	NMAB	Nelson Mandela Bay B
EC109	Kou-Kamma	EC134	Lukanji	EC154	Port St Johns	NMAC	Nelson Mandela Bay C
EC121	Mbhasane						




Figure 9. Proportion testing positive by health sub-district in the Eastern Cape Province for the week of 19-25 June 2022. Areas shaded white represent districts in which either (i) no tests were reported (ii) all tests were negative or (iii) the confidence interval exceeded 30%.




Figure 10. Proportion testing positive by health sub-district in Northern Cape Province for the week of 19-25 June 2022. Areas shaded white represent districts in which either (i) no tests were reported (ii) all tests were negative or (iii) the confidence interval exceeded 30%.

MAN	Mangaung	FS181	Masilonyana	FS191	Setsoto	FS196	Mantsopa
FS161	Letsemeng	FS182	Tokologo	FS192	Dihlabeng	FS201	Moghaka
FS162	Kopanong	FS183	Tswelopele	FS193	Nketoana	FS203	Ngwathe
FS163	Mohokare	FS184	Matjhabeng	FS194	Maluti a Phofung	FS204	Metsimaholo
FS164	Naledi	FS185	Nala	FS195	Phumelela	FS205	Mafube




Figure 11. Proportion testing positive by health sub-district in Free State Province for the week of 19-25 June 2022. Areas shaded white represent districts in which either (i) no tests were reported (ii) all tests were negative or (iii) the confidence interval exceeded 30%.

ETHN	eThekwin North	KZN233	Indaka	KZN273	The Big 5 False Bay
ETHS	eThekwin South	KZN234	Umtshezi	KZN274	Hlabisa
ETHS	eThekwin South	KZN235	Okhahlamba	KZN275	Mtubatuba
ETHW	eThekwin West	KZN236	Imbabazane	KZN281	Mfolozi
KZN211	Vulamehlo	KZN241	Endumeni	KZN282	uMhlathuze
KZN212	Umdoni	KZN242	Nqutu	KZN283	Ntambanana
KZN213	Umkumbe	KZN244	Mzinga	KZN284	uMlalazi
KZN214	uMuziwabantu	KZN245	Umvoti	KZN285	Mthonjaneni
KZN215	Ezingoleni	KZN252	Newcastle	KZN286	Nkandla
KZN216	Hibiscus Coast	KZN253	Emadlangeni	KZN291	Mandeni
KZN221	uMshwathi	KZN254	Dannhauser	KZN292	KwaDukuza
KZN222	uMngeni	KZN261	eDumbe	KZN293	Ndwedwe
KZN223	Mpofana	KZN262	uPhongolo	KZN294	Maphumulo
KZN224	Impendle	KZN263	Abaqulusi	KZN431	Ingwe
KZN225	The Msunduzi	KZN265	Nongoma	KZN432	Kwa Sani
KZN226	Mkhambathini	KZN266	Ulundi	KZN433	Greater Kokstad
KZN227	Richmond	KZN271	Umhlabyalingana	KZN434	Ubuhlebezwe
KZN232	Emnambithi/Ladysmith	KZN272	Jozini	KZN435	Umkhulu




Figure 12. Proportion testing positive by health sub-district in KwaZulu-Natal Province for the week of 19-25 June 2022. Areas shaded white represent districts in which either (i) no tests were reported (ii) all tests were negative or (iii) the confidence interval exceeded 30%.

NW371	Moretele	NW383	Mafikeng	NW396	Lekwa-Teemane
NW372	Madibeng	NW384	Ditsobotla	NW397	Kagisano/Molopo
NW373	Rustenburg	NW385	Ramotshere Moiloa	NW401	Ventersdorp
NW374	Kgetlengrivier	NW392	Naledi	NW402	Tlokwe City Council
NW375	Moses Kotane	NW393	Mamusa	NW403	City of Matlosana
NW381	Ratlou	NW394	Greater Taung	NW404	Maquassi Hills
NW382	Tswaing				




Figure 13. Proportion testing positive by health sub-district in North West Province for the week of 19-25 June 2022. Areas shaded white represent districts in which either (i) no tests were reported (ii) all tests were negative or (iii) the confidence interval exceeded 30%.

EKUE1	Ekurhuleni East 1	GT421	Emfuleni	JHBA	Johannesburg A	TSH1	Tshwane 1
EKUE2	Ekurhuleni East 2	GT422	Midvaal	JHBB	Johannesburg B	TSH2	Tshwane 2
EKUN1	Ekurhuleni North 1	GT423	Lesedi	JHBC	Johannesburg C	TSH3	Tshwane 3
EKUN2	Ekurhuleni North 2	GT481	Mogale City	JHBD	Johannesburg D	TSH4	Tshwane 4
EKUS1	Ekurhuleni South 1	GT482	Randfontein	JHBE	Johannesburg E	TSH5	Tshwane 5
EKUS2	Ekurhuleni South 2	GT483	Westonaria	JHBF	Johannesburg F	TSH6	Tshwane 6
		GT484	Merafong City	JHBG	Johannesburg G	TSH7	Tshwane 7




Figure 14. Proportion testing positive by health sub-district in Gauteng Province for the week of 19-25 June 2022. Areas shaded white represent districts in which either (i) no tests were reported (ii) all tests were negative or (iii) the confidence interval exceeded 30%.

MP301	Albert Luthuli	MP307	Govan Mbeki	MP316	Dr JS Moroka
MP302	Msukaligwa	MP311	Victor Khanye	MP321	Thaba Chweu
MP303	Mkhondo	MP312	Emalahleni	MP322	Mbombela
MP304	Pixley Ka Seme	MP313	Steve Tshwete	MP323	Umgindi
MP305	Lekwa	MP314	Emakhazeni	MP324	Nkomazi
MP306	Dipaleseng	MP315	Thembisile	MP325	Bushbuckridge




Figure 15. Proportion testing positive by health sub-district in Mpumalanga Province for the week of 19-25 June 2022. Areas shaded white represent districts in which either (i) no tests were reported (ii) all tests were negative or (iii) the confidence interval exceeded 30%.




Figure 17. Number of antigen tests by province and overall percentage antigen tests, South Africa, 1 November 2020 – 25 June 2022. WC Western Cape; EC Eastern Cape; FS Free State; KZN KwaZulu-Natal; GA Gauteng; NC Northern Cape; NW North West; MP Mpumalanga; LP Limpopo




Figure 18. Mean number of days between date of specimen collection and date of test result for PCR tests by week of test result, South Africa, 29 May – 25 June 2022.




Figure 19. Mean number of days between date of specimen collection and date of test result for PCR tests in the public sector by week of test result and province, South Africa, 29 May – 25 June 2022. WC Western Cape; EC Eastern Cape; FS Free State; KZN KwaZulu-Natal; GT Gauteng; NC Northern Cape; NW North West; MP Mpumalanga; LP Limpopo




Figure 20. Mean number of days between date of specimen collection and date of test result for antigen tests by week of test result, South Africa, 29 May – 25 June 2022.

Methods

Testing for SARS-CoV-2 began on 28 January 2020 at the NICD and after the first case was confirmed on 5th March 2020, testing was expanded to a larger network of private and NHLS laboratories. Laboratory testing was conducted for people meeting the case definition for persons under investigation (PUI). This definition was updated several times over the reporting period but at different times included (i) symptomatic individuals seeking testing, (ii) hospitalised individuals for whom testing was done, (iii) individuals in high-risk occupations, (iv) individuals in outbreak settings, and (v) individuals identified through community screening and testing (CST) programmes which were implemented in April 2020 and was discontinued from the week beginning 17th May. CST was implemented differently in different provinces, and ranged from mass screening approaches (including asymptomatic individuals) to screening of individuals in contact with a confirmed case to targeted testing of clusters of cases. Respiratory specimens were submitted to testing laboratories. Testing was performed using reverse transcriptase real-time PCR, which detects SARS-CoV-2 viral genetic material. Laboratories used any one of several in-house and commercial PCR assays to test for the presence of SARS-CoV-2 RNA. Testing for SARS-CoV-2 using rapid antigen-based tests was implemented towards the end of October 2020. Results of reported rapid antigen-based tests are included in this report, however data are incomplete and efforts are ongoing to improve data completeness.

Test results were automatically fed into a data warehouse after result authorisation. We excluded specimens collected outside South Africa and duplicate entries of the same test for an individual. From week 49 of 2020 onwards, test data were reported from the Notifiable Medical Conditions Surveillance System (NMCCS). Date of specimen receipt in the laboratory was used when date of specimen collection was missing. Proportion testing positive (PTP) was calculated as the number of positive tests/total number of tests and presented as percentage by multiplying with 100. Testing rates were calculated using mid-year population estimates from Statistics South Africa and expressed as tests per 100,000 (2019 estimates were used from week 10 – 40 of 2020, 2020 estimates were used from week 41 2021 to week 1 of 2022 and 2021 estimates were used from week 2 of 2022 and onwards). Laboratory turnaround times were calculated as the mean number of days between specimen collection and reporting of the result. Categorical variables were compared using the chi-squared test, with a P-value<0.05 considered statistically significant.

Health district and sub-district (in the metros) level results were mapped based on geo-locatable public (approximately 99% of public sector facilities in the country) and private (approximately 79% of private testing facilities) sector testing facilities. Estimates of overall prevalence were derived using regression techniques. Estimates were adjusted to produce district-specific positive test prevalences based on the national average age and sex profile of testing for that week. This adjustment allows more accurate comparison of the proportion testing positive across districts. Districts with fewer than 20 tests reported during the week have been excluded from the analysis.

Limitations

- A backlog in testing of samples by laboratories affects the reported number of tests. As a result, numbers tested during this period may change in subsequent reports.
- If higher-priority specimens were tested preferentially this would likely result in an inflated proportion testing positive.
- Different and changing testing strategies (targeted vs. mass testing, PCR vs. antigen-based tests or prioritisation of severe or at-risk cases during epidemic waves) used by different provinces makes percentage testing positive and number of reported tests difficult to interpret and compare.

- Health district and sub-district level were mapped based on the testing facility and not place of residence.
- Patient admission status was categorised based on the reported patient facility and may not reflect whether the patient was actually admitted to hospital.
- Antigen tests may be underestimated as they are used in a number of different settings and results may not be reported.