

# **COVID-19 Weekly Testing Summary**


## Week 40 of 2022

This report summarises national laboratory PCR testing for SARS-CoV-2, the virus causing COVID-19, in South Africa. This report is based on data for specimens reported up to 8 October 2022 (week 40 of 2022).

NOTE: From week 28 onwards, only PCR tests are included in the report (i.e. excluding antigen tests).

#### Highlights:

- In the period 1 March 2020 through 8 October 2022, 21,210,108 PCR tests for SARS-CoV-2 have been reported nationally. The number of PCR tests reported in week 40 of 2022 (n=15,162) was 28.5% lower than the number of PCR tests reported in the previous week (n= 21,193 in week 39).
- In week 40 the PCR testing rate was 25 per 100,000 persons. The overall PCR testing rate decreased from the previous week (33 per 100,000 persons in week 39).
- The PCR testing rate in week 40 was highest in Gauteng (36 per 100,000 persons) and lowest in Limpopo (4 per 100,000 persons).
- In week 40, the percentage testing positive was 7.3%, which increased significantly from the previous week (6.6% in week 39, p < 0.05).
- The percentage testing positive in week 40 was highest in Gauteng (10.7%), followed by Western Cape (8.1%), Limpopo (7.1%), Eastern Cape (7.0%), Free State (6.5%), and North West (6.4%). The percentage testing positive was <5.0% in all other provinces.
- In week 40, compared to the previous week, the percentage testing positive increased significantly in Gauteng (p < 0.05). The percentage testing positive did not change in all other provinces (p  $\geq$  0.05).
- The percentage testing positive in week 40 was highest in the ≥80 year's age group (16.6%), followed by 70-74 years (13.8%), 65-69 years (11.4%) and 75-79 years (10.7%) age groups.



**Figure 1.** Number of SARS-CoV-2 PCR tests reported by date of specimen collection, South Africa, 4 October 2020 – 8 October 2022. Blue line shows the 7-day moving average of the number of PCR tests reported. Grey bars highlight weekend days and public holidays.



**Table 1.** Weekly number of SARS-CoV-2 PCR tests and positive tests reported, South Africa, 2 January – 8 October 2022

Week	Week	No. of PCR tests	No. of positive PCR	Percentage	testing
number	beginning	n (%)	tests	positive (%)	coung
1	02-Jan-22	176318 (0.8)	45453	25.8	
2	09-Jan-22	160319 (0.8)	27666	17.3	
3	16-Jan-22	150046 (0.7)	19105	12.7	
4	23-Jan-22	153462 (0.7)	20128	13.1	
5	24-Jan-22	148525 (0.7)	17694	11.9	
6	06-Feb-22	147889 (0.7)	16071	10.9	
7	13-Feb-22	140051 (0.7)	14761	10.5	
8	14-Feb-22	132986 (0.6)	13025	9.8	
9	27-Feb-22	121529 (0.6)	10269	8.4	
10	06-Mar-22	109453 (0.5)	8171	7.5	
11	13-Mar-22	117606 (0.6)	7722	6.6	
12	20-Mar-22	105318 (0.5)	7261	6.9	
13	27-Mar-22	106354 (0.5)	7986	7.5	
14	03-Apr-22	93774 (0.4)	7863	8.4	
15	10-Apr-22	80638 (0.4)	8849	11.0	
16	17-Apr-22	92538 (0.4)	17228	18.6	
17	24-Apr-22	97817 (0.5)	23648	24.2	
18	01-May-22	116957 (0.6)	32989	28.2	
19	08-May-22	126873 (0.6)	34321	27.1	
20	15-May-22	115221 (0.5)	26649	23.1	
21	22-May-22	99444 (0.5)	17368	17.5	
22	29-May-22	84460 (0.4)	10604	12.6	
23	05-Jun-22	77895 (0.4)	7497	9.6	
24	12-Jun-22	63270 (0.3)	4484	7.1	
25	19-Jun-22	61548 (0.3)	3358	5.5	
26	26-Jun-22	47977 (0.2)	2164	4.5	
27	03-Jul-22	43148 (0.2)	2012	4.7	
28	10-Jul-22	38515 (0.2)	1871	4.9	
29	17-Jul-22	36440 (0.2)	1658	4.5	
30	24-Jul-22	34781 (0.2)	1561	4.5	
31	31-Jul-22	34376 (0.2)	1360	4.0	
32	07-Aug-22	28643 (0.1)	1171	4.1	
33	14-Aug-22	30891 (0.1)	1253	4.1	
34	21-Aug-22	28839 (0.1)	1195	4.1	
35	28-Aug-22	28128 (0.1)	1144	4.1	
36	04-Sep-22	27301 (0.1)	1263	4.6	
37	11-Sep-22	27024 (0.1)	1351	5.0	
38	12-Sep-22	24225 (0.1)	1329	5.5	
39	25-Sep-22	21193 (0.1)	1405	6.6	
40	02-Oct-22	15162 (0.1)	1108	7.3	





Figure 2. PCR testing rate per 100,000 persons by province and week of specimen collection, South Africa, 21 March 2021 – 8 October 2022

Week start date (week number) of sample collection



Date of specimen collection

**Figure 3.** Percentage of PCR tests positive for SARS-CoV-2 by date of specimen collection, South Africa, 1 March 2020 – 8 October 2022. Blue line shows the 7-day moving average of the percentage testing positive. Grey bars highlight weekend days and public holidays.




Table 2. Weekly number of PCR tests and positive tests reported by province, South Africa, 18 September – 8 October 2022

		18 - 24 Sep 2022		25 Sep - 1 Oct 2022		2 - 8 Oct 2022		Change in percentage positive	
Province	Population <sup>a</sup>	No. of	No. positive	No. of	No. positive	No. of	No. positive	Testing rate per	from previous
		tests	tests (%)	tests	tests (%)	tests	tests (%)	100,000	week <sup>b</sup>
<b>Western Cape</b>	7113776	3866	196 (5.1)	2781	229 (8.2)	2381	193 (8.1)	33	-0.1%
<b>Eastern Cape</b>	6676590	1142	45 (3.9)	965	51 (5.3)	690	48 (7.0)	10	1.7%
Northern Cape	1303047	277	2 (0.7)	202	1 (0.5)	152	1 (0.7)	12	0.2%
Free State	2932441	959	55 (5.7)	798	54 (6.8)	552	36 (6.5)	19	-0.2%
KwaZulu-Natal	11513575	5773	131 (2.3)	5194	168 (3.2)	3955	124 (3.1)	34	-0.1%
North West	4122854	832	64 (7.7)	773	45 (5.8)	488	31 (6.4)	12	0.5%
Gauteng	15810388	9283	745 (8.0)	8486	778 (9.2)	5769	619 (10.7)	36	1.6%
Mpumalanga	4743584	1683	57 (3.4)	1534	48 (3.1)	930	39 (4.2)	20	1.1%
Limpopo	5926724	410	34 (8.3)	456	31 (6.8)	240	17 (7.1)	4	0.3%
Unknown		0	0(0.0)	4	0 (0.0)	5	0(0.0)		
Total	60142978	24225	1329 (5.5)	21193	1405 (6.6)	15162	1108 (7.3)	25	0.7%

<sup>&</sup>lt;sup>a</sup> 2021 Mid-year population Statistics SA

<sup>&</sup>lt;sup>b</sup> Current week compared to previous week



**Figure 4.** Weekly percentage testing positive (PCR tests only) by province, South Africa, 18 September – 8 October 2022. The horizontal blue line shows the national mean for week 40, beginning 2 October 2022




Figure 5. Percentage testing positive (PCR tests only) by age group and week of specimen collection, South Africa, 3 October 2021 – 8 October 2022

#### **Methods**

Testing for SARS-CoV-2 began on 28 January 2020 at the NICD and after the first case was confirmed on 5th March 2020, testing was expanded to a larger network of private and NHLS laboratories. Laboratory testing was conducted for people meeting the case definition for persons under investigation (PUI). This definition was updated several times over the reporting period but at different times included (i) symptomatic individuals seeking testing, (ii) hospitalised individuals for whom testing was done, (iii) individuals in high-risk occupations, (iv) individuals in outbreak settings, and (v) individuals identified through community screening and testing (CST) programmes which were implemented in April 2020 and was discontinued from the week beginning 17th May. CST was implemented differently in different provinces, and ranged from mass screening approaches (including asymptomatic individuals) to screening of individuals in contact with a confirmed case to targeted testing of clusters of cases. Respiratory specimens were submitted to testing laboratories. Testing was performed using reverse transcriptase real-time PCR, which detects SARS-CoV-2 viral genetic material. Laboratories used any one of several in-house and commercial PCR assays to test for the presence of SARS-CoV-2 RNA. Testing for SARS-CoV-2 using rapid antigen-based tests was implemented towards the end of October 2020 and results of reported rapid antigen-based tests were included in this report until the week 27 report (week starting 3 July 2022). However, as of the week 28 report (week starting 10 July 2022), this report was updated to only include reported PCR tests due to incomplete and delayed reporting of antigen-based tests.

Test results were automatically fed into a data warehouse after result authorisation. We excluded specimens collected outside South Africa and duplicate entries of the same test for an individual. From week 49 of 2020 onwards, test data were reported from the Notifiable Medical Conditions Surveillance System (NMCSS). Date of specimen receipt in the laboratory was used when date of specimen collection was missing. Proportion testing positive (PTP) was calculated as the number of positive tests/total number of tests and presented as percentage by multiplying with 100. Testing rates were calculated using mid-year population estimates from Statistics South Africa and expressed as tests per 100,000 (2019 estimates were used from week 10 of 2020 to week 40 of 2021, 2020 estimates were used from week 1 of 2022, and 2021 estimates were used from week 2 of 2022 onwards). Categorical variables were compared using the chi-squared test, with a P-value<0.05 considered statistically significant.

### **Limitations**

- A backlog in testing of samples by laboratories affects the reported number of tests. As a result, numbers tested during this period may change in subsequent reports.
- If higher-priority specimens were tested preferentially this would likely result in an inflated proportion testing positive.
- Different and changing testing strategies (targeted vs. mass testing, PCR vs. antigen-based tests
  or prioritisation of severe or at-risk cases during epidemic waves) used by different provinces
  makes percentage testing positive and number of reported tests difficult to interpret and
  compare.