


GS-SA Network for Genomic Surveillance in South Africa (NGS-SA)

SARS-CoV-2 Sequencing Update 9 December 2022


























The genomic data presented here are based on South African SARS-CoV-2 sequence data downloaded from GISAID (www.gisaid.org) on 09 Dec 2022 at 14h40




Data license: https://www.gisaid.org/registration/terms-of-use/

Elbe, S., and Buckland-Merrett, G. (2017) Data, disease and diplomacy: GISAID's innovative contribution to global health. Global Challenges, 1:33-46. DOI: 10.1002/gch2.1018 PMCID: 31565258

Shu, Y., McCauley, J. (2017) GISAID: Global initiative on sharing all influenza data – from vision to reality. EuroSurveillance, 22(13) DOI: 10.2807/1560-7917.ES.2017.22.13.30494 PMCID: PMC5388101




GISAID genomes vs total cases, 2020 - 2022 (N=47 435)





Number and percentage of clades by epiweek in South Africa, 2021 – 2022 (40 590*)






Detection Rates: Beta, Delta, C.1.2, recombinants, and Omicron

Detection rates of variants being monitored in South Africa 75 Percentage 25 **Total sequences** 1162 827 1681 1600 627 939 666 1408 3578 3279 1928 1789 5222 2821 1214 1484 2495 1036 May-20 Jun-20 Jul-20 Aug-20 Sep-20 Oct-20 Nov-20 Dec-20 Jan-21 Feb-21 Mar-21 Apr-21 May-21 Jun-21 Jul-21 Aug-21 Sep-21 Oct-21 Nov-21 Dec-21 Jan-22 Feb-22 Mar-22 Apr-22 Jun-22 Jun-22 Jun-22 Sep-22 Oct-22 Nov-22 Nov-22 Nov-22 Nov-22 Nov-22 Nov-22 Nov-22 Nov-20 Nov-20 Dec-20 Jun-20 Ju Month Omicron Omicron Omicron Omicron Omicron Beta (22F/XBB.*) (22A/BA.4)(22C/BA.2.12.1) (21L/BA.2.*) (22E/BQ.1.*) Omicron Omicron Omicron Omicron Delta XBA (21K/BA.1.*) (21M/BA.3) (22B/BA.5)(22D/BA.2.75.*)


^{*}Bars represent percentage prevalence of variant for the month; total number sequences collected for the month are given below the bar

Prevalence of Variants of Concern (VOC) and Variants of Interest (VOI) in





South Africa, 2022, n = 14 299*




Surveillance in South Africa

— 11 - 20 **—** 21 - 30

Eastern Cape Province, 2022, n = 679

Genomes added since last report: 0*




*May include genomes from 2020 and 2021 which are not pictured here and are not included in the slide total.

— 6 - 10 **—** 11 - 20 **—** 21 - 30 **—** 31 - 40 **—** 41 - 50



Free State Province, 2022, n = 481

Genomes added since last report: 0*




*May include genomes from 2020 and 2021 which are not pictured here and are not included in the slide total.

— 11 - 20 — 21 - 30 — 31 - 40 — 41 - 50



Gauteng Province, 2022, n = 4452

Genomes added since last report: 3*



*May include genomes from 2020 and 2021 which are not pictured here and are not included in the slide total.


—— 31 - 40

— 11 - 20 **—** 21 - 30



KwaZulu-Natal Province, 2022, n = 1387

Genomes added since last report: 14*



*May include genomes from 2020 and 2021 which are not pictured here and are not included in the slide total.


— 31 - 40

— 11 - 20 **—** 21 - 30



Limpopo Province, 2022, n = 855

Genomes added since last report: 1*



*May include genomes from 2020 and 2021 which are not pictured here and are not included in the slide total.


— 31 - 40

— 11 - 20 **—** 21 - 30



Mpumalanga Province, 2022, n = 1300

Genomes added since last report: 0*



*May include genomes from 2020 and 2021 which are not pictured here and are not included in the slide total.


—— 31 - 40

— 11 - 20 **—** 21 - 30



Northern Cape Province, 2022, n = 463

Genomes added since last report: 15*




Surveillance in South Africa

*May include genomes from 2020 and 2021 which are not pictured here and are not included in the slide total.

North West Province, 2022, n = 620

Genomes added since last report: 0*



*May include genomes from 2020 and 2021 which are not pictured here and are not included in the slide total.


—— 31 - 40

— 11 - 20 **—** 21 - 30



Western Cape Province, 2022, n = 4055

Genomes added since last report: 125*



*May include genomes from 2020 and 2021 which are not pictured here and are not included in the slide total.

—— 31 - 40

— 11 - 20 **—** 21 - 30



Summary

Sequencing update


- All provinces have sequences for September, and all provinces except the Eastern Cape have sequences for October.
- All provinces except the Eastern Cape and the Free State have sequences for November.

Variant of Concern Omicron in South Africa

- Omicron continued to dominate in September (98%), October (98%) and November (99.5%).
- Omicron BA.5 and sub-lineages were dominant in September (78%), October (90%) and currently make up 92% of November data.
- BA.2.75.* was detected at a low prevalence (≤1%) in September, October and November.
- XAY was detected in September, October and November at a low prevalence (≤2%).
- BQ.1 and sub-lineages have been detected in September, October and November
 - September (1%) BQ.1*: n=1 in the Northern Cape, n=6 in the Western Cape, n=1 in Gauteng
 - October (6%) BQ.1*: n=2 in KwaZulu-Natal, n=14 in Gauteng, n=25 in the Western Cape, n=3 in the Northern Cape, n=1 in Mpumalanga
 - November (13%) BQ.1*: n=17 in the Western Cape, n=8 in Gauteng, n=1 in the Northern Cape
- XBB has been detected in October and November
 - October (1%): n=2 in Gauteng, n=3 in the Western Cape
 - November (3%): n=4 in Gauteng, n=2 in the Western Cape
- Delta detected at low frequency until July, and once in October.




Spike protein mutation* profile of Variants of Interest and Concern



- Multiple changes within the two immunogenic regions in S1 (NTD and RBD)
 - Including a three amino acid insertion
- Accumulation of mutations surrounding the furin cleavage site
 - Including combination of N679K and P681H
- Effect of most spike S2 subunit changes have not been defined, but may be linked to immune escape




BA.4 and BA.5 spike mutations*




NTD

RBD

BA.4 whole genome mutation prevalence over time







Epi week (no. of genomes)

Epi week (no. of genomes)

BA.5 whole genome mutation prevalence over time



굦



















UNIVERSITY OF ™ KWAZULU-NATAL

INYUVESI YAKWAZULU-NATALI



ΛΛ

EDCTP







University of Stellenbosch & NHLS Tygerberg Virology



NHLS Greenpoint

This project has

ceived funding from

he European Union's

Horizon Europe

Research and

under grant No.

 $\Lambda \Lambda$

EDCTP

Samrce

Innovation Actions

Annabel Enoch



Susan Engelbrecht **Wolfgang Preiser** Gert van Zyl Tongai Maponga **Bronwyn Kleinhans Shannon Wilson** Karabo Phadu Tania Stander Kamela Mahlakwane Mathilda Claassen Diagnostic laboratory staff

UKZN-Inkosi Albert Luthuli Central Hospital



Dr Kerri Francois

Dr Cherise Naicker

Dr Joedene Chetty



Dr Khanvi Msomi Dr Neli Ngcaba Dr Kerusha Govender Dr Tshepiso Mosito Dr Pravi Moodlev Mr Malcolm Ellapen Dr Aabida Khan Mr Kubendran Reddy Dr Lili Gounder

The COVID-19 Bench team

University of KwaZulu-Natal & Africa Health Research Institute



KRISP at UKZN: Tulio de Oliveira Richard Lessels Houriivah Tegally Eduan Wilkinson Jennifer Giandhari Sureshnee Pillav

Emmanuel James San

AHRT AFRICA
RESEARCH
RESEARCH
RESEARCH
RESEARCH

Alex Sigal Sandile Cele Willem Hanekom

University of Cape Town, NHLS & Western Cape Government

health



NHLS-UCT

Carolyn Williamson Nei-yuan Hsiao Diana Hardie Kruger Marais Stephen Korsman

Mary-Anne Davies Hannah Hussey Andrew Boulle Masudah Paleker Theuns Jacobs

Erna Morden

WCG-UCT









UCT, IDM and CIDRI-Africa

Deelan Doolabh Arash Iranzadeh Lynn Tyers Innocent Mudau Nokuzola Mbhele Fezokuhle Khumalo Thabang Serakge Bruna Galvão Linda Boloko Arghavan Alisoltani

(U. California)

Robert Wilkinson Darren Martin Nicola Mulder Wendy Burgers Ntobeko Ntusi Rageema Joseph







Zoonotic arbo and respiratory virus program **Centre for Viral Zoonoses Department Medical Virology/ NHLS Tshwane Academic division University of Pretoria**



ZARV research program/UP

Marietjie Venter (Head: ZARV) Adriano Mendes (Postdoc) Amy Strydom (Postdoc) Michaela Davis (MSc, intern medical scientist) Carien van Niekerk



NHLS Tshwane

Prof Simnikiwe Mayaphi (HOD)

Funders:

GIZ/BMBF: African Network for Improved diagnostics and epidemiology of common and emerging infectious agents (ANDEMIA) G7 Global Health fund, Robert Koch Institute, Dr Fabian Leendertz

National Institute for Communicable Diseases

Centre for HIV and STIs

Constantinos Kurt Wibmer

Cathrine Scheepers

Thandeka Movo

Frances Ayres

Zanele Molaudzi

Bronwen Lambson

Tandile Hermanus

Prudence Kgagudi

Brent Oosthuysen

Mashudu Madzivhandila

Tandile Hermanus

Jinal Bhiman



Centre for Respiratory Diseases & Meningitis

Anne von Gottberg Thabo Mohale **Daniel Amoako** Josie Everatt Boitshoko Mahlangu Noxolo Ntuli Anele Mnguni Amelia Buys Cardia Fourie Noluthando Duma Linda de Gouveia Jackie Kleynhans Nicole Wolter Sibongile Walaza

Mignon du Plessis

Stefano Tempia

Mvuyo Makhasi

Cheryl Cohen

NICD Groups

Penny Moore

Lynn Morris

NICD COVID-19 response team NICD SARS-CoV-2 Sequencing Group

Sequencing Core Facility

Zamantungwa Khumalo Annie Chan Morne du Plessis Stanford Kwenda Phillip Senzo Mtshali Mushal Allam Florah Mnyameni Arshad Ismail












University of the **Free State**



UFS

Dominique Goedhals Armand Bester Martin Myaga Peter Mwangi **Emmanuel Ogunbayo** Milton Mogotsi Makgotso Maotoana Lutfiyya Mohamed





Sabeehah Vawda **Felicity Burt** Thokozani Mkhize Diagnostic laboratory staff









Additional support and collaborators











FIOCRUZ

NHLS Koeleka Mlisana Zinhle Makatini **Eugene Elliot** Florette K. Treurnicht

Kathleen Subramoney Oluwakemi Laguda-Akingba **Shareef Abrahams** Greta Hoyland Gloria Selabe Elias Bereda Jeannette Wadula

Hyrax Biosciences Simon Travers

Cape Town HVTN Laboratory Erica Anderson-Nissen Anneta Naidoo

Ndlovu Research **Hugo Tempelman** CJ Umunnakwe

Lancet Allison J. Glass Raquel Viana

Ampath Terry Marshall Cindy van Deventer **Eddie Silberbauer**

Pathcare Vermaak Andries Dreyer Howard Newman Riaan Writes Marianne Wolfaardt Warren Lowman

Bridge-the-Gap Raymond Rott

Cytespace Africa Laboratories Christa Viljoen

ARC-OVI Lia Rotherham **CAPRISA**

Salim Abdool Karim Nigel Garret

UKZN - Big Data Francesco Pettruccione Ilya Sinayskiy

University of Oxford José Lourenço

FioCruz, Brazil Vagner Fonseca Marta Giovanetti Luiz Carlos Junior Alcantara

Jean Maritz

Africa CDC

John Nkengasong Sofonias Tessema

Netcare

Richard Friedland Craig Murphy Caroline Maslo Liza Sitharam

DSI Glaudina Loots

SA MRC Glenda Gray

Pathcare N1 City

Nadine Cronje Petra Raimond Kim Hoek




















South African genomes submitted per submitting lab, 2020 - 2022 (N=47 435)



NGS-SA Labs

CERI: Centre for Epidemic Response

and Innovation

KRISP: KZN Research Innovation and

Sequencing Platform

NDLOVU: Ndlovu Research

Laboratories

NICD: National Institute for

Communicable Diseases

NHLS: National Health Laboratory


Service

SU: Stellenbosch University

UCT: University of Cape Town

UFS: University of the Free State

UP: University of Pretoria





Currently circulating Variants of Concern (VOC)

WHO label	Pango Iineage•	GISAID clade	Nextstrain clade	Additional amino acid changes monitored°	Earliest documented samples	Date of designation
Omicron*	B.1.1.529	GR/484A	21K, 21L, 21M, 22A, 22B, 22C, 22D	+S:R346K +S:L452X +S:F486V	Multiple countries, Nov-2021	VUM: 24-Nov-2021 VOC: 26-Nov-2021

^{*} Includes BA.1, BA.2, BA.3, BA.4, BA.5 and descendent lineages. It also includes BA.1/BA.2 circulating recombinant forms such as XE. WHO emphasizes that these descendant lineages should be monitored as distinct lineages by public health authorities and comparative assessments of their virus characteristics should be undertaken.

[•] Only found in a subset of sequences

Omicron subvariants under monitoring

Pango lineage [#] (+ mutation)	GISAID clade	Nextstrain clade	Relationship to circulating VOC lineages	Spike genetic features	Earliest documented samples
BA.5** (+R346X or +K444X or +V445X or +N450D or +N460X)	GRA	22B	BA.5 sublineages (e.g. BF.7, BF.14, BQ.1)	BA.5 + one or more of these mutations: S:R346X, S:K444X, S:V445X , S:N450D or S:N460X	07-02-2022
BA.2.75***	GRA	22D	BA.2 sublineage	BA.2.75: BA.2 + S:K147E, S:W152R, S:F157L, S:I210V, S:G257S, S:D339H, S:G446S, S:N460K, S:Q493R reversion BA.2.75.2: BA.2.75 + S:R346T, S:F486S, S:D1199N	31-12-2021
BJ.1****	GRA	21L	BA.2 sublineage (B.1.1.529.2.10.1.1)	BA.2+S:V83A, S:Y144-, S:H146Q, S:Q183E, S:V213E, S:G339H, S:R346T, S:L368I, S:V445P, S:G446S, S:V483A, S:F490V, S:G798D, S:S1003I	06-09-2021
BA.4.6	GRA	22A	BA.4 sublineage	BA.4+S:R346T, S:N658S	20-07-2020
XBB ^{\$}		recombinant	Recombinant of BA.2.10.1 and BA.2.75 sublineages, i.e. BJ1 and BM.1.1.1, with a breakpoint in S1	BA.2+ S:V83A, S:Y144-, S:H146Q, S:Q183E, S:V213E, S:G252V, S:G339H, S:R346T, S:L368I, S:V445P, S:G446S, S:N460K, S:F486S, S:F490S	13-08-2022
BA.2.3.20 [§]	GRA	21L	BA.2 sublineage	BA.2+ S:M153T, S:N164K, S:H245N, S:G257D, S:K444R, S:N450D, S:L452M, S:N460K, S:E484R	15-08-2022

^{*} these subvariants are tracked under Omicron unless/until sufficient evidence arises that the virus characteristics are substantially different from what is known about the VOC they belong to. If this evidence arises, WHO will decide, in consultation with the TAG-VE, if designation of the emerging variant warrants a separate WHO label.

[#] includes descendent lineages

^{**} additional mutations outside of the spike protein: N:G30-, N:S33F, N:E136D, ORF1a:Q556K, ORF1a:L3829F, ORF1b:Y264H, ORF1b:M1156I, ORF9b:P10F, ORF9b:D16G, ORF9b:M26-, ORF9b:A29I, ORF9b:V30L.

^{***} additional mutation outside the spike protein: ORF1a:S1221L, ORF1a:P1640S, ORF1a:N4060S; ORF1b:G662S; E:T11A

^{****} additional mutations outside of the spike protein: Mutations: M:D3Y, N:T282I, ORF1a:K47R, ORF1b:G662S, ORF1b:S959P, ORF7a:I110T

^{\$} additional mutations outside of the spike protein: E:T11A, ORF1a:K47R, ORF1b:G662S, ORF1b:S959P, ORF8:G8*

[§] additional mutations outside of the spike protein: ORF1a:T727I, ORF1a:I1714T, ORF1a:M2169V, ORF1a:T2174I, ORF1a:T2648I, ORF1a:A2909V, ORF1a:Q3922R, ORF1b:T1404M, ORF3a:L140F, ORF9b:D89E

Previously circulating Variants of Concern

WHO label	Pango Iineage•	GISAID clade	Nextstrain clade	Earliest documented samples	Date of designation
Alpha	B.1.1.7	GRY	20I (V1)	United Kingdom, Sep-2020	VOC: 18-Dec-2020 Previous VOC: 09-Mar-2022
Beta	B.1.351	GH/501Y.V2	20H (V2)	South Africa, May-2020	VOC: 18-Dec-2020 Previous VOC: 09-Mar-2022
Gamma	P.1	GR/501Y.V3	20J (V3)	Brazil, Nov-2020	VOC: 11-Jan-2021 Previous VOC: 09-Mar-2022
Delta	B.1.617.2	G/478K.V1	21A, 21I, 21J	India, Oct-2020	VOI: 4-Apr-2021 VOC: 11-May-2021 Previous VOC: 7-Jun-2022

[•] Includes all descendant lineages. See the cov-lineages.org and the Pango network websites for further details.

https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/ accessed 12 August 2022

Submission of routine specimens for sequencing

- representative of multiple geographic regions (provinces/districts/health facilities) from individuals of
 - all ages
 - over as many time periods during the SARS-CoV-2 epidemic in South Africa
- requested that testing laboratories in both the private and public sectors, submit respiratory samples to their closest NGS-SA sequencing laboratory on a routine basis (ideally every week) as follows, depending on the capacity of the testing laboratory:
 - All positives samples should be sent every week (NGS-SA laboratory will perform random sampling as described below) OR
 - A weekly selection of approximately 10%-20% of randomly selected positive samples should be sent every week. Number of selected samples will depend on the size of laboratory and how many other laboratories are drained by the submitting laboratory.

Submission of special interest specimens for sequencing

In addition to routine samples mentioned above, please send specimens separately to above and clearly marked if:

- Suspected vaccine breakthrough (≥14 days after vaccine), especially if hospitalised and clinically severe
- Suspected re-infection (≥90 days after previous episode), especially if hospitalised and clinically severe
- Prolonged shedding with high SARS-CoV-2 viral loads (i.e. Ct values less than 30 for more than 1 month post-primary diagnosis) in immunocompromised individuals
- Possible animal-to-human transmission
- Suspected cases of importation from another country, especially countries known to harbour SARS-CoV-2 variants of concern or countries with little available information
- Clusters of "unusual" cases (e.g., in terms of disease presentation, patient groups affected, etc.)