



## **INTERIM SITUATION REPORT, 19 May 2023**

### **(Based on laboratory testing data up until 11 May 2023)**

Issued by the National Institute for Communicable Diseases based on laboratory testing data

#### **Highlights**

- The NICD has tested 5916 serum samples for measles since epidemiological week 40, 2022, of which 1004 (17%) were confirmed positive, all from outbreak-affected provinces. In the past weeks (week 18 up until week 19, 11/05/2023) there have been 7 laboratory-confirmed measles cases detected across the country, of which the majority were from Limpopo (7).
- The percentage of samples testing positive (PTP) increased from 10% of 96 samples tested in week 17 to 28% of 61 samples tested in week 18.
- Measles outbreak has been declared in all the provinces in South Africa except for the Eastern Cape. In week 18, seven (7) new cases were reported from Limpopo province. No new cases were reported in Mpumalanga, Northern Cape, Western Cape and KwaZulu-Natal provinces in the past four weeks.
- The measles strain detected in Limpopo province and North West province is genotype D8 which is similar to the strain in Zimbabwe in the 2022 outbreak.
- In the provinces where an outbreak has been declared, the most affected age groups are still the 5-9-year-olds (43%) with a considerable proportion of cases reported among the 1-4 (23%) and 10-14 age groups (20%).
- Vaccination campaigns included all children including those aged 1 to 14 years of age and have been extended across provinces to obtain improved coverage rates.
- The majority of cases (67%) were reported from primary healthcare facilities, and the highest proportion of cases reported from hospitals (59%) was reported in children under the age of one.
- Nationally, the reproduction number as of 2023-05-03 was estimated to be 0.44 (0.19 – 0.91), suggesting that infection incidence is decreasing. There is a 97% chance that the reproduction number was below 1 as of 2023-05-03.
- At the provincial level, the reproduction number as of 2023-05-03 was estimated to be 0.63 (0.4 – 0.84) in Limpopo, 0.53 (0.14 – 1.3) in Mpumalanga, 0.5 (0.31 – 0.73) in Gauteng, and 0.36 (0.18 – 0.67) in North West, suggesting that infection incidence is decreasing. Note that the wide credible interval for Mpumalanga is due to the small number of cases in the estimation period; the high upper bound does not reflect an increase in transmission over previous reports.
- At the district level, the reproduction number as of 2023-05-03 was estimated to be 0.76 (0.5 – 1) in Waterberg, suggesting that infection incidence is decreasing. The reproduction number in all other districts that have had at least 40 detected cases since week 40 of 2022 is estimated to be below 1.

#### **Outbreak Overview**

From epidemiological week 40, 2022 (ending 8 October 2022) to week 19, 2023 the NICD has tested 5916 serum samples for measles of which 1004 (17%) were confirmed measles cases. The number of samples submitted and the percentage of laboratory-confirmed measles-positive cases are shown in Figure 1. From epidemiological week 40 of 2022 to week 19 of 2023, 998 laboratory-confirmed cases were reported from eight provinces with declared measles outbreaks; Limpopo (432 cases), Mpumalanga (108 cases), North West (217 cases), Gauteng (170 cases), Free State (30 cases), Western Cape (14), KwaZulu-Natal (20) and Northern Cape (7) (Table 1). The geographical distribution of cases across South Africa from week 40 of 2022 until week 19 of 2023 is shown in Figure 2. The number of blood samples and throat swabs submitted to the NICD for measles serology and PCR testing decreased from 90 in week 17 to 61 in week 18 (Figure 3).




Figure 1. Number of serum samples submitted to the NICD for measles, week 40 2022, until week 18, 2023, and the number (dark green) and % tested positive (red line), by epidemiological week using the date the specimen was collected. \*Data from week 19 will be updated in next week's situation report, when complete data from samples collected that week becomes available.

\*Note: Data is subject to change as new results are added or updated. Please contact Mr Tshepo Motsamai ([tshepom@nicd.ac.za](mailto:tshepom@nicd.ac.za)) to update data elements




Figure 2. Distribution of laboratory-confirmed measles cases by testing site (red dots – the size of the dot indicates the number of cases from that facility) and district of South Africa (deepening colour of blue indicates the total number of cases by sub-district), from week 40 to week 19, 2023.




Figure 3. Number measles of tests conducted from week 40 2022, until week 19, 2023, by province and epidemiological week using the date the specimen was collected. \*Data from week 18 represent partial data, and will be updated in next week's situation report, when complete data from samples collected that week becomes available.

\*Note: Data is subject to change as new results are added or updated. Please contact Mr Tshepo Motsamai ([tshepom@nicd.ac.za](mailto:tshepom@nicd.ac.za)) to update data elements

## Reproduction Number



The figure below shows the national time-varying reproduction number over the past 90 days. The estimated reproduction number reached a peak of approximately 1.3 in mid-January and has since declined, crossing the threshold value of 1 in mid-February. Nationally, the reproduction number as of 2023-05-03 was estimated to be 0.44 (0.19 – 0.91), suggesting that infection incidence is decreasing.




Figure 4. National time-varying reproductive estimate from early December 2022 to early May 2023.

The weekly report on nowcasts and forecasts for measles in South Africa is available at <https://www.sacema.org/sacema-nicd-measles-forecast/>

\*Note: Data is subject to change as new results are added or updated. Please contact Mr Tshepo Motsamai ([tshepom@nicd.ac.za](mailto:tshepom@nicd.ac.za)) to update data elements

Table 1. Cases of laboratory-confirmed measles tested by the NICD from all provinces in South Africa from epidemiological week 40, 2022 to week 19, 2023. Outbreak-associated cases are contained within the red bordered cells\* (FS=Free State; GP=Gauteng; KZN=KwaZulu-Natal; LP=Limpopo; MP=Mpumalanga NW=North West; NC=Northern Cape, WC = Western Cape). \* A measles outbreak is classified as three or more confirmed laboratory measles cases reported within 30 days of onset of disease, in a district. \*Data from week 19 represents partial data, and will be updated in next week's situation report, when complete data from samples collected that week becomes available.

Epi Week	EC	FS	GP	KZN	LP	MP	NW	NC	WC	Total
40, 2022			1		2					3
41, 2022					5					5
42, 2022			1		4		1		1	7
43, 2022	1				11					12
44, 2022				1	19	2				22
45, 2022		1	1		12	3	1	1	1	20
46, 2022			1	1	9	8				19
47, 2022		1	2		18	15	4	1	1	42
48, 2022			1		18	17	4			40
49, 2022		3	2	2	10	14	18	1	1	51
50, 2022			3		16	6	30			55
51, 2022		3	3	1	7	5	28			47
52, 2022		2	1		7	5	24	1		40
01, 2023		3	1		7	1	13		1	26
02, 2023		1	2		3	4	7			17
03, 2023	1	4	9		9	5	11		1	40
04, 2023	1	2	10	2	9	5	9		1	39
05, 2023		2	12	2	20	2	14		1	53
06, 2023	1	1	17	3	19	3	10		1	55
07, 2023			19	3	26	2	9	1	1	61
08, 2023	2	1	14	1	20	5	8	2		53
09, 2023		3	19	1	26	4	8		1	62
10, 2023		1	17		27		11			56
11, 2023		1	9	1	22					33
12, 2023			7		29	1	4		1	42
13, 2023			6	1	22		2			31
14, 2023	0		2	1	17					20
15, 2023			3		11	1			2	17
16, 2023			3		5					8
17, 2023					10					10
18, 2023		1	3		12	0	1	0	0	17
19, 2023	0	0	1	0	0	0	0	0	0	1
<b>Total</b>	<b>6</b>	<b>30</b>	<b>170</b>	<b>20</b>	<b>432</b>	<b>108</b>	<b>217</b>	<b>7</b>	<b>14</b>	<b>1004</b>

\*Note: Data is subject to change as new results are added or updated. Please contact Mr Tshepo Motsamai ([tshepom@nicd.ac.za](mailto:tshepom@nicd.ac.za)) to update data elements




Figure 5. The epidemiological curve of the number of laboratory-confirmed measles cases in South Africa from week 40, 2022 to week 19 2023 (ending week 18 up until week 19, 11/05/2023) by specimen collection dates and by province, indicating the weeks in which outbreaks were declared in Limpopo, Mpumalanga, North West, Gauteng, Free State, Western Cape, Northern Cape and Kwa-Zulu Natal provinces. \*Data from week 19 represent partial data, and will be updated in next week’s situation report, when complete data from samples collected that week becomes available.

The age of laboratory-confirmed cases across the eight provinces ranges from two months to 67 years (Table 2). The majority of cases 427, (43%) were in the 5-9-year age group, followed by 233 (23%) in the 1-4-year age group and 197 (20%) in the 10-14-year age group. The attack rates are highest among age groups 1-4 and 5-9 (Table 2). In the provinces where a measles outbreak has been declared, 109 (10.9%) of the 998 cases were vaccinated, 131 (13.1%) were unvaccinated, and the vaccination status of 758 (76.5%) is unknown (Table 3). The age groups with the highest number of vaccinated cases are those aged 1-4 years and those aged 5-9 years (Table 4). Whilst the NICD is presently not able to provide data on hospital admission rates nor measles mortality rates, Table 5 reflects the number and proportion of laboratory-confirmed measles cases that originate from hospitals as opposed to primary healthcare facilities. Whilst cases that are seen at hospitals may not necessarily be admitted, this proportion gives us an indication of the severity of illness, as patients consulted tertiary care facilities.

\*Note: Data is subject to change as new results are added or updated. Please contact Mr Tshepo Motsamai ([tshepom@nicd.ac.za](mailto:tshepom@nicd.ac.za)) to update data elements

Table 2. Age distribution of laboratory-confirmed measles cases from epidemiological week 40, 2022 to week 19, 2023, in provinces with a declared measles outbreak with age-specific attack rates.

Age group	FS		GP		LP		MP		NW		WC		NC		KZN		Total	
	# cases	AR	# cases	AR	# cases	AR	# cases	AR	# cases	AR	# cases	AR	# cases	AR	# cases	AR	# cases	AR
<1 year	4	7.51	16	6.07	19	14.40	5	5.52	8	9.90	3	2.50	0	0.00	1	0.41	56	5.55
1-4 years	11	5.24	33	3.18	85	15.94	32	9.17	56	17.79	6	1.29	2	1.97	8	0.83	233	5.86
5-9 years	12	4.49	70	5.58	195	29.26	38	8.89	99	25.28	1	0.18	3	2.38	9	0.76	427	8.74
10-14 years	3	1.04	28	2.28	103	15.43	22	4.75	36	8.82	2	0.34	2	1.59	1	0.08	197	3.97
≥15 years	0	0.00	23	0.19	30	0.76	11	0.32	18	0.60	2	0.04	0	0.00	1	0.01	85	0.22
<b>Total</b>	<b>30</b>	<b>1.03</b>	<b>170</b>	<b>1.06</b>	<b>432</b>	<b>7.27</b>	<b>108</b>	<b>2.29</b>	<b>217</b>	<b>5.18</b>	<b>14</b>	<b>0.19</b>	<b>7</b>	<b>0.53</b>	<b>20</b>	<b>0.17</b>	<b>998</b>	<b>1.85</b>

FS= Free State; GP= Gauteng; KZN=KwaZulu-Natal; LP=Limpopo; MP=Mpumalanga; NW=North West; WC=Western Cape; NC= Northern Cape; AR = attack rate per 100,000 children within the age-band, denominators from mid-year population estimates, 2022, StatsSA

Table 3. Vaccination status for laboratory-confirmed measles cases from epidemiological week 40, 2022 to week 19, 2023 in provinces with a declared measles outbreak.

Vaccination status	FS	GP	LP	MP	NW	WC	NC	KZN	Total
Vaccinated	9	13	37	15	18	7	3	7	109 (10.9%)
Unvaccinated	3	13	66	18	30	0	0	1	131 (13.1%)
Unknown	18	144	329	75	169	7	4	12	758 (73.9%)
<b>Total</b>	<b>30</b>	<b>170</b>	<b>432</b>	<b>108</b>	<b>217</b>	<b>14</b>	<b>7</b>	<b>20</b>	<b>998</b>

Table 4: Age distribution of vaccinated persons from epidemiological week 40, 2022 to week 19, 2023 in provinces with a declared measles outbreak.

Age group	FS	GP	LP	MP	NW	WC	NC	KZN	Total
< 1 year	2	2	3	0	0	3	0	0	10
1 – 4 years	2	4	12	2	5	3	0	3	31
5 – 9 years	2	7	17	9	12	1	2	4	54
10 – 14 years	3	0	5	4	1	0	1	0	14
≥15 years	0	0	0	0	0	0	0	0	0
<b>Total</b>	<b>9</b>	<b>13</b>	<b>37</b>	<b>15</b>	<b>18</b>	<b>7</b>	<b>3</b>	<b>7</b>	<b>109</b>

Table 5. The facility type where laboratory-confirmed measles cases have been identified, for epidemiological week 40, 2022 to week 19, 2023, South Africa. Submission of a specimen from a hospital may suggest (but is not firm evidence) that the patient was admitted. The number of admissions will be lower than the number of cases reported from hospitals.

Reporting Health Facility	<1 year	1-4 years	5-9 years	10-14 years	≥15 years	Total
From PHC/CHC/other	23	154	293	137	58	665(67)
From a hospital (%)	33 (59)	79 (34)	134 (31)	60 (31)	27 (32)	333(33)
<b>Total</b>	<b>56</b>	<b>233</b>	<b>427</b>	<b>197</b>	<b>85</b>	<b>998</b>



## An Overview of the Outbreak in the Limpopo Province

In total, 432 cases of laboratory-confirmed measles were reported between epidemiological week 40, 2022 to week 19, 2023 with the majority of the measles cases reported in the Waterberg, Greater Sekhukhune and Mopani districts. Figure 6 shows an epidemiological curve from week 40, 2022 to week 19 of 2023 in Limpopo province. Waterberg district reported the highest number of measles cases which is 164 cases, Mopani district reported 97 cases, Greater Sekhukhune district reported 128 cases, Vhembe district reported 35 cases and Capricorn district reported eight cases. Dilokong Hospital reported 50 cases out of the 128 from Greater Sekhukhune. Amongst the 35 cases reported from the Vhembe district, 35 cases originated from Makhado (Louis Trichardt Hospital and Clinic). In the Waterberg district, 52 cases of 164 have been reported from Witpoort Hospital in Lephalale. The age of measles cases across Limpopo ranged from 4 months to 42 years.

Measles virus infection affected mostly the age group 5-9 years (Table 2), with an attack rate of 29.26 per 100,000 persons. This was followed by the 1-4 years age group with an attack rate of 15.94 per 100,000 persons. Of the 432 measles cases in Limpopo province, 329 (76%) had an unknown –vaccination status, 37 (9%) were vaccinated, and 66 (15%) were unvaccinated (Table 3).




Figure 6. The epidemiological curve of the number of laboratory-confirmed measles cases by districts of Limpopo Province from epidemiological week 40, 2022 to week 19, 2023 by specimen collection dates

## North West

A total of 217 laboratory-confirmed measles cases have been reported in North West Province since epidemiological week 40, 2022 (Figure 7). An outbreak was declared in North West province on 02 December 2022 (epidemiological week 48, 2022) after three laboratory-confirmed cases were reported in Ngaka Modiri Molema district. The majority of the laboratory-confirmed cases are among children aged 5-9 years, with 99 cases and an attack rate of 25.28 per 100,000 persons, followed by those aged 1-4 years with 56 cases, with an attack rate of 17.79 per 100,000 persons (Table 2). A total of 18 of the 216 cases were vaccinated and 169 had unknown vaccination status (Table 3). Of these 217 cases, the majority (180) were reported from the Ngaka Modiri Molema district, with 73 cases reported from a single clinic, Lonely Park Clinic in Mahikeng. Twenty-two cases were reported from Bojanala Platinum district, eight cases from Dr Kenneth Kaunda district, and seven cases from Dr Ruth Segomotsi Mompoti district.




Figure 7. The epidemiological curve of the number of laboratory-confirmed measles cases in districts of North West Province from epidemiological week 42, 2022 to week 19, 2023 by specimen collection date.

## Gauteng

A total of 170 laboratory-confirmed measles cases have been reported from epidemiological week 40, 2022 to week 19, 2023 in Gauteng Province displayed in Figure 8. An outbreak was declared on 06 December 2022 (epidemiological week 49, 2022) after three laboratory-confirmed measles cases were reported at a single health facility, Ethafeni Clinic in the City of Ekurhuleni Metropolitan Municipality. To date, the majority of cases, 109, have been reported from the City of Ekurhuleni, 34 from the City of Tshwane, 17 cases from the City of Johannesburg, and ten cases from West Rand. Amongst these 170 cases, 144 have unknown vaccination status while 13 cases were vaccinated (Table 3). Of the 109 cases in Ekurhuleni, 14 were identified at Daveyton's main clinic in Ekurhuleni.




Figure 8. The epidemiological curve of the number of laboratory-confirmed measles cases in districts of Gauteng Province from epidemiological week 40, 2022 to week 19, 2023 by specimen collection dates.

## Free State

There are currently 30 laboratory-confirmed measles cases in this province since epidemiological week 40, 2022 (Figure 9). An outbreak was declared on 20 December 2022 (epidemiological week 51, 2022) in Free State province after three laboratory-confirmed measles cases were reported in the Thabo Mofutsanyana district. Of the 30 cases, 20 have been reported from the Thabo Mofutsanyana district, seven from the Fezile Dabi district, two cases from Xhariep district and one case from the Lejweleputswa district. Of these 20 cases reported from Thabo Mafutsanyana district, five were reported by Bethlehem clinic. The vaccination status of 18 cases is unknown, whereas three cases were not vaccinated, and nine were (Table 3).




Figure 9. The epidemiological curve of the number of laboratory-confirmed measles cases in districts of Free State Province from epidemiological week 40, 2022 to week 19, 2023 by specimen collection dates.

## Conclusion

Overall, the incidence of measles appears to be decreasing across the country, with reproduction numbers for all provinces  $<1$ . However, Limpopo province is still contributing to cases seen in the previous week. Continuous surveillance for measles cases is recommended. Prevention and control of measles outbreaks can only be achieved through vaccination. The national measles vaccination coverage remains low. It is never too late to vaccinate – children over the age of 6 months to 15 years were targeted in the National supplemental immunization campaign rolled out in all provinces on 06 Feb 2023. The NICD continues to report on a large number of cases

with unknown vaccination status. We urge the district and province to complete vaccine status on the investigation forms for completeness of data. Clinicians across the country are urged to be on the lookout for measles cases. For more information about measles, case definition, notification, investigation and guidelines for measles management including vaccination, please refer to our website: <https://www.nicd.ac.za/diseases-a-z-index/measles/>. Health care workers are encouraged to submit reports on any adverse events following immunization (AEFI) through the Med Safety application (<https://medsafety.sahpra.org.za/>) or through submitting a case reporting form to their district surveillance officer.