



Wastewater-based genomic epidemiology for SARS-CoV-2 surveillance in South Africa

24 November 2023

Sample collection dates up to 23 November 2023
(Epidemiological week 47)

F. Els*, N.S Msomi*, N.L Ndlovu, V. Mabasa, E. Phalane, M.E Macheke, S. Gwala, T. Mangena, L. Monametsi, L.P Rabotapi, S. Maposa, M. Yousif#, K. McCarthy#

*Joint first authors
#Joint last authors

Centre for Vaccines and Immunology,
National Institute for Communicable Diseases,
A division of the National Health Laboratory Service

Summary: SARS-CoV-2 transmission and genomics based on evaluation of wastewater at sentinel sites across RSA

Wastewater levels

Epidemiological weeks 33-47

- From weeks 33-47 the cumulative SARS-CoV-2 levels measured at wastewater treatment works (WWTW) has remained around 2 log (100) genome copies/ml of wastewater. This has followed on from the increases observed in weeks 31-33, when the cumulative SARS-CoV-2 levels in wastewater at sentinel sites in South Africa showed increases to levels above 2 log (100) genome copies/ml of wastewater, up from below one log copy/ml in epidemiological week 22 (first week in June 2023). From week 45, levels measured below 2 log genome copies/ml of wastewater.
- In weeks 33-47 increases and/or higher levels have been seen in Gauteng (Northern WWTW), Eastern Cape (Kwanobuhle WWTW), Cape Town (Zandvleit WWTW) and Free State (Bloemspruit WWTW and Sterkwater WWTW).
- Correlation with syndromic surveillance for influenza-like illness (ILI) and severe acute respiratory infection findings (SARI) is required to determine the clinical and public health significance of ongoing transmission.
<https://www.nicd.ac.za/diseases-a-z-index/disease-index-covid-19/surveillance-reports/weekly-respiratory-pathogens-surveillance-report-week/>

Wastewater genomics

Epidemiological weeks 33-41

- No new genomics results were obtained for week 41
- Genomics results were obtained for weeks 33-39 for the heat map and mutational profile.
- **Omicron lineage BA.2.86 followed by JB.2, XBB.1.4, XBB.1.5.81 and XBB sub-lineages** were the dominant lineages circulating in wastewater samples between August and September 2023
- In clinical samples, **BA.2.86** was also the dominant lineage circulating between August and September 2023, followed by **XBB.1.5** and **XBB.1.5.81**.
- The **Omicron lineage BA.2.86** is circulating in KwaZulu-Natal in eThekweni (in the catchments of Northern and Central WWTWs), and in Gauteng, in the City of Johannesburg (in catchments of Northern and Goudkoppies WWTWs), in Ekurhuleni (in the catchments of Olifantsfontein, Vlakplaats, and Hartebeesfontein WWTWs), and the City of Tshwane (in the catchment of Daspoort WWTP). It is also circulating in Eastern Cape in Buffalo City (in the catchment of Mdantsane WWTW), in Western Cape, in the City of Cape Town (in the catchment of Borches Quarry WWTW), and Free State, in Mangaung (catchments Bloemspruit and Sterkwater WWTWs).

Interpretation: Ongoing transmission of SARS-CoV-2 due to Omicron lineages including the new lineage BA.2.86.

Wastewater-based Epidemiology for COVID-19


How is wastewater tested for SARS-CoV-2?

For a full description of this process, see the photo essay developed in collaboration with the Gauteng City Region Observatory <https://www.gcro.ac.za/outputs/photo-essays/detail/photo-essay-sewersheds-what-can-wastewater-tell-us-about-community-health/>

For a technical description and analysis of wastewater levels and results see

<https://pubmed.ncbi.nlm.nih.gov/37506905/>
<https://www.medrxiv.org/content/10.1101/2022.12.15.22283506v1> (accepted by Nature Communications, publication pending)

SARS-CoV-2 is not transmitted by faeco-oral route. Wastewater with SARS-CoV-2 is not infectious



Wastewater-based Epidemiology for COVID-19

What does wastewater testing for SARS-CoV-2 mean?

Left vertical axis:

Number of lab confirmed cases in *the metro or district where the water treatment plant is located*

Bars:

Number of lab confirmed clinical cases in specimens submitted to NICD from persons in the metro/subdistrict where the plant is located

Horizontal axis:

Epidemiological weeks from 2021 to 2023

Coloured lines:

Changes in wastewater SARS-CoV-2 results over time for different treatment facilities

Coloured squares:


Epi weeks during which samples were collected

Facets:


Indicates the platform used to test samples

Right vertical axis:

Log (ie 10 to the power x) copies of SARS-CoV-2 genome per ml of wastewater. So $\log 2=10^2=100$ copies per millilitre, $\log 3=10^3=1000$ copies per millilitre



South Africa at a glance: Summed total of clinical and genome copies



★ Chloroform start and end date


■ Sample Collection ● South African SARS-CoV-2 Wastewater Levels

Changes in levels of SARS-Cov-2 (line graph) in in-flowing untreated wastewater from plants tested by NICD, compared with laboratory-confirmed cases from Tshwane, Johannesburg, Ekurhuleni, eThekweni, Mangaung, Nelson Mandela, Buffalo City, and City of Cape Town (grey bars), by epidemiological week, 2021-2023.

South Africa at a glance: Circulating variants as determined by Freyja deconvolution of sequence data

- SARS-CoV-2 variants in wastewater as determined by the 'Freyja' tool (Scripps Institute)
 - Allows determination of variants in each wastewater sample
- Results from sequencing data ending in epi week 39 (29 September 2023)
 - Omicron lineages XBB.1.5 and Omicron were circulating from August to September, with XBB* dominance in September.



Inferred variants in wastewater samples from South African wastewater treatment plants by month, between April 2021- September 2023



South Africa at a glance: Circulating lineages as determined by Freyja deconvolution of sequence data

- Results from sequencing data ending in epi week 39 (29 September 2023)

- Omicron lineages XBB.1.41 and BA.2.86 were circulating from August to September with BA.2.86 dominance in September.
- Omicron XBB sub-lineages were in circulation throughout August.
- The predominant lineages circulating in clinical samples in the recent week are BA.2.86 followed by XBB.1.5, XBB.1.9 and XBB sub-lineages.



Lineages detected by Freyja in weeks 33-39:

- XBB.1.41
- XBB.1.41.1
- XBB.1.41
- XBB.1.5.28
- FY.5
- XBB.1.5.81
- BA.2.86

What mutations in the spike protein tell us about the circulating lineages

Summary:

Summary of lineages associated with most common mutations associated

Epi Week:


Epidemiological week during which samples were collected

Site Names:

Sites from which wastewater samples were collected

Reference lineages:

Reference lineages with signature mutations with which wastewater samples are compared for lineage determination




Amino acid mutations and frequency – Spike protein

XBB* is a recombinant of BA.2.10.1 and BA.2.75 that is characterised by one or more of the following mutations in the spike protein: V83A, Y144-, H146Q, Q183E, V213E, G252V, G339H, R346T, L368I, V445P, G446S, N460K, F486S, F490S

List of variants and sub-lineages of interest and concern
<https://www.who.int/en/activities/tracking-SARS-CoV-2-variants>

BA.2.86 is a highly mutated sub-lineage of BA.2, recently circulating in Denmark, Israel and the United States of America and is characterised by one or more of the following mutations in the spike protein: R21T, S50L, H69-, V70-, V127F, F157S, R158G, N211-, L212I, V213G, L216F, H245N, A264D, I332V, K356T, R403K, V445H, N450D, L452W, N481K, V483-, E484K, E554K, A570V, P621S, 1670V, P681R, S939F, P1143L, Ins16:MPLF*

V213E, R346T
 Mutations in spike protein associated with XBB* sub-lineages




1149
1135
1121
1107
1093
1079
1065
1051
1037
1023
1009
995
981
967
953
939
925
911
897
883
869
855
841
827
813
799
785
771
757
743
729
715
701
687
673
659
645
631
617
603
589
575
561
547
533
519
505
491
477
463
449
435
421
407
393
379
365
351
337
323
309
295
281
267
253
239
225
211
197
183
169
155
141
127
113
99
85
71
57
43
29
15
1

S939F
 Spike protein mutation associated with the BA.2.86 lineage

P143L
 Spike protein mutation associated with the BA.2.86 lineage

Heatmap showing patterns of emerging mutations in the spike region of SARS-CoV-2, collected from April, 2021 - August, 2023. Mutations appearing in yellow have a low read frequency, those appearing in orange have a medium read frequency and those appearing in red have a high read frequency. Mutations are included and updated weekly.

Gauteng - Tshwane



★ Chloroform start and end date

■ Rooiwal Sample Collection ■ Daspoort Sample Collection ◆ Daspoort Wastewater Treatment Works ◆ Rooiwal Wastewater Treatment Works

SARS-CoV-2 levels and Genomic Results in Epi week 47:

- In Epi week 47, SARS-CoV-2 levels in Daspoort WWTW decreased and are low.
- SARS-CoV-2 levels in Rooiwal WWTW have increased slightly in Epi week 44. Levels remain low. No new results for Epi week 47 are available.


*** Sequencing data ending in Epi week 38 in Rooiwal and 39 in Daspoort. No new sequencing data available.**

- Omicron lineages XBB.1.5.49, XBB.1.16, XBB.1.9.1 and JB.2, were circulating in Daspoort during Epi week 39, with BA.2.86 dominating.
- BA.2.86 was also the dominant lineage detected in Rooiwal during epiweek 38. Other lineages including JB.2, XBB.3 and FY.5 were also in circulation.

SNP Analysis:

- A combination of mutations (V127F, L212I, V213G, L216F, H245N, A264D, I332V, K356T) associated with lineage BA.2.86 were found in both Daspoort and Rooiwal.


Daspoort Wastewater Treatment Works



Legend for Daspoort Wastewater Treatment Works:

- Alpha: BA.1, BA.2, BA.2.1, BA.2.10, BA.2.74, BA.2.86, BA.4, BA.4.6, BA.5
- BA.5.1, BA.5.3, BE.1.1, BE.1.2, Beta, BF.1*, BQ.1, BA.2.86.1, BA.4, BA.4.6, BA.5
- FL.25, FY.1, FY.3, FY.5, HS.1, JB.2, Kappa, Other, XAS, XBB, XBB.1.16
- XBB.1.16.17, XBB.1.22, XBB.1.41.1, XBB.1.42.1, XBB.1.5, XBB.1.5*, XBB.1.5.15, XBB.1.5.17, XBB.1.5.25, XBB.1.5.28, XBB.1.5.45
- XBB.1.5.49, XBB.1.5.7, XBB.1.5.81, XBB.1.5.91, XBB.1.9, XBB.1.9.1, XBB.2.7, XBE


Rooiwal Wastewater Treatment Works



Legend for Rooiwal Wastewater Treatment Works:

- Alpha: BA.1, BA.2, BA.2.86, BA.4, BA.4.6, BA.5
- BE.1.2, BE.6, BE.7, BE.8, BE.9, Beta, BF.7, BQ.1, Delta, FL.25
- FY.5, HS.1, JB.2, Kappa, Other, XAA, XAH, XBB, XBB.1.16, XBB.1.28
- XBB.1.28.1, XBB.1.41, XBB.1.41.1, XBB.1.5, XBB.1.5*, XBB.1.5.28, XBB.1.5.40, XBB.1.5.45, XBB.1.5.81, XBB.1.5.83
- XBB.1.5.91, XBB.1.9, XBB.1.9.1, XBB.3, XBB.6, XBE

Gauteng - Johannesburg



SARS-CoV-2 levels and Genomic Results in Epi week 47:

- As of Epi week 45, SARS-CoV-2 levels in Goudkoppies WWTW have decreased. Levels remain are low. No new results for Epi week 47 are available.
- In Northern WWTW, SARS-CoV-2 levels decreased slightly, but remain moderate in Epi week 45. No new results for Epi week 47 are available.


*** Sequencing data ending in Epi week 36 in Goudkoppies and 31 in Northern. No new sequencing data available.**

- During epiweek 36, Omicron lineage BA.2.86 was dominating in epiweek 36. Other lineages in circulation included XBB.1.5.81, XBB.1.41.1, JB.2 as well as XBB.1.16.17
- Omicron lineages XBB.1.5.28, XBB.1.41.1, XBB.2.4, XBB.1.42.1 and XBB.2.9 were circulating during Epi week 31 in Northern Gauteng.


SNP Analysis:

- A combination of mutations (V127F, L212I, V213G, L216F, H245N, A264D, I332V, K356T) associated with lineage BA.2.86 were found in Goudkoppies.
- SNP analysis could not be performed as the SARS-CoV-2 sequencing coverage in the Northern Johannesburg samples collected during Epi week 34 was too low for meaningful interpretation


Goudkoppies Wastewater Treatment Works



Northern Wastewater Treatment Works



Gauteng - Ekurhuleni



SARS-CoV-2 levels and Genomic Results in Epi week 47:

- The SARS-CoV-2 levels in Hartebeesfontein WWTW increased and levels remain low in Epi week 47.
- As of Epi week 43, there was a slight increase in SARS-CoV-2 levels in Vlakplaats WWTW, after a decrease from Epi week 35. No results for Epi week 47 are available.


** Sequencing data ending in Epi week 38 in Vlakplaats and 39 in Hartebeesfontein. No new sequencing data available.*

- Omicron lineages BA.2.86, JB.2, XBB.1.41.1 and XBB.1.5.81 were circulating in Vlakplaats during epiweek 38
- Lineages JB.2, XBB.1.41.1 and XBB.1.5.81 were circulating during Epi week 39 at the Hartebeesfontein water treatment plant, with BA.2.86 dominating during week 39.



SNP Analysis:

- A combination of mutations (V127F, L212I, V213G, L216F, H245N, A264D, I332V, K356T) associated with lineage BA.2.86 were found in both Ekurhuleni treatment plants.

ERWAT Vlakplaats Wastewater Treatment Works



Hartebeesfontein Wastewater Treatment Works



KwaZulu-Natal - eThekweni

SARS-CoV-2 levels and Genomic Results in Epi week 47:


- SARS-CoV-2 levels in Central WWTW in Epi week 39 showed a sharp increase from low levels in Epi week 38 (1 log copy/ml) to moderate levels (2 log copies/ml), followed by a decrease. Levels remain moderate to low in Epi week 45, and no new results are available for Epi week 47.
- SARS-CoV-2 levels increased from low to moderate in week Epi 36 in Northern WWTW, after which there was a slight decrease. Levels remain moderate in Epi week 45 and no new results are available for Epi week 47.

*** Sequencing data ending in Epi week 39 in Central eThekweni and in eThekweni North. No new sequencing data available.**

- XBB, XBB.1.5.28, JC.1, JB.2 were circulating in Epi week 39, with a BA.2.86 dominance at the Central eThekweni water treatment plant.
- In eThekweni North, XBB.1.42.1, JB.2, and BA.2.86 were circulating during Epi week 39


SNP Analysis:

- A combination of mutations (V127F, L212I, V213G, L216F, H245N, A264D, I332V, K356T) associated with lineage BA.2.86 were found in both eThekweni wastewater treatment plants.




★ Chloroform start and end date ◆ Central Sample Collection ■ Central Wastewater Treatment Works (KZN) ◆ Northern Sample Collection ■ Northern Wastewater Treatment Works (KZN)

Central Wastewater Treatment Works




■ BA.1	■ BA.2	■ BA.2.1	■ BA.2.37	■ BA.2.75	■ BA.2.86	■ BA.4	■ BA.4.6	■ BA.4.8	■ BA.5
■ BA.5.2	■ BA.5.3.5	■ BE.1	■ BE.1.1	■ BE.1.2	■ BE.6	■ BE.7	■ Beta	■ BF*	■ BF.7
■ BN.1	■ BQ.1	■ CH.1.1	■ Delta	■ DR.1	■ EG.2	■ FL*	■ FL.2.4	■ FY.5	■ HS.1
■ JB.2	■ JC.1	■ Other	■ XAK	■ XAY	■ XBB	■ XBB.1.16	■ XBB.1.16.2	■ XBB.1.22.2	■ XBB.1.41
■ XBB.1.41.1	■ XBB.1.5	■ XBB.1.5*	■ XBB.1.5.28	■ XBB.1.5.84	■ XBB.1.5.91	■ XBB.1.9	■ XBB.1.9.1	■ XBB.2	■ XBE

Northern Wastewater Treatment Works



■ BA.1	■ BA.2	■ BA.2.16	■ BA.2.74	■ BA.2.86	■ BA.4	■ BA.5
■ BE.1.1	■ BQ.1	■ CH.1.1.19	■ CH.1.1.20	■ CJ.1	■ Delta	■ FL*
■ FL.25	■ HK.2	■ HS.1	■ JB.2	■ JB.2.1	■ Other	■ XAS
■ XBB	■ XBB.1.16	■ XBB.1.16.17	■ XBB.1.22.2	■ XBB.1.41	■ XBB.1.41.1	■ XBB.1.42.1
■ XBB.1.5	■ XBB.1.5*	■ XBB.1.5.28	■ XBB.1.5.81	■ XBB.1.5.91	■ XBB.1.9	■ XBB.1.9.1

Eastern Cape – Nelson Mandela



★ Chloroform start and end date


■ Brickfield Sample Collection ◆ Brickfield Pre-treatment Works ■ Kwanobuhle Sample Collection ◆ Kwanobuhle Wastewater Treatment Works

SARS-CoV-2 levels and Genomic Results in Epi week 47:


- No new results were obtained in Epi week 47, however, in Epi week 37, a sharp decrease in SARS-CoV-2 levels were seen in Kwanobuhle WWTW, followed by a slight increase. SARS-CoV-2 levels remain low.
- SARS-CoV-2 levels increased from low to moderate from Epi week 33 to Epi week 34 in Brickfield Pre-treatment works. No new results are available for Epi week 47.

*** Sequencing data ending in Epi week 29 in Brickfield and 32 in Kwanobuhle. No new sequencing data available.**

Brickfield Pre-treatment works



Kwanobuhle Wastewater Treatment Works




- SARS-CoV-2 sequencing coverage in the Brickfield samples collected during Epi weeks 30-32 are too low for meaningful interpretation
- Omicron lineages XBB.1.5.81, XBB.1.5.32 and XBB.1.5.27 were circulating in Kwanobuhle during epi week 32.

SNP Analysis:

- SNP analysis could not be performed as the SARS-CoV-2 sequencing coverage in the Brickfield and Kwanobuhle samples collected during Epi weeks 30-39 were too low for meaningful interpretation.

Eastern Cape – Buffalo City



SARS-CoV-2 levels and Genomic Results in Epi week 47:

- In Epi week 47, SARS-CoV-2 levels in Mdantsane WWTW decreased and levels remain low.
- SARS-CoV-2 levels in East Bank WWTW in Epi week 47 remain low after decrease was observed in Epi week 38.


** Sequencing data ending in Epi week 38 in Eastbank and 39 in Mdantsane. No new sequencing data available.*

- Omicron lineage BA.2.86, XBB.2.9, XBB.2.9 and GG.1 were circulating in Eastbank during Epi week 38.
- Lineages CH.1.1.11, BA.2.86, XBB.1.5.81 and XBB.1.5.88 were circulating in Mdantsane during Epi week 39.

SNP Analysis:


- A combination of mutations (V127F, L212I, V213G, L216F, H245N, A264D, I332V, K356T) associated with lineage BA.2.86 were found in both Eastbank and Mdantsane.

East Bank Wastewater Treatment Works




- group
- A
- Alpha
- BA.2
- BA.2.86
- BA.4
- BA.5
- BE.1.2
- Delta
- GG.1
- Other
- XBB.2.4
- XBB.2.9

Mdantsane Wastewater Treatment Works



- group
- BA.1
- BA.2
- BA.2.86
- BA.2.86.1
- BA.3
- BA.4
- BA.4.6
- BA.5
- BE.1
- BE.1.2
- BE.9
- CH.1.1.11
- CJ.1
- Delta
- FY.3
- FY.5
- HS.1
- JB.2
- JB.2.1
- Other
- XAP
- XBB.1.22
- XBB.1.41.1
- XBB.1.5.32
- XBB.1.5.56
- XBB.1.5.81
- XBB.1.5.88
- XBB.1.5.91

Free State – Mangaung



SARS-CoV-2 levels and Genomic Results in Epi week 47:

- In Bloemspruit WWTW, a 2-fold increase in SARS-CoV-2 levels were seen in Epi week 44. Levels are moderate. No new results are available for Epi week 47.
- A sharp increase in SARS-CoV-2 levels were seen in Sterkwater WWTW from Epi week 38. In Epi week 42 levels decreased and increased again in Epi week 43 and remain moderate. No new results for Epi week 47 are available.


*** Sequencing data ending in Epi week 39 in Bloemspruit and 38 in Sterkwater. No new sequencing data available.**

- Lineage JB.2 was the dominant lineage circulating in Bloemspruit during week 39, with BA.2.86, FL.25, FY.5 and XBB.2.4 also circulating in the area.
- XBB.1.41.1 was the dominant lineage circulating in Sterkwater during week 38. HS.1, JB.2, XBB.1.22 and JB.2.1 were also circulating during this time.


SNP Analysis:

- A combination of mutations (V127F, L212I, V213G, L216F, H245N, A264D, I332V, K356T) associated with lineage BA.2.86 were found in both Bloemspruit and Sterkwater.

Bloemspruit Wastewater Treatment Works



Sterkwater Wastewater Treatment Works




group

Alpha	BA.1	BA.2	BA.2.19	BA.2.86	BA.4	BA.5	BA.5.1
BA.5.2	BE.1.1	BE.1.2	Beta	BQ.1	Delta	EG.2	EG.5.1.1
FL*	FL.12	FL.25	FL.4.8	FY.5	HK.3.1	JB.2	Other
XBB	XBB.1.16.17	XBB.1.22	XBB.1.41	XBB.1.41.1	XBB.1.5	XBB.1.5*	XBB.1.5.17
XBB.1.5.28	XBB.1.5.81	XBB.1.9	XBB.1.9.1	XBB.2.4	XBE		

group

BA.1	BA.2	BA.4	BA.5	BA.5.1	BA.5.3	BE.1.1
BE.1.2	BE.7	Beta	BF.7	BQ.1	Delta	HS.1
JB.2	JB.2.1	JC.1	Other	XAS	XAV	XBB
XBB.1	XBB.1.22	XBB.1.41.1	XBB.1.5	XBB.1.5*	XBB.1.5.17	XBB.1.5.25
XBB.1.5.81	XBB.1.5.83	XBB.1.9	XBB.1.9.1	XBB.2	XBE	

Western Cape – City of Cape Town




SARS-CoV-2 levels and Genomic Results in Epi week 47:


- After a sharp increase in SARS-CoV-2 levels in Epi week 37, SARS-CoV-2 levels in Borcheds Quarry WWTW remain moderate. No new results are available in Epi week 47.
- In Epi week 41, a 2-fold increase in SARS-CoV-2 levels were observed in Zandvleit WWTW. In Epi week 43, SARS-CoV-2 levels remain moderate. No new results are available in Epi week 47.

*** Sequencing data ending in Epi week 33 in Bloemspruit and in Sterkwater. No new sequencing data available.**

Borcheds Quarry Wastewater Treatment Works



Zandvleit Wastewater Treatment Works



- During epiweek 33, lineages XBB.1.5.81, XBB.1.5.28, JB.2, and XBB.1.16 were detected in Borcheds Quarry.
- Omicron lineage XBB.1.5.28, XBB.1.5.1, XBB.1.22, JB.2.1 and XBB.1.16.26 were circulating in Zandvleit in week 33.

SNP Analysis:

- A combination of mutations (V127F, L212I, V213G, L216F, H245N, A264D, I332V, K356T) associated with lineage BA.2.86 were found in both Borcheds Quarry and Zandvleit.

COLLABORATORS



FUNDERS



TEAM

